

Eidgenössische Technische Hochschule Zürich Swiss Federal Institute of Technology Zurich

## Exercise - Even matrices

You are still part of a team to develop a pseudorandom number generator. Your generator has already passed some simple statistical tests, but now it is time to get serious. You arrange a list of bits produced by the generator in an n by n matrix. If

$$M = \begin{pmatrix} x_{1,1} & x_{1,2} & & x_{1,n} \\ x_{2,1} & x_{2,2} & \cdots & x_{2,n} \\ & \vdots & \ddots & \\ x_{n,1} & x_{n,2} & & x_{n,n} \end{pmatrix}$$

was a truly random matrix of bits, then it would have the property that the sum

$$\sum_{i'=i_1}^{i_2} \sum_{j'=j_1}^{j_2} x_{i',j'}$$

is even for about half of the quadruples  $(i_1,i_2,j_1,j_2)$ ,  $1 \leqslant i_1 \leqslant i_2 \leqslant n$  and  $1 \leqslant j_1 \leqslant j_2 \leqslant n$ .

To check whether this is the case for your generator, you need to be able to count the number of such quadruples.

**Input** The first line of the input contains the number  $t \le 15$  of test cases. Each of the t test cases is described as follows.

- It starts with a line that contains an integer n such that  $1 \le n \le 200$ .
- This is followed by n lines, where the i-th line contains the n integers  $x_{i,1}$  ...  $x_{i,n}$ , separated by a space, such that  $x_{i,j} \in \{0,1\}$ , for all  $i \in \{1,\ldots,n\}$  and  $j \in \{1,\ldots,n\}$ .

Output For each test case output a single line that contains the number of quadruples  $(i_1,i_2,j_1,j_2)$  where  $1 \le i_1 \le i_2 \le n$  and  $1 \le j_1 \le j_2 \le n$  and for which the sum

$$\sum_{i'=i_1}^{i_2} \sum_{j'=j_1}^{j_2} x_{i',j'}$$

is even.

**Points** There are three groups of test sets, worth 100 points in total.

- 1. For the first group of test sets, worth 30 points, you may assume that  $1 \le n \le 10$ .
- 2. For the second group of test sets, worth 50 points, you may assume that  $1 \le n \le 50$ .
- 3. For the third group of test set, worth 20 points, there are no additional assumptions.

Corresponding sample test sets are contained in testi.in/out for  $i \in \{1, 2, 3\}$ .

## Sample Input

## 3 2 1 1 1 1 3 1 0 1 0 1 0 0 0 1 4 1 1 0 0 0 0 1 1 1 0 1 0 0 1 0 1

## Sample Output

5 15 52