



BIOLOGY TWO

	AB	Ab	aB	ab
AB	AABB 	AABb 	AaBB 	AaBb
Ab	AABb 	AAbb 	AaBb 	Aabb
aB	AaBB 	AaBb 	aaBB 	aaBb
ab	AaBb 	Aabb 	aaBb 	aabb

Heredity





Genetics



Evolution

BIOLOGY


~ SPRING 2022 ~



	Lesson	Topic	Page(s)
Unit 1 Heredity	1	What is a species?	6-7
	2	Life finds a way	8-9
	3	Activity: Vegetative propagation with potato	10-15
	4	Mendel's famous experiment	16-17
	5	Inheritance explained	18-21
	6	Deep dive: Why aren't Zebras domesticated?	
	7	The laws of heredity 1	22-25
	8	The laws of heredity 2	26-29
	9	Deep dive: Punnett Square Practice	
	10	Pet pedigree puzzle	30-33
	11	Bigger Punnett squares	34-37
	12	Heredity Quiz Show	38-41
Unit 2 Genetics	13	What is a gene?	42-44
	14	Chromosomes	45-47
	15	Deep dive: Fingerprints	
	16	Meiosis and mistakes	48-49
	17	The Genetic Code 1	50-51
	18	Activity: Gummy worm karyotypes	52-55
	19	The Genetic Code 2	56-57
	20	Blood types explained	58-60
	21	Activity: What's your blood type?	61-63
	22	Mutations	64-65
	23	Nature and nurture	66-67
	24	Deep dive: Why Red Delicious are the least delicious apples	68-69
	25	Calico Cats & X-linked traits	70-71
	26	Modifying genes and gene therapy	72-73
	27	Genetics Quiz Show	74-77

The course is structured with at least two videos per week. The third activity of each week is a deep dive activity or a quiz show. Students who complete and submit all activities and assignments will receive a bonus certificate of completion.

	Lesson	Topic	Page(s)
Unit 3 Evolution	28	Endangered vs invasive species	78-79
	29	Darwin and the Galapagos	80
	30	Activity: Journey of the HMS Beagle	81-83
	31	Survival of the fittest	84-86
	32	Selection and Adaptation	87-88
	33	Deep dive: When Sherpas climb Mt. Everest	
	34	Evolution: what it is and what it isn't	89-90
	35	Genetic drift	91-92
	36	Activity: Drifting marbles and population models	93-94
	37	Species and gene pools	95-96
	38	Taxonomy and similar traits	97-99
	39	Deep dive: DNA and Pandas	
	40	The fossil record	100-101
	41	Phylogenies and family trees	102-103
	42	Activity: Build your own phylogeny	104
	43	Australia vs New Zealand	105
	44	Timeline of life on Earth	106-107
	45	Evolution Quiz Show	108-111
		Appendix	112-136



How to use this course:

For younger students (2nd-5th grade) we recommend a “get the basics” approach that focuses on the Mon/Wed classes. For older or more advanced students (6th-8th grade) we recommend completing each of the Friday activities and assignments to reinforce the concepts covered.

Option One: Get the Basics

Ideal for younger students

Participate in Mon-Wed classes only. Friday activities, articles, and other assignments are optional. They can be skipped, saved for later, or completed depending on your preference.

Tips for best learning:

- Read the pages that go with each lesson before watching the video. Take 10-15 minutes before class to see if you can fill in the blanks. If you can't fill in everything, that's okay! Listen during class to see if you can complete the page.
- If a lesson moved too fast, rewatch it later to help learn the concepts.
- Download the answer key for the notes, but don't look at the answers until after you give things a try yourself!

Option Two: The Full Course

Ideal for older students & advanced learners

Participate in Mon-Wed classes and do the Friday activities and reading assignments. Complete all quizzes and assignments to receive a certificate of completion at the end of the course.

Project Supply List:

Lesson 3 - Vegetative Propagation with Potato

- Potato or sweet potato (for quicker results, use one where the eyes have already begun to sprout)
- Knife
- 4 large pots or burlap sacks
- Potting soil

Lesson 18 - Gummy Worm Karyotypes

- Enough gummy worms or other oblong-shaped colorful candy to create 16 pairs of chromosomes
- Chocolate (to melt and cover some of the gummy worms.)
- Knife or scissors

Lesson 21 - What's the Blood Type?

- Home blood typing kit (optional)

Lesson 24 - Apple Variety Taste Test

- Several varieties of apple including Red Delicious, Granny Smith, and three other types.
- Knife

Lesson 30 - Journey of HMS Beagle

- Printable world map
- Paper and scissors
- Crayons, colored pencils, or markers

Lesson 36 - Model a Gene in a Population

- 20 black marbles and 20 white marbles or ANY other items that can be drawn randomly in a sample! For example, beans, candy, coins, or dice marked with tape.

Lesson 42 - Build your own phylogeny

- Paper and art supplies

Lesson 45 - History of Earth Timeline

- Paper, tape, and crayons, markers, or colored pencils

Which science standards are covered in this course?

The standards listed below are the national science standards for K-12 education in the United States. They are commonly referred to as Next Generation Science Standards or NGSS.

MS-LS1-4. Use argument based on empirical evidence and scientific reasoning to support an explanation for how characteristic animal behaviors and specialized plant structures affect the probability of successful reproduction of animals and plants respectively. **Lessons 28, 29, 31, 32**

MS-LS1-5. Construct a scientific explanation based on evidence for how environmental and genetic factors influence the growth of organisms. **Lesson 23**

MS-LS3-1. Develop and use a model to describe why structural changes to genes (mutations) located on chromosomes may affect proteins and may result in harmful, beneficial, or neutral effects to the structure and function of the organism. **Lessons 13, 14, 17, 19, 22, 28**

MS-LS3-2. Develop and use a model to describe why asexual reproduction results in offspring with identical genetic information and sexual reproduction results in offspring with genetic variation. **Lessons 2, 4, 5, 7, 10**

MS-LS4-1. Analyze and interpret data for patterns in the fossil record that document the existence, diversity, extinction, and change of life forms throughout the history of life on Earth under the assumption that natural laws operate today as in the past. **Lessons 38, 40, 41, 43**

MS-LS4-2. Apply scientific ideas to construct an explanation for the anatomical similarities and differences among modern organisms and between modern and fossil organisms to infer evolutionary relationships. **Lessons 38, 41**

MS-LS4-3. Analyze displays of pictorial data to compare patterns of similarities in the embryological development across multiple species to identify relationships not evident in the fully formed anatomy. **Lesson 38**

MS-LS4-4. Construct an explanation based on evidence that describes how genetic variations of traits in a population increase some individuals' probability of surviving and reproducing in a specific environment. **Lessons 31, 32, 34**


MS-LS4-5. Gather and synthesize information about the technologies that have changed the way humans influence the inheritance of desired traits in organisms. **Lesson 26**

MS-LS4-6. Use mathematical representations to support explanations of how natural selection may lead to increases and decreases of specific traits in populations over time. **Lessons 35, 36**

Lesson	Topic	NGSS	Page(s)
2	Life finds a way	MS-LS3-2	8-9
4	Mendel's Famous Experiment	MS-LS3-2	16-17
7	The laws of heredity 1	MS-LS3-2	22-25
8	The laws of heredity 2	MS-LS3-2	26-29
10	Pet Pedigree Puzzle	MS-LS3-2	30-33
13	What is a gene?	MS-LS3-1	42-44
14	Chromosomes	MS-LS3-1	45-47
17	The genetic code 1	MS-LS3-1	50-51
19	The genetic code 2	MS-LS3-1	56-57
22	Mutations	MS-LS3-1	64-65
23	Nature and nurture	MS-LS1-5,	66-67
26	Modifying genes and gene therapy	MS-LS4-5	72-73
28	Endangered vs invasive species	MS-LS1-4, MS-LS3-1	78-79
29	Darwin and the Galapagos	MS-LS1-4	80
31	Survival of the fittest	MS-LS1-4, MS-LS4-4	84-86
32	Selection and adaptations	MS-LS1-4, MS-LS4-4	87-88
36	Drifting marbles and population models	MS-LS4-6	93-94
37	Species and gene pools	MS-LS4-2, MS-LS4-3 MS-LS4-6	95-96
38	Taxonomy and similar traits	MS-LS4-4	97-99
40	The fossil record	MS-LS4-1, MS-LS4-2, MS-LS4-3	100-101
41	Phylogenies and family trees	MS-LS4-1	102-103
44	Timeline of life on Earth	MS-LS4-1	106-107

What makes a species?

The domestic dog includes hundreds of different breeds with diverse personalities, shapes, and sizes. The biggest dog, an English Mastiff named Zorba, weighed more than 143 kg (315 pounds) and measured 94 cm (37 inches) tall from ground to shoulder. The smallest dog, a Chihuahua named Milly, is just under 10 cm (4 inches) tall and weighs just 0.5 kg (about 1 pound). Mastiffs, Chihuahuas, Dachshunds, and Greyhounds are all called dogs, but they are incredibly different animals!



1. What makes a dog a dog? Share your opinion:

Answers will vary. Some possible characteristics to include could be that dogs are four legged, have fur, are carnivorous, and are social animals.

2. If the characteristics you described above were used as the official definition of a dog, would cats also be called dogs?



Answers will vary.

3. List three similarities and differences between cats (*Felis*) and dogs (*Canis*):

Similarities between cats and dogs:	
	<ul style="list-style-type: none"> Quadrupeds (4 legs) Have fur Both kept as pets by humans Both have whiskers Both can be trained. Both have tails. Both have breeds with distinct traits. Both have wild relatives (wolf, coyote, lynx, bobcat)

Differences between cats and dogs:


- Dogs are pack animals, while cats are solitary hunters.
- Dogs are diurnal (active during the day) while cats are primarily nocturnal
- Dogs bark/howl (can't purr) while cats purr and meow (can't bark).
- Cats have retractable claws, but dogs don't.
- Cats climb trees. Dogs can't.
- Dogs are omnivores and scavengers. Cats are obligate predators.
- Cats have 30 teeth, dogs have 42
- Cats can't taste sugar, dogs can.




4. Coyotes, wolves, and dogs can all interbreed, creating hybrids like the wolfdog, coywolf, coydog or dogote. Lions and tigers can also be bred to create hybrids, which are called ligers or tigons. In your opinion, would it be possible to cross a cheetah with a wolf? If so, what would you call this hybrid? A woltah? A cheef?

Answers will vary.

Selective breeding can cause big changes



Breeders prized the "squished face" appearance of the Pug and chose dogs with shorter and shorter snouts. Because of their face shape, modern Pugs are susceptible to eye trouble and breathing problems.



In 1900, Shetland Sheepdogs were reported to weigh between 7 and 10 pounds and have medium length fur. The modern Shetland Sheepdog has doubled in size and has much longer fur.

5. Suppose that a settlement on Mars uses Great Danes for pulling sleds while a different settlement uses Chihuahuas for hunting cockroaches that hide in small spaces. If the people in the first settlement selectively breed their Great Danes to be larger and stronger while the people in the second settlement breed their Chihuahuas to be smaller and faster, what would you expect Mars dogs to look like after approximately 1,000 years have passed? Would Great Danes and Chihuahuas be different species from each other then?

Answers will vary.

Martian Great Dane in 3022



Have fun drawing! Does your Martian Dane look similar to a bear?

Martian Chihuahua in 3022

Have fun drawing! Does your Martian Chihuahua look similar to a weasel?



Life finds a way

The **ability to reproduce** is one of the key characteristics in most definitions of life. There are two main strategies: asexual reproduction, which creates identical clones, and sexual reproduction which creates offspring that are different from each parent.




Record the primary advantages and disadvantages in each category below. Consider the following:


- Speed of reproduction: Is the next generation formed and independently growing within minutes or days? Does it take years to form an independent next generation?
- Genetic diversity: is the diversity high (which provides greater resistance to disease) or low (which makes a population more at risk from disease)?
- Does reproduction require a mate? Could one individual reproduce?




Match each reproductive strategy with the correct definition:




Genetic information from two individuals is combined to create offspring that are genetically different from both parents. This reproductive strategy is common among animals, fungi, and plants.




An organism is split into pieces. Each of those fragments then develops into adult individuals that are clones of the original organism. Molds, lichen, sponges, certain worms, and sea stars all use this reproductive strategy.



The division of one cell into two identical cells. This reproductive strategy is used by many types of single-celled life forms including bacteria and archaea.



A specialized cell is formed that can withstand harsh environmental conditions such as drought or extreme heat. Once conditions improve, this cell germinates and begins growing. This reproductive strategy can be either sexual or asexual. Commonly used by fungi.




A form of reproduction occurring in plants where a fragment or cutting is separated from the parent plant and grows as a separate individual. Commonly used with food crops such as bananas, sweet potato, sugarcane, and pineapple.

Hands-on Science Project

PROPAGATE A POTATO

MATERIALS:



Potato or sweet potato

Knife

Rubbing Alcohol


4 identically-sized pots or bags
Use a large pot or bucket with drainage holes, or a large burlap sack.

Potting soil
Enough to fill all 4 containers


For best results, use sprouting potatoes

Most commercially-available potatoes are treated with chlorpropham, a compound that prevents sprouting. For best results, select a tuber where you can see small sprouts or swelling in the "eyes" of the tuber. If you do not see any sprouts or signs of budding, let the potato sit in a dark location at room temperature until sprouts form.


Potatoes are tubers, specialized root structures that store starch. Tubers contain buds that can grow into new plants!



1. Place the potato in a dark and warm area (such as a kitchen cupboard) and let it sit there until sprouts have formed at the eyes. This may take between 1 to 3 weeks.




2. When the potato has sprouts, clean the knife with rubbing alcohol. Use the clean knife to cut the potato into 4 pieces that are roughly equal in size. Make sure that each piece of potato has at least one or two "eyes" with sprouts. If there are many sprouts on the piece of potato, break off some of the sprouts so there will be no more than two sprouts per quadrant.



3. Let the cut pieces sit at room temperature for 24 hours. This drying process helps prevent mold from growing on the potatoes after they are planted.



4. Fill the pots or burlap bags 1/3 full with potting soil. Place one potato piece in each container so that the potato is approximately 5 cm or 2 inches below the surface of the soil.



5. Water the soil in each pot so that it is moist. Place two of the pots in a very sunny location and the other two pots in a shady location.

Place two pots in a bright location, either under lamps or next to a window that receives several hours of full sunlight.

6. Care for the plants by keeping the soil moist with regular watering. Do not let the soil become too wet or waterlogged, and be careful that it doesn't dry out completely.

Indoors or Outdoors?




This activity can be done either indoors or outdoors. If you have space and it's winter time, we recommend indoors! If you have cats or dogs, take care that they don't eat the above-ground leaves and stems of the potato plant, which are poisonous. If you would prefer to plant your potatoes outdoors, plant them no earlier than 4 to 6 weeks before the average last frost. They are cold hardy and can handle some freezing or near-freezing temperatures. Whether the plants are indoors or outdoors, they will need at least 10 to 12 weeks of growth to produce a crop of new tubers.

7. After sprouting, continue to add additional layers of potting soil until the pot is full. Record your observations about the size and appearance of the above-ground plants at regular intervals.

8. After 10 to 15 weeks or when the plants begin to die back, pour out the potting soil and sort through it to find the new potato tubers. (See page 13 for more detailed instructions).

How many tubers did you find? Which plants produced the most potatoes?

Fill the pots with more soil as the plants grow.



New potatoes!

Observations just after the first sprouts emerge:







Full sun



Partial sun or shade

Noting the date is a vital part of recording research!

DATE:
1/02/2022







Draw the soil line and a simple representation of your tubers in these pots!

Observations will vary! These entries are examples from when we planted potatoes.

Write a line or two about what you observe.

Observations 1 week after the first sprouts emerge. Draw the approximate size and shape of the potato plants growing in each container.

DATE:
1/09/2022




The leaves are starting to curl on the plant in pot A. I think it may be sick.

Observations 3 weeks after the first sprouts emerge:

DATE:

1/23/2022




We added more soil at week 1. Plant A got sick with a fungus and died on 1/16/2022.

Observations 6 weeks after the first sprouts emerge:

DATE:

2/13/2022




The full sun plant looks much more robust and larger than the part shade plants.

Observations 10 weeks after the first sprouts emerge:

DATE:

3/6/2022



Leaves are yellowing - The plants seem to be getting ready to die back.

DATE:

3/10/2022

POTATO HARVEST

Before harvesting your potatoes, answer the following questions:

1. Which plants have the most above-ground material? Count the approximate number of leaves on each plant and record their numbers.

The plant with the most above-ground mass: B

The number of leaves on each plant:

NA. Died on week 3 125

A

20

B

35

C

D

2. How different were the light conditions between your "full sun" and "partial sun" plants? If possible, study the locations for one day and count the number of hours of direct light each plant experienced.


Full sun = window with 9 hours of direct light each day

Part shade = window with 2 hours of direct light.


4. Plants get their energy from photosynthesis, which requires light. Which plants were able to do the most photosynthesis during the experiment? What did the plant do with the energy it gathered from photosynthesis?

The full sun plants gathered most energy by photosynthesis. They used some of that energy to make tubers!

Full sun



Partial sun or shade



Draw the above ground portion of your potato plants. Then draw a guess of what you think the root structure will look like. How many tubers do you expect to find?

When to harvest potatoes

Farmers usually wait until the tops of the potato plants have turned yellow and died back before harvesting. This allows the plant to store the most starch in the tubers.

With potatoes grown indoors, you can wait for the plants to go dormant or you can harvest them anytime after 10-12 weeks of growth. The choice is yours. Harvesting before 10 weeks will result in very small (pea and marble size) potatoes.

How to harvest potatoes

If the plants are in containers, take them outside and empty the dirt from the container. Sift through all the dirt and separate the tubers from the roots and potting soil.

If the plants are in the ground, loosen the soil around them with a shovel. Pull up the plants and then thoroughly explore the area with shovel and hands to be sure you found all of the tubers.

Wash and dry the potatoes and store them in the fridge until ready to eat. If the skin of a new potato is scratched or damaged during harvesting, then they won't store for as long and should be eaten sooner rather than later.

POTATO HARVEST

After harvesting your potatoes, answer the following questions:

1. Which plants produced the greatest number of tubers? Count the record the number of tubers from each plant.

AB
17 potatoes
(Most potatoes)C
9 potatoesD
11 potatoes

2. Which plant produced the greatest biomass of tubers? In other words, which plant grew the most food?
If you don't have a scale to weigh the potatoes, that's okay! Compare the size of the potatoes from each plant and draw pictures to represent the potato production of each.

AB
C
D

3. The main variable studied in this experiment was light. Ideally, everything else would have been the same for each potato plant. But in real life, experiments don't always have ideal conditions! Were there any other variables that impacted your plants, such as disease, an injury, waterlogged roots, or drought stress?

Yes. Plant A got sick with a disease and died on week 3. It showed signs of illness just

5 days after sprouting.

CONCLUSIONS


DATE:
3/10/2022

Your potatoes were genetically identical clones. Did they grow into identical plants or did they look different from each other? What impact did light have on the production of your potatoes? Do you have any advice for someone who wants to grow their own potato clones?

They looked very different! The plants with full light produced more potatoes and larger potatoes than the plants with part shade. Only advice is to really carefully wash and clean potatoes before cutting them.

Potato Blight and the Irish Potato Famine

From 1845-1852, Ireland experienced a devastating famine when a fungus-like microorganism called *Phytophthora infestans* caused a serious disease known as potato blight. The blight caused the Irish to lose about $\frac{3}{4}$ of their most important food crop and resulted in over a million people dying of starvation and millions emigrating away from the island in the following years when the potato harvest was poor. Ireland is possibly the only country in the world that has a lower population today (5 million) than in 1840 (about 6.5 million).



How Potatoes are Grown

Potatoes can be infected by many different pathogens including viral, fungal, bacterial, and protozoan! Because they are susceptible to so many diseases, many of which are hard to detect, the crops of modern potato farmers actually start out in test tubes!

Tissue Culture (Also called microppropagation)




Growing cells in an artificial medium like agar or broth.

It's called tissue culture and it all begins with a small piece of plant about the size of a flake of pepper. This flake of green leaf tissue is taken from a parent plant which was grown in a controlled environment free from viruses.

The disease-free bit of the plant (meristem) is placed in a test tube that has all the nutrients that the plant needs in order to grow. The test tube remains in the sterile lab until the cells have grown into a "plantlet," a small plant with distinct roots, stem, and leaves.

When the plantlets are large enough, they are moved to a greenhouse and planted in the ground. Then, just like regular potatoes, they grow for a few months, forming miniature tubers. After harvesting, these small tubers (called seed potatoes) are sorted and stored until it's time to plant the large crop of potatoes in regular fields.

WHAT'S THE DIFFERENCE BETWEEN A COOKED SWEET POTATO AND A FLYING PIG?



ONE IS A HEATED YAM AND THE OTHER IS A YEETED HAM!

Certified seed potatoes have been tested multiple times to be sure they are free from bacterial, fungal, and viral infections.


When the seed potatoes are ready to be planted, a large machine uses sharp spikes or suction cups to grab each seed potato and drop it into the designated furrow of soil.

The buds on the seed potatoes sprout into plants and as the plants grow, so does their network of tubers underground. It is now up to the farmer to care for the potatoes by irrigating and fertilizing them. They'll be ready to harvest between 80 to 115 days after planting. At harvest, they're carefully dug up and stored between 10 and 14 days in cool temperatures ($\approx 15^{\circ}\text{C}$) to allow the skins time to harden and minor injuries to seal. Then they're sent to a grocery store near you!




WHY DOES EVERYONE WANT TO EAT US?

WE MUST BE DELICIOUS!



SEED POTATOES



POTATO SEEDS



PRODUCED FROM FLOWERS THAT GROW INTO GREEN TOMATO-LOOKING BERRIES.

Why is Idaho famous for its potatoes?

The domestic potato originated in the mountains of Peru. Idaho has a similar climate and geology to this area with warm sunny days, cool nights, and fertile volcanic soil that is well-draining and high in nutrients. The high elevation, geologic history, location, and surrounding areas all combine to make Idaho ideal for potato production. The state grows about 13 billion pounds of potatoes every year.


Idaho even has a museum dedicated to the potato with an exhibit of the world's largest potato chip!

The question of inheritance


Throughout history people have had different ideas about how traits are inherited. The three ideas described below were common when Mendel was alive. Which of these ideas best explains what you have observed about traits and how they are inherited? Rate them from one star (not accurate, e.g. there are lots of counter examples for this idea) to four stars (this idea explains inheritance very well).



Lamarckism is the belief that acquired traits can be inherited. Giraffes are able to lengthen their necks by constantly stretching, so their offspring will inherit a longer neck.



Pangenesis is the idea that every part of the body produces small particles called gemmules which are then passed on to offspring.



Blending inheritance is the idea that each trait will be inherited by the offspring at about the average level of each parent. For example, your hair color falls between your parents' hair colors.




Gregor Mendel

Mathematician, biologist, abbot, and founder of genetics

Gregor Mendel was a monk at St. Thomas Abbey. During Mendel's lifetime, the question of how traits were passed from one generation to the next was one of the great mysteries of science. Popular ideas to explain inheritance included Lamarckism, the idea that acquired traits could be passed on to offspring.

In the 1850s, Mendel conducted experiments to discover how inheritance worked. He studied the characteristics of pea plants, such as green and yellow seeds. His experiments were conducted in a small garden next to the abbey.


His conclusions were incredibly advanced for his time and he is widely considered to be the founder of modern genetics.



FILL IN THE BLANKS USING THESE WORDS:



garden inheritance traits pea genetics

A famous experiment



Peas are self-fertile. They do not need pollinators and unless a person intervenes, the pollen on the stamens will fertilize the stigma and produce a pea pod filled with seeds. To cross-pollinate two different pea plants, Mendel used tweezers to remove the stamens from one flower (1) and used a paintbrush to gather pollen from the flower of a different plant (2). The pollen was then applied to the stigma of the first plant to produce hybrid seeds (3).


MEET THE PARENT PEAS!



A pea plant with smooth seeds is cross-fertilized with a plant that has wrinkled seeds.

ALL of the seeds in that first generation (called the F_1 generation) are smooth.

When those first generation seeds (F_1) are self-fertilized, the next generation shows a 3 to 1 ratio of smooth to wrinkled seeds.



We see the same result with flower color.

If you were Mendel, how would you explain these results? Why is one trait disappearing in the F_1 generation and then coming back again in the next?

Answers will vary.

Disappearing and reappearing traits?

Mendel's results explained

FILL IN THE BLANKS USING THESE WORDS:


alleles inherited generation ignored traits dominant

When crossing two true-breeding pea plants with different traits, Mendel found that the F₁, or first generation of plants looked identical to one of the parent plants. The trait from the other parent had "disappeared." But when these plants were self-pollinated, they were no longer true-breeding. Both of the traits appeared with a consistent ratio: 1/4 of the plants had the trait that had disappeared while 3/4 of the plants had the other trait.

To explain this phenomenon, Mendel said that each plant inherited two factors, one from each parent. Some factors were dominant and others were recessive. Today, we call these factors genes or alleles.


He published his research, but it was ignored for more than thirty years. Mendel wasn't recognized for his work until other scientists discovered the same facts in the 1900s.

THE TIME TRAVELING SCIENTIST




Genotype

the genetic information



The gene for flower color.*

In this example, we use "P" to represent the allele for purple flower color and "p" to represent the allele for white flower color.



purple allele

white allele


*We'll learn more about what genes are (segments of DNA) and where they are located (on chromosomes) in the next unit. For now, we can think of them as little bosses telling the organism what to do.




Phenotype

the observed trait


In this example, there are two phenotypes: purple flowers and white flowers. Write the expected phenotype in each box below. Then color the flowers accordingly.



A plant with two alleles for purple flowers will produce:
purple




A plant with two alleles for white flowers will produce:
white




A plant with one allele for purple and one allele for white will produce:
purple


In the example above, the purple allele is **dominant** and the white allele is **recessive**. The traits below also have a dominant allele (represented by a capital letter) and a recessive allele (lower case letter). Use this information to indicate the phenotype that will be produced from each pair of alleles.




FOR PLANT HEIGHT, I HAVE INSTRUCTIONS FOR DWARF PLANTS. WHAT DO YOU THINK ABOUT THAT?




IT'S PERFECT!




The phenotype will be:
tall




THE SEEDS MUST BE SMOOTH!! NO WRINKLES!!




I WAS HOPING FOR WRINKLES, BUT WHATEVER YOU SAY.




The phenotype will be:
smooth



ONLY YELLOW SEEDS!! YES! YELLOW SEEDS FOREVER!!!




GREEN SEEDS




The phenotype will be:
green

The traits Mendel studied follow a **dominance / recessive** pattern. The trait of wool color in sheep does as well, which explains why black sheep are less common than white. Color in the sheep below with the correct phenotype and then label each block below as either "Homozygous dominant," "Homozygous recessive," or "Heterozygous."




Mendel's results explained (in more detail)


Write either "heterozygous" or "homozygous" in each of the labels below.




The plants in the parent generation are true-breeding or homozygous for their trait, which is plant height. The tall plants always produce seeds that grow into tall plants. The dwarf plants always produce seeds that grow into dwarf plants.



The F₁ generation received one allele from each parent plant and are heterozygous for plant height. All of the plants in this generation are tall. The plants will be self-fertilized to form the F₂ generation.




In the F₂ generation, a quarter of the plants show the phenotype of dwarfism. These plants are homozygous recessive. Two of the three tall plants are heterozygous, which means they are no longer true-breeding for the trait of plant height.



In cases of incomplete dominance, the heterozygous phenotype is different than either of the homozygous phenotypes!

Most human traits are controlled by multiple genes and have complicated inheritance patterns. We sometimes think of eye color as being a trait that follows Mendelian inheritance patterns with brown eyes being dominant and blue eyes being recessive.

But it's not that simple! With more than 16 different genes playing a role in the process, two blue-eyed parents can indeed have a brown-eyed child. The same is true with many other traits including hair color, earlobes, and more.



* For further reading, see Genotype-phenotype associations and human eye color by Desiree White in the Journal of Human Genetics. Published Oct 14, 2010.

Draw lines to match each term with the correct description:

Recessive

Only one copy (allele) of a gene is needed for the trait to be expressed. For example, if the trait for large feathers is represented by B, then both BB and Bb would result in big feathers.

Dominant

The copies (alleles) of the gene are both the same. This word comes from the Greek word "homos," which means same, and "zugos," which means yoked. BB bb

Homozygous

The genetic information of an organism. The term can also refer to the genetic information of a specific trait.

Heterozygous

There are two different versions (alleles) of the gene. This word comes from the Greek words "heteros," which means different, and "zugos," which means yoked.

Genotype


The trait will only be expressed if there are two identical copies (alleles) of the gene. For example, if small feathers are represented by b, then only the genotype of bb will produce small feathers.

Phenotype

The observable characteristics or traits of an individual, such as having either small or large feathers.

Laws of heredity

Yujun buys two pure-bred cats, each with a pedigree showing that all of their ancestors have the same trait for hair length. One cat is long haired. The other is short haired. Now the cats are going to have kittens. Yujun knows that hair length is a recessive/dominant trait but doesn't remember which trait is dominant. Match each of the possible outcomes with the correct conclusion.




1. SEGREGATION - the alleles are distributed randomly

Mendel's first law states that the chance of inheriting an allele (or genetic factor) is random. For example, in the case of yellow or green seeds in pea plants, if the parent is heterozygous (Yy) there is a 50% chance the offspring will receive the dominant yellow (Y) allele and a 50% chance they will receive the recessive green (y) allele. This is called the principle of segregation.

Try it yourself!

Supplies: 2 cups and 4 objects to represent dominant and recessive alleles. The objects should have the same shape and size but different colors. You could use dried peas, beans, marbles, dice, marshmallows or any items of the same size. Just make sure they are marked or colored so that you can tell them apart.




→ **STEP 1**

Place 1 object of each color in each cup. Assign one of the colors to be dominant (A) and the other to be recessive (a).



→ **STEP 2**

Without looking, draw one object from each cup.



→ **STEP 3**

Record whether you picked two dominant (AA), one of each (Aa), or two recessive (aa) in the tables below. Go back to **STEP 1** and repeat until the table is full.

$\text{OO} \rightarrow \text{AA}$
 $\text{Oo} \rightarrow \text{Aa}$
 $\text{oo} \rightarrow \text{aa}$

AA	Aa	aa
----	----	----

→ **STEP 4**

Color each cell green if it holds aa and yellow if it holds AA or Aa. When complete, do you expect the table will have more yellow or green squares? Expect more yellow

AA																				
AA																				

Count the number of yellow and green cells in your table.

Answers will vary! The first time we did this, we had 26 yellow and 14 green. Our second attempt had 31 yellow and 9 green.

26

Yellow

14








Green

Mendel's first law predicts 30 yellow (AA and Aa) and 10 green (aa). Were your results similar to this? If you repeated this activity 3 more times, would you expect the overall ratio to be closer to or further away from $\frac{3}{4}$ yellow and $\frac{1}{4}$ green?

Yes. If repeated more, we would expect the ratio to come closer to $\frac{3}{4}$ yellow and $\frac{1}{4}$ green.

Punnett Squares

Since the probability of getting one allele or another is 50/50, we can use a tool called a Punnett square to see all of the possible combinations AND how likely they are to occur! For example, if a round shape is dominant and pear shape is recessive, a punnet square shows us how likely it is that two heterozygous aliens would produce a pear-shaped alien:



Practice some crosses!

Fill in the Punnett squares for each of the following crosses. Two eyes (B) is a dominant trait. One eye (b) is recessive. Write down the chance of producing a one-eyed alien for each cross. If no squares are "bb," then a one-eyed alien can't be produced from that cross (0% chance). If one square is bb, the chance is 25%. If two squares are bb, the chance is 50%. If three squares are bb, the chance is 75%. If all four squares are bb, the chance is 100%.

Bb x Bb

We usually just write "bB" as "Bb" so that all the heterozygous genotypes look alike.

B	BB	Bb
b	Bb	bb

25 % chance of bb

BB x bb

B	b	
B	Bb	Bb
b	Bb	bb

0 % chance of bb

BB x BB

B	BB	BB
B	BB	BB

0 % chance of bb

Bb x bb

b	b	
B	Bb	Bb
b	bb	bb

50 % chance of bb

BB x Bb

B	BB	Bb
B	BB	Bb

0 % chance of bb

bb x bb

b	b	
b	bb	bb
b	bb	bb

100 % chance of bb

With cats, white fur is dominant (W) and black hair (w) is recessive. **Before** you fill out each Punnett squares, check a box to make a prediction about whether the cross will produce black kittens. **After** filling out the Punnett square, mark the percentage of black kittens the cross produced.

Will there be black kittens?

yes no

Ww x WW

w	w	
W	Ww	Ww
w	ww	ww

Black kittens produced:

<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
0%	25%	50%	75%	100%

Will there be black kittens?

yes no

WW x Ww

W	W	
W	WW	Ww
W	WW	Ww

Black kittens produced:

<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
0%	25%	50%	75%	100%

Will there be black kittens?

yes no

WW x WW

w	w	
W	Ww	Ww
W	Ww	Ww

Black kittens produced:

<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
0%	25%	50%	75%	100%

Will there be black kittens?

yes no

Ww x Ww

W	W	
W	WW	Ww
w	Ww	WW

Black kittens produced:

<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
0%	25%	50%	75%	100%

Will there be black kittens?

yes no

WW x WW

w	w	
W	WW	WW
W	WW	WW

Black kittens produced:

<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
0%	25%	50%	75%	100%

Will there be black kittens?

yes no

WW x WW

w	w	
w	WW	WW
w	WW	WW

Black kittens produced:

<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>
0%	25%	50%	75%	100%

2. INDEPENDENT ASSORTMENT - traits are inherited independently

Mendel's second law of inheritance says that the inheritance of one trait (such as a cat having long hair or short) has no influence on the inheritance of another trait (such as the color of the cat's fur). This is called the principle of independent assortment. It applies to a lot of traits, but not all of them! Some genes are inherited together because they are located close together on the same chromosome.


When talking about hair color, it doesn't matter if the hair is long or short; cats with the **W** allele will be white, while cats with the **ww** genotype will be dark-haired.

Genotype	SS	Ss	ss	WW	Ww	ww
Phenotype	SHORT HAIR	SHORT HAIR	LONG HAIR	WHITE HAIR	WHITE HAIR	DARK HAIR

When talking about hair length, it doesn't matter what color the hair is; cats with the genotype of **SS** or **Ss** will have short hair. Long hair is only produced by the genotype **ss**.

Could this pair of cats produce a black long-haired kitten? Check yes or no. Then show why or why not.


- YES
 NO



The only genotype that will give a black long-haired kitten is **wwss**. The kitten can get the recessive alleles from both parents.

Could this pair of cats produce a white long-haired kitten? Check yes or no. Then show why or why not.


- YES
 NO



There is no dominant white allele (**W**) to inherit. Also, the kitten needs the genotype **ss** for long hair, but since it receives one allele from each parent, its genotype for hair length can only be **Ss**.

Could this pair of cats produce a black short-haired kitten? Check yes or no. Then show why or why not.


- YES
 NO



Yes. It is possible to pass on the genotype **wwSs**. The **ww** gives the color black, and **Ss** results in short hair.

Could this pair of cats produce a white long-haired kitten? Check yes or no. Then show why or why not.

- YES
 NO



The kitten can inherit **Wwss**. **Ww** yields white fur, and **ss** yields long hair.

The traits of pea height (tall or dwarf) and the trait of flower color (purple or white) are inherited independently.


When talking about flower color, it doesn't matter if the plants are tall or dwarf; peas with the **P** allele will be purple while peas with the **pp** genotype will be white.

Genotype	TT	Tt	tt	PP	Pp	pp
Phenotype	TALL PLANTS	TALL PLANTS	DWARF PLANTS	PURPLE FLOWERS	PURPLE FLOWERS	WHITE FLOWERS

When talking about plant height, it doesn't matter what color the flower is; peas with the genotype of **TT** or **Tt** will be tall. Dwarf pea plants are only produced by the genotype **tt**.

Could this cross produce a dwarf pea plant with purple flowers? Check yes or no. Then show why or why not.


- YES
 NO



The genotype **tt** yields a dwarf plant and **Pp** makes purple.

Could this cross produce a tall plant with white flowers? Check yes or no. Then show why or why not.


- YES
 NO



The new plant will have a genotype of **tt**, but we need **TT** or **Tt** to get a tall plant.

Could this cross produce a dwarf plant with white flowers? Check yes or no. Then show why or why not.


- YES
 NO



This cross will always result in tall purple flower plants. For this cross, **PP** or **Pp** are the only possible genotypes for flower color and **TT** and **Tt** are the only genotypes for height.

Could this cross produce a tall plant with white flowers? Check yes or no. Then show why or why not.


- YES
 NO



The genotype **Tt** yields a tall plant and **Pp** makes purple.

Could this cross produce a tall plant with purple flowers? Check yes or no. Then show why or why not.


- YES
 NO



The genotype **Tt** yields a tall plant and **Pp** makes purple.

Could this cross produce a dwarf plant with white flowers? Check yes or no. Then show why or why not.

- YES
 NO




All of the offspring will be dwarf plants (**tt**), but the genotype for color will be either **PP** or **Pp**, resulting in a purple plant.


The Munchkin Mutation

Riley is fostering a pregnant Munchkin cat named Percy for their local animal shelter. The short legs of the Munchkin are caused by a dominant mutation (**M**). The mutation is lethal if homozygous (the **MM** genotype fails to develop in utero). Heterozygous (**Mm**) gives the Munchkin phenotype of short legs, and homozygous recessive (**mm**) results in long legs (**wildtype**).


The animal shelter has a waiting list of people ready to adopt Percy's kittens. They want to know what the probability is of the kittens being Munchkins, but no one knows if Percy's mate was a Munchkin (**Mm**) or not (**mm**). Calculate the probability of Munchkin kittens from each possible cross.



Possibility 1: Munchkin x Munchkin cross



Possibility 2: Munchkin x Wildtype cross



What is the probability of Munchkin kittens if both parents are Munchkins?

- 100% - every kitten will be a Munchkin.
- Approximately 67% - 2 out of every 3 kittens will be a Munchkin.
- 50% - there's a 50/50 chance of the Munchkin trait.
- 25% - there's a 1 in 4 chance of the Munchkin trait.

What is the probability of Munchkin kittens if one parent is a Munchkin and the other is wildtype (has long legs)?

- 100% - every kitten will be a Munchkin.
- Approximately 67% - 2 out of every 3 kittens will be a Munchkin.
- 50% - there's a 50/50 chance of the Munchkin trait.
- 25% - there's a 1 in 4 chance of the Munchkin trait.

If Percy is pregnant with only two kittens, what are the possible outcomes for this litter?

Mark all genotype combinations that are possible.

- Mm and Mm Even though we don't know the other parent's genotype, each kitten could be either Mm or mm (inheriting M or m from Percy), so all three possibilities can occur.
- Mm and mm
- mm and mm

Ethics and Genetics


The study of genetics is about more than traits and DNA. It often impacts social and ethical questions as well. Read the following paragraphs and then write your opinion about the concerns associated with each example. There are no "right" answers here! These questions are as much about what you value as they are about heredity and genetics.

A group of dog breeders have developed a new dog breed called a Tardal.* In working to establish the Tardal as an officially recognized breed, full health histories are completed for all Tardal dogs. The studies reveal that Tardals are intelligent, loyal, and have the fastest running speed of any dog on Earth, capable of running at 70 mph. But in addition to these desirable traits, Tardals are 11x more likely than other dogs to experience hip and ankle dislocation. They also have a life expectancy that is significantly shorter than other dogs, living for just 4 to 5 years.


Should this new dog breed be accepted, a move that would cause it to become a common breed among dog owners? Or should people stop breeding Tardals because of the health concerns associated with the breed? Write your recommendation:

Answers will vary. Similar questions to this one impact several dog breeds today. Pugs are more susceptible to eye problems and breathing issues, and large breeds such as Danes and German Shepherds are more likely to suffer from hip dysphasia.

*The Tardal is an imaginary dog breed invented for this lesson.




Remy breeds ball pythons and in one of their recent clutch of eggs they discovered a scale-less snake. Rare python morphs are valuable and Remy knows they can earn money by breeding the snake to produce more pythons without scales. But this scale-less snake has incredibly soft skin. It needs special care when shedding to avoid infection. Even with soft bedding, it is easily injured and often experiences minor cuts and scrapes from everyday movements.



Do you think Remy should breed the scale-less snake? Why or why not?

Answers will vary. Scaleless python morphs exist in real life and some snake breeders argue that they should not be bred because of their susceptibility to injury.



The Governing Council of the Cat Fancy* currently refuses to recognize Munchkins as a breed because of health issues associated with the short legged trait. Some Munchkin owners claim that the cats are healthy and happy with no more health concerns than the average cat. Others claim that the mutation hinders Munchkins ability to move and causes discomfort.

If you were a member of the GCCF, would you be in favor of recognizing Munchkins as a breed, be neutral on the issue, or be in favor of prohibiting further breeding of Munchkins?

Answers will vary.


*An organization that registers pedigreed cats in the United Kingdom

How to Read A Pedigree Chart


A pedigree chart is a family tree that tracks how a trait has been passed along. It typically follows the rules below:

- Males are represented by squares, and females are represented by circles.
- A filled shape means the individual exhibits the trait.
- Horizontal lines between two individuals exhibit mating pairs.
- Vertical lines descend from parents to a horizontal line that connects the offspring.

Example of a pedigree tracking the dominant trait **A**




Example of a pedigree tracking the recessive trait **b**




Each individual receives their alleles from their parents. When tracking recessive traits, sometimes it helps to identify the heterozygous individuals that carry the trait. These **carriers** are marked with half-shading.


In this example, neither parent has the recessive trait, but one of the offspring does!



This means both parents were carriers of the trait! We can show this by shading half of each shape.




Color in this pedigree for tracking **pear shape** in the aliens. Remember, round body shape is dominant (RR) and pear shape is recessive (rr). The phenotype is known for all individuals. Can you determine the genotypes of the aliens indicated by the arrows?





Pet Pedigree Puzzle #1


Profile of a breed with feathered feet



Profile of a breed without feathered feet




The "feathered" foot has long hair on the lower leg that covers the hoof.




A non-feathered foot has short hair.

Below is a pedigree chart for the trait of "feathered feet," the long hair that cover the hooves of certain horse breeds like Clydesdales, Shires, Friesians, and Ardennes.* Each shaded shape represents a horse with feathered feet.




Can a dominant trait skip a generation? What about a recessive trait? Explain.

A dominant trait can't skip a generation. If the allele is present then the dominant trait will be expressed. Recessive traits can skip a generation.



Is the trait mapped in this pedigree chart dominant or recessive? Explain how you know.


The feathered feet trait skipped a generation, so it must be a recessive trait.




AA, Aa, or aa?

When tracking a recessive trait, a half-shaded shape is used to indicate a carrier. Can you identify a horse or horses in this pedigree that must be carriers of the trait? If yes, shade in their shapes.

*While a cross between a feathered horse (such as a Clydesdale) and a non-feathered horse (such as a Thoroughbred) will result in a horse that is non-feathered, the genetics of feathering have not been studied in detail. It's likely to include more than one gene and be more complicated than the hypothetical example used here!





A carrier of the trait!




Pet Pedigree Puzzle #2

At Ella's pet store, albino blood pythons have been the most popular snake breed. Ella has space to breed two snakes and wants to choose a pairing that will produce more albinos. Fortunately, Ella has pedigree charts of the store's four blood pythons: Julius Squeezer, Basil, Medusa, and Naga.


Help Ella make the choice that will give the highest chance of more albino pythons! Remember that albinism is a recessive trait and the pedigrees only show phenotype, not genotype.



Julius Squeezer is the an albino blood python with beautiful cream and orange coloring.



Basil has standard coloring and a very friendly temperament. He is easy to hold.



Naga has standard coloring and an aggressive temperament. Gloves are required for handling.

What genotype is Julius Squeezer? The standard color allele is represented by **A**. The albinism allele is represented by **a**.

AA or Aa Aa aa

What genotype is Basil?

AA or Aa Aa aa

What genotype is Naga?

AA or Aa Aa aa

What genotype is Medusa?

AA or Aa Aa aa


Which snakes should Ella select for breeding and why?

Julius Squeezer and Medusa are the male and female pair most likely to pass on the aa genotype, so they are most likely to have albino offspring.

Mendel's First Law is like flipping a coin...


The probability of an **aa** x **Aa** cross giving albino snakes may be 50/50, but that doesn't mean you'll see that exact ratio in real life! Explore Mendel's first law by using a coin to determine whether these snakes produce albino blood pythons. Color the albinos (**aa**) orange. Use brown to mark the standard coloring (**AA** or **Aa**) which is also called **wildtype**.

Julius Squeezer will always give the albino allele for coloring, but Medusa is heterozygous and can give either the **A** or **a** allele. The first time this pair are bred, they have a clutch of 14 eggs. Flip a coin for each of the 14 snakes to see which allele is given. If it's heads, write **Aa**. If tails, write **aa**. Then color in the shapes to represent the number of blood pythons in the clutch of 14 snakes.




The second time Medusa and Julius Squeezers breed again, they have a clutch of 14 eggs. Flip a coin for each of the snakes to see which allele is given. If it's heads, write **Aa**. If tails, write **aa**. Then color in the shapes to represent the number of blood pythons.

Your ratios will be different depending on the coin flip. But overall, it will likely be approximately half orange/albino and half brown/wildtype.




Assuming Naga is heterozygous (**Aa**), flip a coin to see what proportion of this clutch of eggs would be expected to be albinos. Then color in the results.

If Naga is heterozygous, then approximately half of the eggs would hatch albino snakes.



Assuming Naga is homozygous (**AA**), what proportion of this clutch of eggs would be albinos? Do you need to flip a coin here?


All the offspring will be **Aa**, so there is no need to flip a coin.



Bigger Punnett Squares!

When we consider two traits at once, we concatenate (or join) the genotypes. For example, an alien that is double heterozygous for roundness (**Aa**) and number of eyes (**Bb**) has the genotype **AaBb**. An pear-shaped alien that is heterozygous for eyes would have the genotype **aaBb**.

The double heterozygous alien has an equal chance of passing along each of the 4 allele pairs: **AB**, **Ab**, **aB**, and **ab**. The pear-shaped alien also has an equal chance of passing on each of the 4 possible allele pairs.



Fill in the punnet square below by drawing the correct number of eyes on each alien for the cross **AaBb** x **aaBb**. Remember **BB** or **Bb** result in the phenotype of two eyes, and only **bb** gives the phenotype of one eye.

	aB	ab	aB	ab
AB				


Genotype: Phenotype:

AA	Round body shape	
Aa	Round body shape	
aa	Pear body shape	

Genotype: Phenotype:

BB	Two eyes	
Bb	Two eyes	
bb	One eye	


Create a Punnett square for the cross **AaBb** x **AaBb**. Will you see the rare pear-shaped one-eyed alien (**aabb**) from this cross? If so, what is the chance of seeing it? 1/16, 2/8, 1/4 or 1/2?



Only 1 in 16 will be pear shaped
and one-eyed.


AB	Ab	aB	ab
AB			

Create a Punnett square for the cross **aaBb** x **Aabb**. What is the chance of seeing a round two-eyed alien (**AaBb**) from this cross?




All 4 possible phenotypes occur
1/4 of the time with this cross.

Ab	Ab	ab	ab
Ab			




Genotype:	Phenotype:
AA	Round body shape
Aa	Round body shape
aa	Pear body shape

Genotype:	Phenotype:
BB	Two eyes
Bb	Two eyes
bb	One eye



Create a Punnett square for the cross **AABb×AaBb**. How many of the offspring, on average, will have one eye?
1/16, 2/8, 1/4 or 1/2? Will there be any pear-shaped one-eyed aliens or will they all be round?




Only 1 in 4 will have one eye.

All aliens will be round. There
will be no pear-shaped aliens.

		AB	Ab	aB	ab
AB	AB				
	AABB	AABb	AaBB	AaBb	
Ab	AB				
	AAbB	AAbb	Aabb	Aabb	
Ab	AB				
	AABB	AABb	AaBB	AaBb	
Ab	AB				
	AAbB	AAbb	Aabb	Aabb	

Create a Punnett square for the cross **aabb × AABB**.
What is the chance of seeing a round two-eyed alien
that is heterozygous for both traits (**AaBb**)?




All the offspring will be **AaBb**

because they inherit the dominant
alleles from one parent and the
recessive alleles from the other.


		AB	AB	AB	AB
ab	ab				
	aAbB	aAbB	aAbB	aAbB	
ab	ab				
	aAbB	aAbB	aAbB	aAbB	
ab	ab				
	aAbB	aAbB	aAbB	aAbB	

Genotype:	Phenotype:
AA	Round body shape
Aa	Round body shape
aa	Pear body shape

Genotype:	Phenotype:
CC	Tentacle arms
Cc	Tentacle arms
cc	No arms



A new dominant trait of tentacle arms has been discovered. What will be the results of a cross between a heterozygous tentacled round alien and a pear-shaped no-arms alien? (**aacc×AaCc**) Fill out the square, then write the ratios for each of the outcomes.



1 in 4 are round with tentacles.


1 in 4 are round without tentacles.

1 in 4 are pear-shaped with tentacles.

1 in 4 are pear-shaped without tentacles.

		AC	Ac	aC	ac
ac	AC				
	AaCc	Aacc	aacC	aacc	
ac	AC				
	AaCc	Aacc	aacC	aacc	
ac	AC				
	AaCc	Aacc	aacC	aacc	

Can two round-eyed tentacle-arm aliens produce a
pear-shaped no-arms alien? If yes, what do the
genotypes need to be for this cross, and what is the
chance of the pear-shaped no-armed offspring?




Yes. To get both recessive traits, the parents
must be heterozygous (both **AaCc**). The
chance of producing **aacc** is **1/16**.

For more on Punnett squares, check out pages 112-117 in the appendix! This bonus section include exercises to sharpen your skills for calculating probabilities, and templates for a tryhybrid and tetrahybrid cross!


Heredity Word Search

There are a lot of new words to learn when studying biology. Repetition is the best way to learn them, and word games can be part of that! Find each of the hidden words in the word-search. The words can run in any direction: horizontal or diagonal, and the letters might go left to right or right to left!

GENE	RECESSIVE	CARRIER
ALLELE	PHENOTYPE	PROBABILITY
HOMOZYGOUS	GENOTYPE	HEREDITY
HETEROZYGOUS	CODOMINANCE	OFFSPRING
PUNNETTSQUARE	PEDIGREE	TRAIT
DOMINANT	HYBRID	




Heredity Crossword Puzzle



VERTICAL Words

- What is the name of the father of genetics?
- What is the term for an organism's physical appearance or visible traits?
- The name for a chart of the ancestry or heritage of an individual.
- A trait that will appear in the offspring if one of the parents contributes an allele for it.
- A trait that will only appear in the offspring if both parents contribute an allele for it.
- The scientific study of heredity.
- The term for the condition where both copies of the allele are the same.
- The passing of traits from parents to offspring.
- The term for a possible form of the gene.
- A type of square used to keep track of the possible combination of alleles that can result from a cross.
- A segment of DNA on a chromosome that codes for a specific trait.
- A person who has one recessive allele for a trait and one dominant allele, but does not have the trait.
- The term for the condition where there are two different alleles for a trait.

HORIZONTAL Words



Quiz Time!

ANSWER THE QUESTIONS TO
SEE WHAT YOU LEARNED
ABOUT HEREDITY!

- 1 Which statement is true?
 - A. Members of the same species must be approximately the same size.
 - B. Members of the same species can breed with each other and produce offspring.**
 - C. Scientists always agree on whether animals are in the same species.
 - D. Only animals have species.

- 2 Which type of reproduction will result in the most genetic diversity?
 - A. Fragmentation
 - B. Binary fission
 - C. Vegetative propagation
 - D. Spore formation
 - E. Sexual reproduction**

- 3 Every trait is controlled by a single gene, and each gene is determined by two alleles.
 - A. True
 - B. False**

- 4 Select each true statement below.
 - A. A dominant trait can skip a generation.
 - B. A recessive trait can skip a generation.**
 - C. When two parents share the same genotype, their offspring will share the same genotype.
 - D. When two parents share the same phenotype, their offspring will share the same phenotype.

- 5 What does the notation BB mean to a geneticist?
 - A. Two dominant alleles**
 - B. Two recessive alleles
 - C. At least one dominant allele
 - D. One dominant and one recessive allele

- 6 If a homozygous black guinea pig (BB) is crossed with a homozygous white guinea pig (bb), what is the probability that an offspring will have black fur?
 - A. 0%
 - B. 25%
 - C. 50%
 - D. 75%
 - E. 100%**

- 7 If an organism is homozygous dominant for a trait that follows Mendelian inheritance patterns, which of the following must be true?
 - A. Its offspring will also have a homozygous genotype.
 - B. Its offspring will also have a heterozygous genotype.
 - C. Its offspring will exhibit the dominant phenotype for the trait.**
 - D. The organism's parent(s) also had a homozygous genotype for the trait.
 - E. None of the above

- 8 Select each true statement below.
 - A. If you know an organism's genotype for a trait, then you also know its phenotype.**
 - B. If you know an organism's phenotype for a trait, then you also know its genotype.
 - C. A parent always has the same genotype as its offspring.
 - D. A parent always has the same phenotype as its offspring.


- 9 An organism that has two different alleles for a single trait is said to be heterozygous for that trait.

- 10 An organism has a phenotype for a dominant trait while some of its offspring have the phenotype for the recessive trait. What can we conclude about this organism?
 - A. It is heterozygous for the trait.**
 - B. It is homozygous dominant for the trait.
 - C. It is homozygous recessive for the trait.
 - D. There is not enough information to tell.

- 11 True or False: all alleles are either dominant or recessive.
 - A. True
 - B. False**

- 12 What is a Mendelian trait?
 - A. A trait that involves alleles
 - B. A trait that is passed down by dominant and recessive alleles of one gene**
 - C. A trait where different alleles are equally and independently expressed
 - D. A trait exhibited by pea plants

Below is a pedigree chart for tracking **hornless cattle**, a Mendelian trait. Cattle without horns are called *polled* cattle and are usually viewed as more desirable than cattle with horns. Each of the cattle is numbered for reference.



- 13 Is the hornless trait (polling) a dominant or recessive trait in cattle? Explain how you know.
Polling is dominant. Individuals 5 and 6 have the trait, but 9 doesn't. If polled cattle is a recessive trait, then ALL of the offspring from two polled cattle would be polled. If polled cattle is a dominant trait, then two polled cattle can have a calf with horns. This latter case is what we observe in the pedigree.

- 14 We can assign the alleles A and a to the trait "no horns" exhibited in the pedigree chart. In that case, what is the specific meaning of both A and a?
"A" is the allele for the gene for no horns or polling. "a" is the allele for horns.

- 15 Label each member of the pedigree chart whose genotype can be fully identified. Are there any that you can't determine? Explain.
The genotype for 1, 3, 7, and 9 is aa because they don't have the trait of polling.
The genotype for 4, 5, 6, and 8 is Aa.
2 and 10 can't be determined. They could be either AA or Aa.

- 16 List the numbers of each individual that has horns in the pedigree above:
1, 3, 7, and 9 have horns.


- 17 List the numbers of the individuals that are heterozygous in the pedigree above:
4, 5, 6, and 8 are heterozygous (carriers) for the "no horns" trait.

- 18 If we cross two cattle with horns, could the offspring have no horns? Explain.
No. Having horns is a recessive trait, so two cattle with horns will only pass on recessive alleles causing the offspring to also be recessive. The cross aa x aa can only produce aa.


- 19 If we cross two cattle with no horns, could the offspring have horns? Explain.
Yes. If both parents are heterozygous, then the cross Aa x Aa can produce aa. In that case, two cattle without horns would have offspring with horns.

The Genetic Code

Random words don't communicate ideas.



Words in the correct sequence communicate ideas!



FILL IN THE BLANKS USING THESE WORDS:


four thymine garage repeating genome trees

DNA stands for deoxyribonucleic acid. It's a long strand of repeating units which are called nucleotides. The complete sequence of DNA within an organism is called its genome.

There are four different nucleotides in DNA: adenine, thymine, guanine, and cytosine. They fit together to form pairs: **adenine** pairs with thymine, and **cytosine** pairs with **guanine**.

These pairings are what's being referred to when we read that the human genome contains approximately 3 billion **base pairs**. If you ever forget which base pairs with another, remember, "Apples grow on trees, park your car in a garage."


LABEL THE BASE PAIR, BACKBONE, AND NUCLEOTIDE IN THIS CARTOON REPRESENTATION OF DNA. THEN WRITE A, T, C, OR G IN EACH BOX TO FILL IN THE CODE!



What's a gene?


A gene is a specific sequence of DNA that contains information to make a gene product such as a protein. Most organisms have thousands of genes. There are approximately 20,000 genes in the human genome!

To better understand how a gene works, we'll look at two examples with genes that affect a pigment called melanin.



Albinism

Albino rats have completely white hair and pink eyes because they lack the pigment melanin. Blood vessels behind their transparent eyes are the reason for the pink color. The lack of pigment makes albinos much more sensitive to light. They sunburn easily, and their eyes can be damaged by bright lights.




The cause: a mutation in the DNA causes a protein that makes melanin to stop working. Why would this result in a recessive trait?

Any gene that codes for a working copy of the enzyme will produce melanin and the pigment will be visible in the phenotype. A lack of melanin can only happen when there are two defective genes.

Melanism

Black panthers are actually leopards with melanism. The extra pigment turns their coat completely black, which is not a disadvantage when hunting prey at night! Examples of melanism have been found in many different mammals and reptiles. In each case, the skin, feathers, or fur of the animal is completely black due to the extra pigment.




The cause: a mutation in the DNA causes a protein that makes melanin to work overtime! Why would this result in a dominant trait?


If there is one working copy of the gene, the result will be an over-abundance of melanin pigment.




Write the correct phenotype below the cartoon representing each genotype. Possible phenotypes include albino, melanistic, or wild type (average melanin levels).*




wild type



albino



melanistic



wild type


*This example is over-simplified for educational purposes. While melanism in jaguars is a dominant trait, in other animals melanism is often recessive! There isn't just one type of melanin, and there are multiple enzymes and pigments involved in the process. So in the real world, pigmentation is more complex than displayed here.

Flower Color in Mendel's Peas

Remember the example of flower color that Mendel observed, where purple was dominant and white was recessive? The "bossy" P allele is representing a piece of DNA that has instructions for an enzyme. The P enzyme makes the purple pigment in the flower. It's efficient and fast, and one copy can do just as good a job at making the flower purple as two copies. The recessive allele is representing a piece of DNA with a mutation that makes a non-functional enzyme. It can't make the purple pigment.

Draw pictures in the boxes below representing the working protein that makes purple pigment and the mutated protein that can't make the pigment:

Drawings can be cartoonish and fun or more realistic. The main point is that one drawing shows an enzyme making lots of purple pigment, and the other shows an enzyme that cannot make pigment.



ARE THESE FACT OR FICTION? Write your verdict below each statement.

The Australian lungfish has the largest genome of any animal.

IT'S HEAVY CARRYING AROUND ALL THIS DNA.

FACT - its genome has 43 billion base pairs!

If you type 8 hours a day at 60 words per minute, it would take approximately 2 years to type the human genome!

WHY AM I USING A QWERTY KEYBOARD TO TYPE A, T, C, AND G? MY RIGHT HAND IS SO BORED.

FICTION - it would take 50 years!


Identical twins have the same DNA!

It all depends on what you mean by identical! It can be argued to be FACT in that monozygotic twins come from a single cell. They will almost always have the same alleles or genes.

It can be argued to be FICTION because there can be mistakes in cell division early on in development that could cause twins to have different chromosomes (ex: one twin could be XY, the other XO). Also, as the twins age there will be a relatively small number of differences due to mutations, as well as epistatic changes to how the DNA is expressed. Even with identical genes, twins will have measurable physical differences between their DNA.

For most cases and for the most common use of the word "identical," we think the best answer is "FACT." This is one reason why identical twins have been and continue to be an important part of understanding the role genetics plays in different diseases and traits.

Chromosomes



FILL IN THE BLANKS USING THESE WORDS:

number histones chromosome fly 23 karyotype humans



A **chromosome** is a strand of DNA wrapped around proteins called **histones**.

Each member of the same species will have the same number of chromosomes, but between

different species, the **number** and size of chromosomes can be very different. A fruit **fly** has 8 chromosomes. The black mulberry tree has more than 300! **Humans** have **23** pairs of chromosomes for a total of 46 chromosomes. A

karyotype is a picture of the complete set of chromosomes that have been sorted by length.

Draw lines to match each organism with its karyotype:



The human genome

There are 23 pairs of chromosomes in the human genome, but what do they all do? Lots of things! Remember each chromosome contains *hundreds to thousands* of different genes. Having only one chromosome or having three copies of a chromosome (trisomy) is usually fatal. There are only a few chromosomes where an extra copy can be tolerated.

Vocabulary review! Fill in the blanks below with the correct term:

The condition of having three copies of a chromosome is called:

trisomy


Chromosomes contain both DNA and proteins. The proteins are called:

histones

A sequence of DNA that contains information to make a product such as a protein is called:

a gene

Arranging chromosomes in a karyotype helps scientists organize them for further study. In the human karyotype, the chromosomes are arranged from longest to shortest, with the XY pair at the end.



Write the number of the corresponding chromosome next to each statement:

1 The longest chromosome in the human genome. It contains thousands of genes.

23 This pair of chromosomes determines biological sex.

21 Three copies of this chromosome causes Down Syndrome.

18 Three copies of this chromosome causes Trisomy 18.

22 It was thought to be the shortest human chromosome, but then chromosome 21 was discovered to be shorter!

Sex Determination

FILL IN THE BLANKS USING THESE WORDS:


male 50 eyesight 900 females birds

Most organisms that reproduce sexually have two sexes: male and female. In humans, biological sex is determined by chromosome 23. **Females** have two copies of the X chromosome. The X chromosome is long and has about **900** genes, including important alleles for **eyesight**, blood clotting, hearing, and nerve and muscle function. The Y chromosome has about **50** genes, and their purpose is to start a cascade of hormone changes that cause the development of the **male** physical traits.


In **birds**, some fish, and most reptiles, the sex determination is different. Males have two copies of the same chromosome and females have two different sex chromosomes. This system is called the ZW system.

ZW system - birds, some fish, reptiles, and pistachio trees


Male bird is ZZ



female bird is ZW




Fill out the Punnett square for ZW. On average, how many birds will be male, and how many will be female?




On average, $\frac{1}{2}$ female and $\frac{1}{2}$ male

XY system - mammals, some insects, guppies, and ginkgo trees


Male mammal is XY



female mammal is XX



Fill out the Punnett square for XY. On average, how many mammals will be male? How many will be female?




On average, $\frac{1}{2}$ female and $\frac{1}{2}$ male

Female	Male	Intersex	Klinefelter Syndrome	Turner syndrome
XX	XY	XX or XY	XYY	X
The individual has ovaries that produce eggs through meiosis.	The individual has testes that produce sperm through meiosis.	Physical characteristics don't match traditional definition of male or female. Observed phenotype is sometimes opposite what would be expected from the genotype.	Physical characteristics are usually male. Individual is often infertile. Occurs in about 1 in 1,000 male births.	Physical characteristics are female. Individual is infertile. Higher risk of heart disease, diabetes, and thyroid disorders.

MEIOSIS





Meiosis is a special type of cell division used to produce **gametes**. Gametes have $\frac{1}{2}$ the number of chromosomes as their parent cell and are called **haploid** cells. Combining gametes from two different individuals increases genetic diversity, which is one of the main advantages in sexual reproduction.



Meiosis Mysteries

Seedless watermelons, bananas, and mules all have something in common. They can't reproduce!* Match each organism with the corresponding explanation for why it is sterile or infertile.

*The domestic banana (Cavendish) can reproduce asexually through vegetative propagation but does not have seeds and cannot undergo sexual reproduction. There are hundreds of different varieties of bananas. Most of the wild bananas do produce seeds.



A mutation stopped meiosis from working. People first noticed this mutation several hundred years ago. They liked the result and have been nurturing similar mutations ever since!

One parent has 64 chromosomes. The other parent has 62. The resulting organism has an odd number of chromosomes (63), which causes problems in meiosis.

One parent is diploid (2 copies of each chromosome) and the other parent is tetraploid (4 copies of each chromosome). The resulting triploid organism is infertile or sterile.

Both of the parents have 60 chromosomes. Female offspring are fertile, but males are sterile because of some interesting and complex epigenetic changes to meiosis.

Proteins!


The molecules that run things in the cell

FILL IN THE BLANKS USING THESE WORDS:

acids	proteins	muscles	duplicating
traits	make	amino	structural

Almost all of the traits in animals and plants are ultimately controlled by proteins. Proteins can be enzymes that make things like pigments or perform tasks like duplicating DNA. Proteins can also be structural; they're the main ingredients in the muscles, skin, and bones of all animals. All proteins are polymers made of amino acids.

Below are the 20 amino acids used to make proteins. Each amino acid has a 1-letter abbreviation. The 9 dietary essential amino acids are shaded grey.



Label Hunt: Check some food labels. Can you find amino acids listed in the ingredients?


FOOD ITEM: AMINO ACID IT CONTAINS:

Sugar-free gum phenylalanine
Diet soda phenylalanine

Hint: check sugar-free products such as chewing gum. You'll find one!

A Code of Triplets

A **codon** is a sequence of 3 nucleotides such as adenine, thymine, and cytosine (ATC). Each codon represents an amino acid or an instruction like "start" or "stop." On the previous page, you can see the amino acid tryptophan is specified by the triplet TGG and abbreviated with the letter W. Some amino acids can be specified by multiple codons. For example, both AAA and AAG code for the amino acid lysine.



Can you decipher a secret message from the letters below?

...YRTQUNYOUCANEAATPIEANYDAYMMETBK...

It helps if we break up the phrase into triplets:

...YRT QUN YOU CAN EAT PIE ANY DAY MME TBK...

Now what happens if we shift our starting position by just one letter?

...RTQ UNY OUC ANE ATP IEA NYD AYM MET BK...

The secret message is lost!

 The genetic code is similar. Adding or taking away one base pair can have disastrous effects! 

DNA Decipher Challenge

Break up the letters below into groups of three nucleotides (codons). Match each codon with the abbreviation letter of the associated amino acid by referring to the previous page. Can you finish deciphering the secret message? Use the chart on the previous page OR use the DNA decoder wheel from the appendix.

AGTTGCATTGAAAATGTGAGATCTAATAAACATGTATGATAACGCG
S C I E N C E I S I N M Y D N A

What did one chromosome say to the other?  Do these genes look good on me?

DNA Encryption Challenge

Write out the DNA codons that will code for the word "LEARNING." Use the abbreviation letters for the amino acids. For example, "L" can be coded for by "CTC" or "CTT." Then give the sequence to someone and see if they can decode it!

Tip: For faster code deciphering, use the codon wheel in the appendix!

SCIENCE MAM There are multiple correct answers. The letters below show just one possible code. 51

Hands-on Science Project

GUMMY WORM KARYOTYPES


MATERIALS:



knife or scissors



gummy worms



chocolate




Small bowl for melting chocolate

Make a fruit fly karyotype

The common fruit fly is used as a model organism for biological research because it has a quick life cycle and a relatively small genome with just 4 pairs of chromosomes.

The first pair of chromosomes in the fruit fly karyotype is the X/Y pair that determines biological sex or the female/male trait. The other chromosomes are named 2, 3, and 4. But chromosome 4 is so small that it only has a handful of genes compared to the others!




Instructions:

1. Using the picture on the right as a guide, cut the gummy worms to the appropriate relative lengths. Make multiple copies of each chromosome type.
2. Carefully melt chocolate in a microwave, and dip about half of chromosomes 2, 3, and X into chocolate. It is not necessary to coat chromosome 4 or the Y chromosome with chocolate since we will not be evaluating genes on those chromosomes. After dipping, let the gummy worms cool.
3. Arrange two of each chromosome type on paper to make a fruit fly karyotype. If the chromosome is dipped in chocolate, this indicates that it contains a dominant trait. If the chromosome is not dipped in chocolate, this indicates a recessive trait.
4. See if another person can use the key below to decipher the traits that your gummy worm karyotype represents! Then color in the fly correctly.
5. Rearrange the gummy worms to create another karyotype. Repeat the exercise again.

Karyotype Key




- XX chromosomes = female
- XY chromosome pair = male
- Dominant trait in X = red eyes
- Recessive trait in X = white eyes
- Dominant trait in 2 = functional wings
- Recessive trait in 2 = miniature wings (flightless)
- Dominant trait in 3 = yellow body color
- Recessive trait in 3 = ebony or black body color

FOR EXAMPLE:





This karyotype matches a female red-eyed, flying, yellow fruit fly.



There are many different ways to arrange a fruit fly puzzle! Here are a few examples of potential karyotypes and answers.




Color and describe the fruit fly that matches this karyotype! Don't forget to draw in the wings.



Color and describe the fruit fly that matches this karyotype! Don't forget to draw in the wings.



Color and describe the fruit fly that matches this karyotype! Don't forget to draw in the wings.




Hands-on Science Project

GUMMY WORM MEIOSIS

MATERIALS:



In this activity, the cell inherited **chocolate-covered** gummy worm chromosomes from one parent and **non-chocolate** chromosomes from the other parent.



Meiosis with gummy worms

1. Interphase: Arrange 4 pairs of gummy worms on a plate just as in the karyotype activity. One set of gummy chromosomes should be covered in chocolate. The others should be uncovered. Prepare for duplication by making sure you have an exact match for each gummy worm on the plate.


2. Prophase I: When duplicated, the identical chromosomes are linked together and called sister chromatids. Represent this step by adding an identical gummy worm to each gummy worm on the plate.

3. Recombination: During prophase one, the matching chromosomes (homologous pairs) come together to form a tetrad. While they're together, they swap pieces of DNA! With your knife, cut pieces from two matching chromatids and trade them!

4. Metaphase I: Line up all of the tetrads in a straight line and lay out the yarn or noodles and candies to represent the spindle fibers and centrioles!

5. Anaphase I: The spindle fibers pull the homologous chromosomes apart. Each side should have 8 gummy worm pieces in total.

6. Telophase I and cytokinesis: The cell splits to become two cells, each with 8 gummy worm pieces in total. Notice that now there is more variety with the chromosomes due to recombination.



7. Prophase II: A new set of spindle fibers are prepared.

8. Metaphase II: The chromosomes line up in a straight line. Centrioles form and spindle fibers attach.

9. Anaphase II: The spindle fibers pull the sister chromatids apart to opposite sides of the cell.

10. Telophase II and cytokinesis: Each cell splits into two cells, with one copy of each chromosome. Notice how the cells produced (gametes) have half the number of chromosomes as the starting cell.


Why do the gametes have half the number of chromosomes? What is the purpose?

The gametic cells are haploid ($\frac{1}{2}$ the number of chromosomes) so that they can combine with another haploid cell to produce a new diploid cell. The purpose is to increase genetic diversity!

Can any of the 4 daughter cells or gametes produced by meiosis be identical? Why or why not?

The daughter cells of a single mitotic division will not be identical.

Recombination and the random orientation of the chromosomes in metaphase I and II create a high amount of diversity in the daughter cells/gametes.




What happens if, during anaphase II, a spindle fiber pulls two chromatids to one side instead of pulling the chromatids apart?

One of the gametes will have an extra chromosome. During the reproductive process, this cell would result in a trisomy (3 copies of one chromosome). In most cases, trisomy is fatal.

There are only a few chromosomes where a trisomy can develop and then be born.

The Genetics of PKU

Most people have an enzyme that can change the amino acid phenylalanine into tyrosine. But some people inherit a rare genetic disease called phenylketonuria or PKU. This disease is recessive. It's caused by a mutation that codes for a non-functional copy of an enzyme called phenylalanine hydroxylase. The working enzyme converts the amino acid phenylalanine to tyrosine. If phenylalanine cannot be converted, it can build up to toxic levels in the blood and cause permanent damage to nerve cells.



BEFORE NEWBORN SCREENING

A person born with PKU was healthy at birth, but as they grew up they developed permanent damage to brain tissue and experienced intellectual disability, psychiatric disorders, and seizures.

AFTER NEWBORN SCREENING

A person born with PKU is put on a special low-protein diet that restricts the amount of phenylalanine. As long as the diet is followed, they do not experience the negative effects of PKU.




Some people think the mutations responsible for PKU originated in Europe. Does this data support or contradict that idea?

COUNTRY	INCIDENCE OF PKU*
United States	1 in 25,000
Austria	1 in 5,764
Finland	1 in 112,000
France	1 in 9,091
Italy	1 in 4,000
Japan	1 in 125,000
Philippines	1 in 116,006
Thailand	1 in 227,273

supports





contradicts

* From The Genetic Landscape and Epidemiology of Phenylketonuria by Alicia Hillert et al. Published in the American Journal of Human Genetics. Vol 107, Aug 6, 2020.



Aspartame is an artificial sweetener that is 200 times sweeter than sucrose. Why are food products with aspartame required to contain a warning label about phenylalanine?

Because foods containing aspartame will be broken down into phenylalanine. High amounts of phenylalanine are dangerous to people with PKU. Because they cannot convert it into tyrosine, it can build up to toxic levels in the bloodstream.




Transcription - from DNA to RNA

FILL IN THE BLANKS USING THESE WORDS:

membrane DNA adenine transported uracil

In human cells, DNA is kept in the nucleus. Because it is surrounded by the nuclear membrane, other parts of the cell (like ribosomes) can't access it directly. The information in DNA is transported between the nucleus and the rest of the cell by a single-stranded molecule called RNA. RNA is very similar to DNA, but instead of thymine, it contains a nucleotide called uracil. The four units of RNA are adenine, uracil, cytosine, and guanine, or A,U,C, and G.



Translation


FILL IN THE BLANKS USING THESE WORDS:

translation RNA ribosomes codons protein

RNA molecules called mRNA are used by ribosomes to make protein. The ribosome reads every three nucleotides from the strand of RNA and matches them to an amino acid. These three-letter segments of RNA are called codons. For example, if the ribosome reads a codon of three guanines (GGG), it will add the amino acid glycine to the protein it is making. Since the sequence of nucleotides is being translated into a sequence of amino acids, this process is called translation.

LABEL EACH BOX IN THE DRAWING BELOW USING THESE WORDS:

nucleotide RNA ribosome amino acid protein




Blood Types

FILL IN THE BLANKS USING THESE WORDS:


recessive codominant type proteins antigens AB

Blood Type A+



Red blood cells have small proteins and carbohydrates on the surface of their cells, which are called antigens. Different antigens result in different blood types.


Blood Type B-



Blood type is essential to know for performing surgery and other medical procedures. The most well known blood type systems are ABO and Rh.

The ABO phenotype is controlled by one of three alleles: A, B, or O. The A and B alleles are codominant. Someone with both alleles has blood type AB. The A and B alleles are both dominant over O.

Blood Type O+














The Rh gene has a dominant allele (+) and a recessive allele (-). The dominant allele produces a small protein on the surface of the blood cells. Someone with this protein is called "Rh positive." Two copies of the recessive allele results in cells with no Rh protein on their surfaces, a condition which is called "Rh negative."

Decipher the blood type!

Use the information in the antigen key to label each blood type.

Antigen Key

A antigen	♀
B antigen	♂
Rh factor	Y



Genotype & Phenotype

HOW DO WE SHOW BLOOD TYPE WHEN THERE ARE THREE ALLELES?



USE CAPITAL "I" FOR THE A AND B ALLELES WITH A SUPERSCRIPT TO TELL THEM APART. FOR THE O ALLELE, WE USE A LOWERCASE "i".




Blood Alleles
 I^A, I^B, i
 $ii \rightarrow$ Type O
 $I^A I^B \rightarrow$ Type AB
 $I^A i \rightarrow$ Type A
 $I^B i \rightarrow$ Type B


UGH! THAT'S CONFUSING. WHY CAN'T WE JUST USE THE LETTER "O" FOR THE TYPE O ALLELE??



WELL, "O" ISN'T A GOOD LETTER TO USE FOR ALLELES BECAUSE IT CAN BE HARD TO TELL THE DIFFERENCE BETWEEN CAPITAL AND LOWERCASE "O."



SO DON'T CAPITALIZE IT! OR MAKE A NEW LETTER! COMBINE A AND B TOGETHER. ANYTHING IS BETTER THAN USING "I" AND SUPERSCRIPTS.




ANYTHING IS BETTER?? HAVE YOU EVER TRIED TO WRITE A LOWERCASE "A" AND "B" AT THE SAME TIME?




We are using A, B, and O to represent the ABO blood types. Someone with type A blood can either be homozygous (AA) with two alleles that code for enzymes that make the A antigen or they can be heterozygous (AO) with one functional allele. The recessive allele (O) creates a non-functioning enzyme that can't make any antigens.

Phenotype Genotype

Type A	AA or AO
Type B	BB or BO
Type AB	AB
Type O	OO
Rh +	++ or +-
Rh -	--



The phenotypes of a family are displayed below. What are the genotypes of the parents?




The genotype of parent with O+ blood is:

- OO++
 OO+-
 Something else

The genotype of parent with B- blood is:

- BB ++
 BB +-
 BB --
 BO ++
 BO +-
 BO --
 Something else

The phenotypes of a family are displayed below. What are the genotypes of the parents and the child with A+ blood?



The parents are:

- Both homozygous for the Rh factor (both AB++)
 One homozygous, one heterozygous (AB++ and AB+-)
 Both heterozygous for Rh factor (both AB+-)
 We are unable to determine genotype with the given information.

The genotype of the child with A+ blood is:

- AA++ AO+-
 AO++ Either AA++ or AA+-
 AA- Something else

Fill out the Punnett square below for the cross AO⁺ × BO⁺. Include both the genotype and the phenotype. Two squares have been completed as examples. Then answer the questions.

	B+	B-	O+	O-
A+	AB+	AB+	A+	A+
	AB++	AB+-	AO++	AO+-
A-	AB+	AB-	A+	A-
	AB+-	AB--	AO+-	AO--
O+	B+	B+	O+	O+
	BO++	BO+-	OO++	OO+-
O-	B+	B-	O+	O-
	BO+-	BO--	OO+-	OO--

How many different blood types could result from this cross?

Eight different blood types: AB+, AB-, A+, A-, B+, B-, O+, O-

Which blood type is most likely from this cross?

AB+, A+, B+, and O+ are all equally likely (3/16 chance of each)

What is the chance of heterozygous A+ / B+ parents producing a child with O- blood?

1/16

Not all blood types are equally likely! Use the distribution chart for human blood types to fill in the blanks below. As a bonus question, do you know what blood type is called the “universal donor”?


The most common blood type is O+.

The least common blood type is AB-.

Rh positive blood types are more common than Rh negative blood types.


The universal donor blood type is O-.

Distribution of Blood Types



ARE THESE FACT OR FICTION? Write your verdict below each statement:

Cows can have more than 800 possible blood types.



FACT


Mosquitoes prefer type A blood over all other types.

EVERYONE COME AND GET THIS HUMAN RIGHT HERE!

THEIR BLOOD IS A+!

FICTION - they prefer type O or B, depending on the species. One study found that mosquitos were 2x more likely to bite someone with type O blood than type A. In another study, a different species showed a strong preference for type B. See this article for more: <https://www.nature.com/articles/s41598-021-03765-z>

One pint of blood contains 2.4 TRILLION red blood cells!



FACT- For more general reading on blood cells, see: <https://scienctrek.org/scienctrek/topics/blood/facts.cfm>



Hands-on Science Project

WHAT'S YOUR BLOOD TYPE?




Using a blood type kit requires the use of a lancet to provide a blood sample, so this should only be done while supervised and with an adult's permission. Be sure to follow the directions of the blood typing kit used. If not using a blood type kit, decipher the results on the following page to determine the mystery blood types.

MATERIALS:



Blood type kit
Kits typically contain a card, antiseptic, lancet, dropper, sticks or swabs for mixing, and instructions.



Glass of water

A clock or timer









INSTRUCTIONS:

Always follow the directions from your blood typing kit! Most likely, they will look something like this:

1. Place a drop of water on each circle located on the card to activate the serum.
2. Prep your finger by sterilizing it with an alcohol wipe.
3. Prick your finger using the lancet. Then gently massage your finger to allow a drop of blood to form.
4. Touch one of the provided sticks to the blood. Then rest it on one of the circles on the card. Stir gently to mix.
5. Using a new stick each time, repeat until all circles are filled.
6. Gently mix the blood sample with the water on each serum circle.
7. Let the card sit for at least 1 minute. The circle(s) that have agglutinates or clumping indicate your blood type. No clumping in the A or B circles means you are type O. The "Anti-D" or "Anti-Rh" space will clump if you are Rh positive.
8. Using the alcohol wipe, clean your finger and apply a bandaid.
9. Write your name and blood type on the card. Then, when the blood has dried, place the protective sheet on top to save your results.

WHY IS THERE A 'CONTROL' CIRCLE?

If a person is careless, they can cross-contaminate their sample. This might give a result of AB+ when in reality, the blood type should be A-, B-, B+, or A+. To help avoid this error, a control spot is added. If agglutination is visible in the control circle, the results are invalid! The test must be run again to determine blood type.




HOW IT WORKS


The serum on the card in the testing kit contains A, B, and Rh antibodies. When blood is mixed with these antibodies, the antigens attempt to find their match.

When you see the agglutination in your sample, you are witnessing the antigens and antibodies bonding together, thus making it visible what the blood type is.

One of the parents in this family does not know their blood type. Is there enough information from the other individuals to determine what their blood type is? Can the genotype be determined as well? Record the blood types indicated by the cards for each offspring in the boxes below.



One of the parents in this family does not know their blood type. Is there enough information from the other individuals to determine what their blood type is? Can the genotype be determined as well? Record the blood types indicated by the cards for each offspring in the boxes below.



Complete a Punnett square of blood genotypes for each cross below. Then write a summary of what blood types you could expect in the offspring. Use language that would be understandable to a non-scientist.

AB+- × OO+-

	O+	O+	O-	O-
O+	AO++	AO++	AO+-	AO+-
B+	BO++	BO++	BO+-	BO+-
A-	AO+-	AO+-	AO--	AO--
B-	BO+-	BO+-	BO--	BO--

3/8 (or 6/16) will have A+ blood.

3/8 (or 6/16) will have B+ blood.

1/8 (or 2/16) will have A- blood.

1/8 (or 2/16) will have B- blood.

AO-- × BO+-

	B+	B-	O+	O-
A-	AB+-	AB--	AO+-	AO--
A-	AB+-	AB--	AO+-	AO--
O-	BO++	BO--	OO+-	OO--
O-	BO++	BO--	OO+-	OO--

Each blood type (AB+, AB-, A+, A-, B+, B-, O+, and O-) will show up 1/8 (or 2/16) of the time.

AO+- × BB--

	B-	B-	B-	B-
A+	AB+-	AB+-	AB+-	AB+-
A-	AB--	AB--	AB--	AB--
O+	BO+-	BO+-	BO+-	BO+-
O-	BO--	BO--	BO--	BO--

1/4 (or 4/16) will have AB+ blood.

1/4 (or 4/16) will have AB- blood.


1/4 (or 4/16) will have B+ blood.

1/4 (or 4/16) will have B- blood.

MUTATIONS


IN COMIC BOOKS

HOW DID YOU TURN INTO A FLYING SUPER HERO?



IN REAL LIFE

HOW DID YOU DIE?




FILL IN THE BLANKS USING THESE WORDS:

growing replicates enzymes sequence


DNA replication is an essential part of living and growing. It is happening right now inside millions of cells in your own body! Every time a cell replicates, DNA polymerase and other enzymes are there to duplicate the DNA. If something damages or breaks the DNA, repair enzymes are there to fix it. These enzymes are incredibly accurate, but sometimes they make mistakes. When that mistake causes a change to the DNA sequence, we call it a **mutation**.

CAUSES OF MUTATIONS




Radiation

High energy particles from sun, x-rays, and radioactive materials cause damage to DNA.




Random Chance

Something goes wrong in the copying process without getting corrected.




Mutagens

Chemicals that damage DNA or increase the likelihood of mutations




Viruses

Some viruses insert their own DNA into the genome of host cell. (This DNA then becomes part of the host DNA and is passed on to the next generation.)



Is the mutation that causes lactase persistence harmful, neutral, or beneficial? Support your answer with an example.

The ability to digest lactose is beneficial. It allows people to consume another food source that would not otherwise provoke gastrointestinal distress. During a famine, lactase persistence provides a survival advantage.



Lactose - the sugar in milk!
(galactose + glucose)



Lactase - the enzyme that can cut lactose in half


ARE THESE FACT OR FICTION? Write your verdict below each statement:

Hydrogen peroxide produces molecules called free radicals that damage DNA.




FACT

Getting a sunburn increases the number of mutations in your DNA.




FACT - Sunlight contains UV-radiation that can harm skin cells and DNA.

Some people have genes that let them smell chemicals in urine after someone has eaten asparagus.





We'll say **FACT**, but recent research shows it's more complicated than we previously thought. <https://udel.edu/~mcdonald/mythasparagusurine.html>

Nature & Nurture



Three genetically identical dandelion seeds landed in three different environments. Match each plant to its environment. Then note one or two characteristics that gave clues to the match.




- Drought is a stress for plants. often grow taller and have elongated stems.
- It should match with the environment that has the most water.

Write each attribute in the Venn diagram below to classify it as being primarily the result of genetics, environmental conditions, or both. There may not be a clear answer for every trait. Be able to justify your classifications.

eye color	dialect	blood type	intelligence
height	hearing	hair color	honesty

We selected the circles that made most sense to us, but many of these traits can be argued to be the result of both!



List five examples of how the growth of an organism can be influenced by its environment:


- There are many possible correct answers!
- 1 Good nutrition helps organisms grow larger.
 - 2 Low levels of light cause plants to grow taller with elongated stems.
 - 3 Fluoride in the water supply can make teeth stronger
 - 4 Poor soil will stunt the growth of plants.
 - 5 Lack of calcium in the diet can lead to weaker bones.

FILL IN THE BLANKS USING THESE WORDS:


dandelions epigenetics both genetic

The traits plants, animals, and other life forms have are often influenced by both their genetics and their environment. Genetic clones such as dandelions can be almost unrecognizable when grown in drastically different environments. Some traits, like perfect pitch, need a combination of genetic factors and experience to develop the trait. And sometimes the environment can directly influence how genes are expressed! The study of how environmental conditions impact gene expression is called epigenetics.


ARE THESE FACT OR FICTION? Write your verdict below each statement:



FALSE - but average height has increased by half that amount. For more, see <https://ourworldindata.org/human-height>



FACT - Cold Spring Harbor Laboratory has a good article with more information: <https://www.cshl.edu/the-non-human-living-inside-of-you/>



FACT - They're called human chimeras. This rare condition can occur when a dizygotic twin dies in utero and the other embryo absorbs the cells. Recipients of organ transplants are another example of a person who contains two different sets of DNA. For more information see: <https://www.sciencedirect.com/science/article/pii/S1769721220302895>

Hands-on Science Project

APPLE VARIETY TASTE TEST

MATERIALS:




5 apples of different varieties



marker



knife



5 plates

Prepare the taste test

1. Go to the store and pick out 5 apples that are different varieties, including one Red Delicious. If possible, choose apples that have stickers or mark them in some way to differentiate which variety is which.

2. At home, use a marker or paper to label each plate with a letter: A,B,C,D,E. Place the sticker from each apple or a label with the name of each apple on the bottom of each plate to help keep track of which apple is which. Be sure the label with the apple variety is not visible to the person taste-testing the apples.

3. For the browning test, cut each apple in half. Place one half face-down on the plate for future taste testing. Cut a piece from the other half and leave it face up and exposed to the air. Then set a timer. Rate each apple variety for browning at 1, 5, 15, and 30 minutes. The taste test can proceed while the apples are being tested for browning.

4. For the taste test, locate volunteers to taste the apples. Cut additional slices from each apple and present the samples to each taste tester. Each apple sample should be identified by letter (A, B, C, D, and E) not by variety name. If possible, give each taste tester their own paper and have them rate all apples for tartness, sweetness, crispness, and overall flavor.

5. If multiple taste testers evaluated the apples, average their results and then record them on the next page.

The Red Delicious - Beautiful but bland

The Red Delicious apple was created in 1872 at an orchard in Iowa. It debuted as a sweet, round, red and yellow fruit named Hawkeye. When Stark Nurseries bought the rights to the apple in the 1890s, they changed the name to Stark Delicious and then later Red Delicious to match another Stark Nursery apple named Yellow Delicious. As the apple became more popular in grocery stores, growers increasingly selected genes for longer shelf life and appearance over flavor, which resulted in a less appetizing fruit. The selection of redder skin caused deselection of flavor because the genes that produced yellow stripes on the original fruit were on the same chromosomes as those of the flavor-producing compounds. Breeding for uniform appearance and shelf life resulted in a thicker skin. One hundred years later, in the 1980s, demand for the Red Delicious declined as other varieties of more flavorful apples entered the market.



For BEST results

Remove the peel from the apple samples so taste testers don't have peel color as a clue about which variety is which!

TARTNESS

how sour the apple is

	A	B	C	D	E
Very tart					

Results will vary! For a more scientific approach, we recommend having 5 volunteers participate in the taste test and then averaging their results. Be sure to remove the stickers or labels from the apples and just call them by their letter designation (A, B, C etc.). You don't want the name of the variety doesn't influence the taster as they rank the apple.

Individual apples can vary in their texture and taste. In general, Granny Smith will be the most sour or tart apple, and Honey Crisp, Fuji, or Gala will be the sweeter varieties.

Red Delicious, Macintosh, and Golden Delicious will be less crisp, or softer, than other varieties. They also typically rate less tart than the aforementioned varieties.

Very sweet

	A	B	C	D	E
Very sweet					

Sweet

	A	B	C	D	E
Sweet					

A little sweet

	A	B	C	D	E
A little sweet					

Not sweet

	A	B	C	D	E
Not sweet					

CRISPNESS

how crisp the apple is

	A	B	C	D	E
Very crisp					

Crisp

	A	B	C	D	E
Crisp					

A little crisp

	A	B	C	D	E
A little crisp					

Not crisp

	A	B	C	D	E
Not crisp					

OVERALL FLAVOR

the most subjective rating of all

	A	B	C	D	E
Delicious flavor					

Good flavor

	A	B	C	D	E
Good flavor					

Okay flavor

	A	B	C	D	E
Okay flavor					

Poor flavor

	A	B	C	D	E
Poor flavor					

BROWNING

how quickly the apple turns brown after being cut

	A	B	C	D	E
Browning visible at 1 min					

Browning visible at 5 mins

	A	B	C	D	E
Browning visible at 5 mins					

Browning visible at 15 mins

	A	B	C	D	E
Browning visible at 15 mins					

Little to no browning visible at 30 min

	A	B	C	D	E
Little to no browning visible at 30 min					

DID YOU KNOW? Apples turn brown after being cut because of a reaction between oxygen in the air and enzymes in the apple called polyphenol oxidases. Applying vitamin C to apple slices (such as with lemon juice) will stop this reaction. But different apple varieties have different amounts of polyphenol enzymes. The Opal variety has such low levels, it's known as a "non-browning" apple.

X-linked traits

FILL IN THE BLANKS USING THESE WORDS:

more inherited chromosomes linked recombination


Genes located on different chromosomes are inherited independently. Genes that are further apart on the same chromosome are also inherited independently because of recombination. But if genes are close together on a chromosome, there's a very good chance they'll be inherited together. We call them linked traits. There's one chromosome pair in particular where linked traits are rather famous: that's pair 23. Genes on the X chromosome are called X-linked traits, and they are more likely to be expressed in men than in women.

Red-green colorblindness


Human vision comes from specialized cells called rods and cones. These photoreceptors work by detecting specific wavelengths of light. The genes with the information to create the color-sensitive red and green cones are located on the X-chromosome.

In people of Northern European ancestry, red-green colorblindness, also called protanopia (no red cones) or deutanopia (no green cones), is present in about 8% of males and 0.6% of females.


Tritanopia (no blue cones) is not x-linked. It arises from a mutation on chromosome 7 and is much rarer than red-green colorblindness.



Identify the biological sex and color vision status for each of the children below. Write "male" or "female" and either "WT" for wild type (normal vision) or "CB" for colorblind.



What proportion of the offspring will be colorblind from a colorblind mother and trichromat (normal vision) father?



On average, 1/2 of the offspring will be colorblind (the male half). All of the females will have trichromat vision but be carriers for colorblindness.

About Calico Cats


In cats, the genes that control orange or black fur color are on the X chromosome. Orange is dominant, and black is recessive. Because it can be difficult to distinguish between upper and lowercase versions of the letter "o," the alleles are typically represented by X^B (orange) and X^b (black).

In females with an XX genotype, early in development, one of the X chromosomes is inactivated and supercoiled into a dense structure known as a Barr body. None of its genes will ever be expressed!

The random inactivation of half of the X chromosomes early in development is called X-inactivation. As a result of this inactivation, some of the cells in XX females express genes from one chromosome while others express entirely different genes!

With calico cats, you can see where the Barr bodies formed because the black fur is from one X chromosome and the orange fur is from another. Calico cats are always female, except in the rare case of Klinefelter syndrome (XXY).

The white fur in a calico cat is caused by the recessive piebald gene, which is located on a different chromosome. It creates unpigmented (white) patches of fur.



What will be the outcomes of the following crosses? Will any of them produce a calico cat?
All cats carry the piebald gene and have patches of white fur.

Black and white "tuxedo" male (X^bY) with an orange and white female (X^BX^b).

All the female cats will be calico (about $\frac{1}{2}$ the cats, on average).

X^b	Y
X^BX^b CALICO	X^BY ORANGE
X^BX^b CALICO	X^BY ORANGE

Orange and white male (X^BY) with a calico female (X^BX^b).

$\frac{1}{4}$ chance of a calico kitten

X^B	Y
X^BX^B ORANGE	X^BY ORANGE
X^BX^b CALICO	X^bY BLACK

Black and white male (X^bY) with a calico female (X^BX^b).

$\frac{1}{4}$ chance of a calico

X^b	Y
X^BX^b CALICO	X^BY ORANGE
x^bx^b BLACK	x^bY BLACK

Orange and white male (X^BY) with a black and white female (X^bX^b).

All females will be calico, so, on average, $\frac{1}{2}$ of the kittens will be calico.

X^B	Y
X^BX^b CALICO	x^bY BLACK
X^BX^b CALICO	x^bY BLACK

Genetically Modified Organisms


WHAT THEY ARE

The genetics of organisms are changing all the time due to natural factors such as mutations and sexual reproduction. But these changes have a high degree of randomness. Genetically modified organisms, or GMOs, have had their genetic code changed in a more deliberate way. They have either had a specific piece of DNA inserted or removed from the genetic code.


Sometimes this piece of new DNA is from the same species. For example, a pumpkin could be genetically engineered to have another trait by taking the gene from a different breed of squash and inserting it into the pumpkin genome. This would be faster than achieving the same result through artificial selection or breeding.

More often, the piece of new DNA is from a different species, such as when human beings engineer bacteria to produce the protein insulin to treat diabetes.

Artificial Selection




Genetic Engineering




EXAMPLE 1: Insulin-producing bacteria

Insulin is a hormone that regulates the amount of glucose in the bloodstream. Someone with diabetes is unable to produce enough insulin. Before 1922, diabetes was a fatal illness. After the discovery of insulin, doctors used insulin from the pancreases of pigs or cattle to treat diabetes. This was an improvement from before, but it would sometimes cause allergic reactions because the insulin in humans is different from the insulin from pigs and cows. In 1978, the bacteria *E. coli* was modified by inserting the human gene for insulin into its DNA. When this bacteria reproduces, it transfers the insulin production gene to its millions of offspring. The insulin protein is then purified and used as medicine.



EXAMPLE 2: Bt corn

Bacillus thuringiensis (Bt) is a common bacteria that has natural insecticidal abilities. It makes a protein known as the **Bt toxin** which is poisonous to most insects. Because of this ability, Bt has been an important pest control tool for nearly a century. In the 1990s, scientists were able to develop bug-resistant plants by moving some of the genes from the bacteria into corn and cotton. Bt crops have been highly effective at combating pests such as European corn borer, rootworm, corn earworm, tobacco budworm, and bollworm. They are the predominant varieties planted in North America. But now the bugs are starting to develop resistance!



Benefits: The new insulin is the same as human insulin and easier to produce.

Drawbacks or concerns: Almost none

Alternatives: The only alternatives are to use the insulin produced by the pancreases of other mammals. But pig and cow insulin is slightly different than human insulin and can cause allergic reactions.

EXAMPLE 3: GURT or Terminator Seeds

Genetic use restriction technology (GURT), also known as terminator seeds or suicide seeds, involves a genetic switch that will be activated to produce a non-viable seed. Advocates of GURT technology say it would benefit farmers by reducing volunteer plants that come up from the previous year after crops have been rotated. Opponents of GURT technology say the only benefit is that a seed company would earn more money because farmers could not save seeds to plant again the following year. Higher food prices and the risk of the terminator gene impacting other populations of plants were of strong enough concern that the United Nations recommended a moratorium on the sale of terminator seeds. India and Brazil have both passed laws banning GURT technology.



EXAMPLE 4: Genetically Modified Mosquitos

The *Aedes aegypti* mosquito spreads diseases such as malaria, dengue fever, Zika, and chikungunya. But only female mosquitoes bite. They need a blood meal in order to produce eggs. Scientists developed genetically modified *A. aegypti* mosquitoes that had a **self-limiting gene**. This gene prevents female mosquitoes from growing to adulthood. The genetically modified mosquitoes also have a fluorescent marker gene that glows under a specific light so that they can be identified in the wild. These mosquitoes have been successfully used in Brazil and Panama to reduce the number of *A. aegypti* mosquitoes and the diseases they cause.



Benefits: The seed producing company makes more money. There will not as many volunteer crops during rotation.

Drawbacks or concerns: Increased food prices, loss of biodiversity.

Alternatives: Don't develop terminator seeds! Harvest crops in a timely manner to discourage seeds falling the field.

Benefits: Fewer mosquitoes means fewer diseases.

Drawbacks or concerns: Other animals depend on mosquitoes for food.


Alternatives: Draining marshy land, nets, spray, and treating sick people to reduce disease. Other GMO efforts focus on making it so the mosquitoes can no longer carry certain diseases.

If you could design a genetically modified organism, what would you design and why?

Answers will vary!

I would design a type of algae that would digest microplastics to help remove microplastics from the oceans.

SALT is not an O!
It has no Gs to M!




Ridiculous label.

THINK ABOUT IT: One of the biggest questions of the upcoming century is whether or not human beings should modify their own genetic code. Using CRISPR and gene therapy, it is possible to alter genes to make a person taller, smarter, or less susceptible to disease.

Should people experiment with this technology, or should genetic modification to human beings be restricted? What do you think?

Genetics Crossword Puzzle



VERTICAL Words

1. A cell organelle where protein synthesis occurs
2. A type of molecule that combines to form proteins
3. A sequence of three nucleotides
4. The complete set of genes or genetic material present in a cell or organism
5. The process of making an RNA copy of a DNA sequence
6. The process of creating haploid cells containing half of the genetic information


HORIZONTAL Words

7. The process of converting messenger RNA to a sequence of amino acids
8. A version of a gene inherited from a parent
9. When the sequence of a gene changes
10. A protein that provides structural support to a chromosome
11. Ribonucleic acid
12. The rearrangement of genetic material by crossing over in chromosomes


DNA MAZE

Collect all the building blocks of DNA to escape the maze. Start at the star and work your way through the maze to escape in the upper right corner. You can only pass through a lock if you have the corresponding key .

Keep an eye out for overpasses and underpasses!



Follow the yellow path to get the **A key**. Then backtrack to the pink path to unlock the **A lock** and get the key labeled **C**. Then follow the pink back to where it intersects with the salmon-red color to access the **C lock**. Continue following the path to get the **G key**. Then turn around and backtrack to the pink and then to the yellow path from the beginning. Go past the **A key** to get on the green path, which will take you to the **G lock**. Right after the **G lock**, you'll get the **T key**. Then backtrack all the way back to the yellow and then the pink. Next, get back on the salmon-red color path to access the blue path. The blue path will take you to the **T lock** so you can escape the maze!



Quiz Time!

ANSWER THE QUESTIONS TO
SEE WHAT YOU LEARNED
ABOUT GENETICS!

- 1 Which individuals will have identical or near-identical genes?
 A. Siblings
 B. Identical twins
 C. Fraternal twins
 D. Parents and their offspring

- 2 How many pairs of chromosomes does the human genome have?
 A. 23
 B. 46
 C. 15
 D. 52

- 3 A strand of DNA reads 'GATACTACT'. The complementary (paired) strand reads
 A. CTATGATGA
 B. CATACTAGT
 C. ADGCTDCT

- 4 How does RNA differ from DNA?
 A. RNA contains deoxyribose, and DNA contains ribose.
 B. RNA contains uracil, and DNA contains thymine.
 C. RNA is double stranded, and DNA is single stranded.

- 5 Is it possible for a man and a woman, each with Type A blood, to have a child that is Type O?
 A. Yes, if the man is heterozygous
 B. Yes, if the woman is heterozygous
 C. Yes, if both parents are heterozygous
 D. No, it is impossible

- 6 In cells, long pieces of DNA are wrapped up with proteins to form
 A. microtubules
 B. RNA
 C. nuclei
 D. chromosomes

- 7 DNA is made up of which repeating subunits?
 A. RNA
 B. nucleotides
 C. genes
 D. bases

- 8 The sequence of DNA in the skin cells of your hand is significantly different in sequence from the DNA in your heart cells.
 A. True
 B. False

- 9 Down Syndrome results from
 A. an extra copy of chromosome 21
 B. a missing copy of chromosome 21
 C. a specific gene mutation
 D. changes to DNA that occur after birth

- 10 Who are more likely to suffer from X-linked diseases and why?
 A. females, because they have two X-chromosomes and are therefore more likely to inherit one with a disease
 B. males, because they have only one X-chromosome and it, alone, determines whether they have the disease.
 C. females, because one of their X-chromosomes is deactivated
 D. males, because the Y-chromosome already carries the gene for the disease.

- 11 Crossing over or recombination takes place during _____.
 A. meiosis
 B. mitosis
 C. segregation
 D. linkage

- 12 A sequence of 3 consecutive nucleotides that codes for a specific amino acid or signals the start or termination of gene is called a(n)
 A. gene
 B. allele
 C. codon
 D. gamete

- 13 Two parents have 8 kids, and all 8 kids have different phenotypes for their blood type. Their blood types are A+, A-, B+, B-, AB+, AB-, O+, and O-. What genotypes do the parents have for blood type?


One parent must have blood type AO+ and the other has type BO-. No other combinations can yield all eight blood types as offspring.

- 14 Can two orange cats produce a calico kitten? Why or why not?

They cannot. Both black and orange alleles are needed for a calico kitten. An orange male cat only carries the orange allele. An orange female cat has two orange alleles.


- 15 If the mother carries the allele for red-green colorblindness and the father has normal vision, what proportion of their offspring will be colorblind?

- A. Half the males, all the females
 B. Half the males, none of the females
 C. All the males, half the females
 D. All the males, none of the females
 E. None of the males, none of the females



- 16 If the father has red-green colorblindness and the mother has normal vision and is not a carrier, what proportion of the offspring will be colorblind?

- A. Half the males, all the females
 B. Half the males, none of the females
 C. All the males, half the females
 D. All the males, none of the females
 E. None of the males, none of the females



- 17 True or false: in a living organism, mutations are always occurring.

- A. True
 B. False

- 18 Lactase persistence is an example of a _____ mutation.

- A. beneficial
 B. neutral
 C. harmful

- 19 If the parents are A+ and B+ and are both heterozygous for both alleles, what are the chances of their child having the universal donor blood type of O-?

To answer this question you can draw a Punnett square or multiply the individual probabilities:
 A. 1/2
 B. 1/4
 C. 1/8
 D. 1/16
 E. 1/32
 For the A+ parent, there is $\frac{1}{2}$ chance of giving O allele and $\frac{1}{2}$ chance of giving Rh- allele
 $\frac{1}{2} * \frac{1}{2} = \frac{1}{4}$ chance of the child inheriting O- from the first parent.
 For the B+ parent, there is $\frac{1}{2}$ chance of them giving the O allele and $\frac{1}{2}$ chance of them giving the Rh- allele.
 $\frac{1}{2} * \frac{1}{2} = \frac{1}{4}$ chance of the child inheriting O- from the second parent.

To obtain the chance of the offspring having the genotype OO-, multiply the two probabilities: $\frac{1}{4} * \frac{1}{4} = 1/16$.

- 20 What is the genotype of the shaded individual in the pedigree?


- A. AO-
 B. AO++
 C. AA++
 D. AA-

We know they are heterozygous for both alleles because of the B- offspring.

- 21 What is the genotype of the striped individual in the pedigree?

- A. BO-
 B. BO--
 C. BB+-
 D. BB--

The phenotype of Rh negative is always homozygous recessive (BB--). The B allele came from the father. If the mother (blood type A+) had given the A allele, the blood type would have been AB-. Since the blood type is B-, the mother must have given the O allele.




Invasive vs Endangered

FILL IN THE BLANKS USING THESE WORDS:

fungus adaptable species traits clone seven

New Zealand Mud Snail




Invasive species like the kudzu vine or the New Zealand mud snail have specific traits that make them more prolific and adaptable than other species. These traits are why they are so successful at taking over.

Endangered species like the white fringeless orchid or Oahu tree snails have traits that make them less adaptable or slower to reproduce.


While a kudzu seed can germinate in virtually any soil, orchid seeds can only germinate by bonding with a specific species of fungus. The fungus they need often depends on old growth forests.

A single New Zealand mud snail can clone itself to start a new population, and each of its offspring is born pregnant! The Oahu tree snail, on the other hand, must be almost seven years old before it is mature enough to mate. When it does reproduce, it needs a partner snail, and the pair will only have one to four offspring.

Oahu Tree Snail



Draw lines to match each trait with the corresponding invasive or endangered organism:



This animal only eats carrion. It needs to lay its eggs in a dead bird or rodent in order to reproduce.

Each plant produces thousands of seeds per year. Seeds can remain viable for 28 years.

This animal can eat almost anything. Its venom allows it to attack and eat prey 1,000 x larger than itself!

This animal can produce 150,000 eggs a year. It lives underwater but can survive on land for up to 4 days.

This plant flowers once every 2 to 7 years and is pollinated by nocturnal beetles and flies.

What about in YOUR climate?

Research the ecology of where you live to discover if there are any invasive or endangered species near you. If there are none in your area, research one invasive and one endangered species for a similar climate.

TROPICAL



Every month of the year averages 18°C / 64°F or warmer. Lots of rain.

DRY



Low precipitation. At least one month warmer than 10°C / 50°F.

TEMPERATE



Temps in coldest month do not average below freezing. The warmest is above 10°C / 50°F.

CONTINENTAL



Coldest month is below freezing, warmest is above 10°C / 50°F.

POLAR



Temperatures for every month of the year average below 10°C / 50°F.

INVASIVE SPECIES

Common name: _____

Scientific name: _____

Average number of offspring produced in one year: _____

Preferred diet or growing conditions: _____

What native species is it competing with? _____

What traits or characteristics contribute to this species being invasive?

There are many different options to choose from! Research a species specific to where you live. A few invasive species in the US and Canada include: Zebra Mussels, Asian Carp, Burmese Python, Purple Loosestrife, Fire Ants, Argentine Ants, Kudzu Vine, Cane Toad, Rusty Crayfish, Tamarisk, Sea Lampreys, Emerald Ash Borer, and Leafy Spurge.

ENDANGERED SPECIES

Common name: _____

Scientific name: _____

Average number of offspring produced in one year: _____

Preferred diet or growing conditions: _____


What traits or characteristics contribute to this species being endangered?

There are also several options to choose from in this category! Research a species specific to your area.

A few endangered species in the US and Canada include: Devil's Hole Pupfish, Oahu Tree Snail, Kemp's Ridley Sea Turtle, Florida Panther, Dusky Gopher Frogs, California Condor, and Ozark Hellbender (a salamander)

Darwin and a journey around the world


In 1831, Charles Darwin secured passage on a ship conducting a mapping expedition that would circle the globe. The captain was interested in science and had more than 400 books onboard.




After visiting South America, the ship sailed to the Galapagos Islands.

In a few days' time, the Beagle will sail for the Galapagos Islands. I look forward with joy and interest to this, both as being somewhat nearer to England and for the sake of having a good look at an active volcano!
- Charles Darwin


Darwin was fascinated by the diversity of life between the different islands. It was possible to tell which island a tortoise came from just by looking at the shape and patterns of the shell!




Santa Cruz was a green island with abundant food for tortoises such as grasses. The tortoises on this island had domed shells.




España was an arid island. The tortoises on this island had saddle-backed shells and long necks.



The islands also had iguanas, which Darwin described as "disgusting imps of darkness."



Darwin noticed that the finches on different islands had differently-shaped beaks. Finches that primarily ate seeds had thicker, stronger beaks. Insect-eating finches had narrower, spear-like beaks.




Hands-on Science Project

MAP THE JOURNEY OF THE H.M.S. BEAGLE


MATERIALS:




scissors




tape or
glue



crayons, colored pencils, or markers




world map and 3-D labels
(templates in appendix)




The Beagle had made one previous trip to South America to survey and map the area. A second journey was planned that would circumnavigate the globe. Charles Darwin secured passage as the ship's naturalist. The journey took 5 years and included many extended stops with exploration expeditions. Darwin wrote frequent letters and kept a detailed diary as they traveled. His observations were instrumental in the famous book he published years later: *On the Origin of Species*.

Map the journey

1. Print and color the world map (pg 120 in the appendix). Then draw the 1831-1836 journey of the H.M.S. Beagle on the map. Use the list of stops and example map below to help chart the course.



2. Match each of the clues on the following page to the circled locations on the world map template.



3. Color the 3-D pop up labels. You can use either the pre-drawn cutouts or create your own (instructions in appendix).

4. Cut and fold each cutout, securing the flaps with glue so that they stand upright.

5. Arrange the 3-D cutouts across the top of the map. Glue them in place and write the corresponding letter or place name next to each cutout.

Selected stops along the way

These are just a few of the stops the H.M.S. Beagle made during its 5 year journey. There were many others!

1. Plymouth, England Dec 27, 1831
2. Cape Verde Islands Jan 1832
3. Punta Alta Sep 1832
4. Concepción Mar 1835
5. Galápagos Sep-Oct 1835
6. Wallerawang Jan 1836
7. Keeling Islands Apr 1836
8. Ascension Island, Jul 1836
9. Falmouth, England Oct 2 1836



7 STOPS MADE BY THE H.M.S. BEAGLE

Draw lines between each description to match it with the correct location. Each description matches a stop made by the H.M.S. Beagle. The pictures on the next page provide additional clues.

It was here Darwin found the fossilized skull of an extinct megatherium (giant ground sloth), which was about the size of an elephant! This skull allowed scientists to assemble a full megatherium skeleton for the first time.

CLIMATE: SUBTROPICAL
CLOSEST OCEAN: ATLANTIC
CLOSEST CONTINENT: SOUTH AMERICA

Darwin encountered marine iguanas, birds, land tortoises, and many other creatures. While studying finches, he noticed distinct differences in their beaks that correlated with the differences in their available food.

CLIMATE: HOT AND DRY
CLOSEST OCEAN: PACIFIC
CLOSEST CONTINENT: SOUTH AMERICA

After a large earthquake, Darwin found mussel beds had been raised above the water level. He also found marine shells in the mountains that proved the land had risen over time.

CLIMATE: SUBTROPICAL
CLOSEST OCEAN: PACIFIC
CLOSEST CONTINENT: SOUTH AMERICA

Though the black and red volcanic island had been described as "hell with the fire put out," Darwin saw potential.

He later convinced the British Admiralty to plant trees on the island. A century and a half later, the island is covered in trees and vegetation with a National Park called, "Green Mountain." The forest of planted trees didn't just change the soil on the island, it changed the weather too!

CLIMATE: HOT AND DRY
CLOSEST OCEAN: ATLANTIC
CLOSEST CONTINENT: AFRICA

Here Darwin saw a platypus and many other marsupials. Despite how different these marsupials were from the animals in England, he observed that they had similar behaviors and filled the same ecological niche as the mammal species back home.

CLIMATE: TROPICAL
CLOSEST OCEAN: PACIFIC
CLOSEST CONTINENT: AUSTRALIA

At these islands, Darwin saw rock made from shells and coral sandwiched between layers of volcanic rock. The sedimentary rock could only form at the beach or underwater, yet here it was high above sea level.

CLIMATE: TROPICAL
CLOSEST OCEAN: ATLANTIC
CLOSEST CONTINENT: AFRICA

Darwin created a theory that coral reefs are formed as volcanic islands gradually subside into the ocean. At these islands, he was able to find evidence to support this theory as he collected many geological, plant, and animal specimens.

CLIMATE: TROPICAL
CLOSEST OCEAN: INDIAN
CLOSEST COUNTRY: INDONESIA

Concepción

Punta Alta

Galápagos Islands

Ascension Island


Cape Verde Islands

Wallerawang


Keeling-Cocos Islands

7 STOPS MADE BY THE H.M.S. BEAGLE


Circle the two locations you would most like to visit!




The Galapagos are part of the country of Ecuador. 97% of the islands belong to a national park and wildlife preserve.




Wallerawang is located 150 kilometers northwest of Sydney near the Blue Mountains.




Of the world's known earthquakes of over 8.5 magnitude, nearly 1/3 have occurred in Chile.




This British territory is a remote Atlantic island located about 1,600 km (1,000 miles) from the coast of Africa and 2,300 km (1,400 miles) from the coast of Brazil.



The Cape Verde Islands (officially named the Republic of Cabo Verde) is a group of volcanic islands located approximately 580 kilometers (360 miles) west of the westernmost point of Africa.



The Territory of Cocos (Keeling) Islands are an Australian territory located in the Indian Ocean. Only 2 of the 27 islands are inhabited.






The city of Punta Alta is located on the Atlantic coastline of Argentina.

What clues from his visits did Darwin use to develop his theories of natural history and evolution?

Darwin found fossils of sea creatures on high mountains that showed shifting geology over time. He saw that plants and animals on different islands had variations that suited their needs for survival on the island, and that organisms with advantageous traits were more likely to survive and reproduce.

Genetic Variation



UNDERLINE THE CORRECT ANSWER TO EACH QUESTION:

Which reproductive strategy is most common among all the species living on Earth?

Asexual Sexual

Which reproductive strategy produces individuals that are genetically identical?

Asexual Sexual

Which reproductive strategy produces the most genetic variation?

Asexual Sexual

Which of the populations above has the highest genetic diversity?

Banana Mud Snail Pigeon

Resources are Limited

Water, nutrients, and food are all finite resources. Consequently, not every individual will survive and reproduce. *But what if every member of a population did?* Use the information that follows to calculate the total mass of octopuses, puffballs, opossums, and elephants. Start with 100 individuals. Assume every individual survives to adulthood and reproduces once and only once. Two generations have been completed as examples.

Pacific Octopus - *Enteroctopus dofleini*

Average mass: 15 kg
Average number of offspring: 100,000

Generation 1: 100 octopuses Note that dividing by 2
½ are female and ½ are male. and then multiplying by
100/2 = 50 breeding pairs
50 × 100,000 = 5,000,000




Generation 2: 5,000,000 octopuses
 $5,000,000/2 = 2,500,000$
 $2,500,000 \times 100,000 = 250,000,000,000$ or 2.5×10^{11}

Generation 3: $=1.25 \times 10^{16}$

Generation 4: $1.25 \times 10^{16} \times 50,000 = 6.25 \times 10^{20}$

Generation 5: $6.25 \times 10^{20} \times 50,000 = 3.125 \times 10^{25}$

Total mass: $(3.125 \times 10^{25}) \times 15 \text{ kg} = 4.6875 \times 10^{26} \text{ kg}$



Star Formation Threshold
FUSION
This much of anything will have strong enough gravity to create a new star!
 $1.6 \times 10^{29} \text{ kg}$

Mass of the Sun
 $2 \times 10^{30} \text{ kg}$

North American Opossum - *Didelphis virginiana*

Average mass: 4 kg
Average number of offspring: 16

Generation 1: 100 opossums

½ are female and ½ are male.
100/2 = 50 breeding pairs
50 × 16 = 800

For each 16 offspring,
there will be 8 new
mating pairs.

Generation 2: 800 opossums

$800/2 = 400 \text{ pairs}$
 $400 \times 16 = 6,400$

Generation 3: $6,400 \times 8 = 51,200$

Generation 4: $51,200 \times 8 = 409,600$

Generation 5: $409,600 \times 8 = 3,276,800$

Total mass: $3,276,800 \times 4 \text{ kg} = 13,107,200 \text{ kg} \approx 1.3 \times 10^7 \text{ kg}$

Elephant - *Loxodonta africana*

Average mass = 5,000 kg
Average number of offspring: 1

Generation 1: 100 elephants

½ are female and ½ are male.
100/2 = 50 breeding pairs
50 × 1 = 50

Generation 2: 50 elephants
 $25 \times 1 = 25$

Generation 3: 12 breeding pairs give 12 offspring.

Generation 4: 6 breeding pairs give 6 offspring.

Generation 5: 3 breeding pairs give 3 offspring.

Total mass: $3 \times 5,000 \text{ kg} = 15,000 \text{ kg} = 1.5 \times 10^4 \text{ kg}$

Did any of the populations exceed the mass of the Earth? Yes: puffballs and octopus.

Did any of the hypothetical populations become large enough for the formation of a star? Yes: puffballs

Did any of the populations experience negative growth? The elephant population dropped.

What do you think?

In the wild, the survival rate for newly-hatched octopuses is less than 1%. What would happen if a certain species of octopus had a smaller birthrate but a higher survival rate?

It depends on how much higher and smaller the rates are! If the octopus only had 10 offspring but it stayed with them and took care of them till they were almost full grown, most of that generation would survive to reproduce. A population of "small birthrate but high-survival rate" octopuses might even grow faster than the population of octopuses where each female has 100,000 offspring.

Did the model of elephant population growth reflect what we observe in the real world? Why or why not? How could the model be adjusted to better reflect real-world conditions?

No. In the real world, elephants have more than one offspring because they have more than one pregnancy. The model would be more accurate if it used the average number of offspring an elephant had over a 25 year period, or during its lifetime.


Survival of the fittest

FILL IN THE BLANKS USING THESE WORDS:

traits family mutations pressure limited survival

Not all members of a species or a family are the same. Some will have inherited different genes and traits than others, and mutations in DNA can cause further differences. If every individual reproduced, then most traits would be passed on. But because resources are limited, the number of offspring that survive is also limited. Over time, the pressure of limited resources ensures that the traits passed on are those best suited for survival. This idea was first described by Charles Darwin and is called "survival of the fittest." The word "fittest" in this phrase does not necessarily mean strongest or most athletic. Instead, "fit" refers to successful reproduction.

The Influenza Virus



The influenza virus has a high **MUTATION** rate. Small changes to its genetic material occur each time it infects a host.


Two new variants of influenza are described: Variant X has mutated to become more transmissible and can survive on surfaces for 36 hours. Variant Y has mutated to become more fragile. It falls apart 5 minutes after leaving a host. Which variant will become more common, and why?

Variant X will become more common. It lasts longer and spreads more easily, so more people will catch Variant X and transmit it to others. Variant Y is much less likely to spread, and when it does spread, it will be less likely to continue to spread.


Selection

Selection happens when one item or trait is favored over another. For example, Sara buys a bag of trail mix and puts it in the kitchen pantry. Sara's brother Vance only eats the raisins from the trail mix, and then pretty soon the trail mix doesn't have any more raisins. They'll all have been "selected" out. This same process occurs in nature too.


Consider a mixed population of mice where half are tan colored and half are black. Identical populations of mice arrive on three different islands. Color the islands according to their descriptions. Then match them with the most likely outcome.




Volcanic island has black rock and hawks that hunt by sight.




Sandstone island has tan rock and hawks that hunt by sight.




Green island has moss-covered rock and no predators.



The mouse population is all tan.




The mouse population is all black.



The mouse population is multicolored (both tan and black).

Case Study: The Peppered Moth

The peppered moth comes in both black and light varieties.



In the countryside, the light grey variety was much more numerous due to its superior camouflage. Its coloring matched that of the tree bark of many trees, allowing it to hide from predators.

In the late 1800s and early 1900s, naturalists noticed that the black peppered moth became more numerous than the lighter variety. The change was especially prominent in areas of London with industrial pollution.

Why was the black peppered moth more common in industrial areas?

The black peppered moth was more likely to survive because it blended in with the dark soot of the industrial areas of London. Predators were more likely to see the lighter moths, so fewer lighter moths survived and reproduced.

Adaptation

FILL IN THE BLANKS USING THESE WORDS:

alligators generations learned beaks climates confuse characteristic

An adaptation is a characteristic that helps an organism survive in its habitat. It takes many generations for adaptations to develop. Birds have adapted different kinds of beaks that help them obtain their particular source of food. Alligators have eyes and nostrils placed on top of their heads, so they can stay mostly submerged. Spots and stripes can camouflage animals or confuse predators. In nearly all climates, plants and animals have adapted to survive and thrive. Learned behaviors are not considered adaptations.

Name a plant or animal that has the indicated adaptation:

Whiskers to help them feel their way through tight spots

Rat

Lives in large herds to protect against predators

Wildebeest

Webbed feet for faster swimming

Duck

Fur that shifts color with the seasons

Rabbit

A look-a-like of a venomous or poisonous animal

King snake

Acute eyesight to see prey at a distance

Peregrine falcon

Hibernation in cold seasons

Groundhog

Camouflage to blend in with surroundings


Chameleon

Toxins excreted through their skin

Poison dart frog


ARE THESE FACT OR FICTION? Write your verdict below each statement.

Some frogs freeze solid in the winter, then thaw out in the spring.



Fact - Wood frogs do exactly this!


The chameleon monkey can change its skin to match its surroundings.



False. No such monkey exists.

Some salamanders have no lungs. They breathe through their skin!

MOUTHS ARE ONLY FOR EATING.







Fact - the arboreal salamander and the California slender salamander, don't have lungs or gills as adults.

Evolution

Evolution is change over time. More specifically, evolution is *descent with inherited modifications*. Mutations are always occurring, which introduces new variations into genetic material. The mutations that have advantages will be selected, and soon the genetic information of a population will be different than it was before. This change is evolution.

Sometimes the change is on a small-scale, such as changes in how often an allele appears in a population. Other times, the change is on a large-scale, such as when different species descend from a common ancestor.

Match each picture with the correct description. Check the box if it is an example of evolution.



Prehistoric horses were grazing animals that walked on four toes. Over millions of years, the fossil record shows a change in the foot structure of the horse, leading to an animal that walks on one toe.

Farmers grew corn with Bt, a pesticide that killed more than 99.9% of corn borer insects. Years later, the farmers discover that Bt is not killing as many insects. Genetic testing shows that many insects have a gene that makes them resistant to Bt.

A mountain range that was once very rugged becomes eroded over time and eventually becomes smaller and covered in trees. Geological evidence also shows that several ancient landslides changed the shape of the mountain range.

A large oak tree experiences many changes each year. It flowers in the spring and produces acorns and lots of leaves. Then in the fall and winter, all of the leaves and acorns fall to the ground. The next spring, the cycle begins again.

Can a single animal evolve? Why or why not?

No. Not according to the biological definition of "evolve." Evolution can only happen through descent with modification, which requires multiple generations.

Can a population of animals evolve? Why or why not?


Yes, but only over time across generations. Genes for new traits can arise through mutations or the proportions of certain genes can change over time either through natural selection or through random processes. This is evolution.

Misconceptions about evolution


Here are three common misconceptions about evolution. Can you spot the errors in each idea or think of a counter example? Correct each idea by writing a better explanation in the boxes below.

THE


Humans evolved from chimpanzees.



Evolution always results in stronger and better organisms.



Humans can't permanently damage an ecosystem because species will evolve the traits they need to survive.






YOUR

Chimpanzees did not evolve into humans. Humans and chimps are related because they both evolved from a common ancestor after diverging 12.1 million years ago.

Natural selection results in improved ability to survive and reproduce in a given environment. Evolution selects for "good enough to reproduce" but this is not always the stronger or "better" organism!

ONCE A SHARK, ALWAYS A

The first spiny sharks evolved during the Silurian Period, nearly 450 million years ago! Their cartilaginous skeleton and streamlined shape are still observed in modern shark species. But does that mean sharks have stopped evolving? Not at all! There have been and continue to be many fascinating variations between sharks. Just consider the following:



Stethacanthus, a Carboniferous shark with an anvil-like fin
Megalodon, a Cenozoic giant more than 65 feet long

Rhincodon or whale sharks, a modern filter-feeding shark

Mitsukurina or goblin shark, a modern deep-sea shark

Sphyrna or hammerhead shark, a modern shark with an elongated head structure called a cephalopel


Genetic Drift

About 500,000 years ago, a group of **Canadian geese** (*Branta canadensis*) were blown off course and landed on the islands of Hawaii.

The environment of Hawai'i was different than North America. On Hawaii, the geese enjoyed a more stable temperature, so they didn't have to migrate. They also had fewer predators, making flight less of a priority.


Over time, the coloring changed, and some of the geese lost the ability to fly. Today, the only surviving species of goose on Hawaii is called the **nēnē** (*Branta sandvicensis*). It's smaller than its Canadian goose cousin, and it doesn't migrate.

When humans introduced new species to the Hawaiian islands, the nēnē nearly went extinct because they were not adapted to predators. In the 1700s, there were estimated to be more than 25,000 nēnē living in the Hawaiian islands. In 1952, there were only 30! Fortunately, conservation efforts brought the nēnē back from the brink of extinction. Today, there are approximately 3,000 nēnē living in the islands of Hawaii.




The Founder Effect


In the original population, butterfly wings are approximately 75% spotted and 25% striped. Very rarely, there are solid color butterflies or "eyespots" pattern butterflies.




What happens when a small group of organisms starts or "founds" a large population? Consider three small groups of butterflies from the main population. Each founds a new population on an isolated island. Assuming the proportions of traits remain the same as the populations grow, what percentage of each type would you expect to see 15 years later?



0 % spotted
25 % striped
25 % solid
50 % eyespot




80 % spotted
20 % striped
0 % solid
0 % eyespot



0 % spotted
50 % striped
50 % solid
0 % eyespot

Population bottleneck


A population bottleneck occurs when a substantial portion of a the population is reduced. Famine and disease are common causes of a population bottleneck. When a small group is geographically separated from the main population (the founder effect) this is sometimes considered to be a type of bottleneck. In the case of the founder effect, other parts of the population exist that did not experience the loss of genetic diversity.



1 The original population of spheres is evenly mixed with solid, dotted, and striped individuals.

2 A bottleneck event drastically reduces the number of individuals. There are no solid spheres in the surviving population.

3 The population rebounds, but the distribution of traits is very different than before.



The Hawaiian Islands



More than 3,000 of the species living in the Hawaiian islands are endemic, meaning they are found nowhere else in the world. The island of Vancouver in British Columbia is slightly larger than the combined area of the Hawaiian islands. But it has less than 20 endemic species. Why are there more endemic species in Hawai'i?

Hawaii is very isolated. There are more than 2,000 miles of ocean surrounding Hawaii in every direction! Without the help of human transportation, it is very difficult for species to get to, or leave the islands. Evolution in an isolated environment is more likely to create endemic species. Vancouver island is only 25 miles from British Columbia.


Drifting Marbles

Read through the instructions for the simulation below. But before trying it out, go to the next page and make some predictions!

This activity simulates genetic drift by considering a population of 20 marbles. To keep things simple, the population is capped at 20, and just a single trait (color) is measured. For the instructions, we chose black and white marbles, but you could use any object. You will want 40 marbles (or peas, beans, pieces of candy, etc.) with half of them being marked or colored so that they can be easily distinguished. Each generation of marbles will pass on their colors to the next generation through random selection as follows.



○ → W




What do you predict will happen with the marble population? Will it stay half white marbles and half black marbles or will the proportions change over time?

The portions will likely change up and down over time. Eventually, it might become all black or all white. After that, it would stay unchanged.

Place a checkmark next to each outcome that will always be true for this simulation:

- There will always be 10 black and 10 white marbles.
- After enough generations, one of the colors will disappear entirely.
- There will always be some black marbles and some white marbles, but the number will vary.
- The number of each color will eventually become stable.

Below are 5 graphs. Which one do you think is the most likely to occur in this simulation of the number of black marbles in each generation?



The answer that eventually levels out with all black is most likely. With enough generations, eventually they will all be either black or all white.

Now go back and carry out the simulation for 20 generations. Was your prediction correct?
Yes, my prediction was essentially correct. After 17 generations, only the white marbles were left. It was a random process, but the limited size of the population caused one color to "die out" eventually.

Consider how the situation would change if we didn't cap the marble population at 20. Suppose each new generation added an additional marble. Do you think that one of the colors would be eliminated from the population?

Maybe. Sometimes in the simulation, one color will be eliminated, but growing population allows both versions to stay. (We ran the simulation many times for thousands of generations and saw both scenarios occur.)

In what type of population in the real world might we see the loss of an allele through genetic drift? Give an example, and explain why genetic drift might occur.

Animals might wash up on an isolated island. If the island is small, then the population will be limited, and we would see genetic diversity disappear over the generations just through random chance in the same way we saw the black marble die out in this simulation.

What is a species?

GENE POOL: All the genes and their alleles in an interbreeding population




The biological concept of a species is often defined as follows: living organisms that can produce fertile offspring belong to the same species. But in practice, it's more useful to think of a species as a **gene pool** where genetic information is exchanged.


For example, lions and tigers can interbreed and produce fertile offspring, but in the wild they live in separate populations. There are two separate gene pools, one for each species. As individuals of a species breed and produce offspring, the genetic information in the gene pool changes. Mutations may arise or disappear, and the frequency of certain traits will increase or decrease.

Life doesn't exactly fit into the categories people have made to help them understand the world. No matter how one defines a species, there are multiple examples where scientists have lively debates about whether a group of individuals are participating in the same gene pool or not.


Match the description of each species or group of animals with the corresponding diagram representing the gene flow within the gene pools.




Orcas are all currently classified as being part of the same species, but they live in ten distinct breeding populations. The different populations have different diets and hunting behaviors. Their social behaviors and communication techniques are different enough that they never interbreed, even when they live in the same geographic area.



Polar bears and grizzly bears are classified as two different species: *Ursus maritimus* and *Ursus arctos horribilis*. While they live in different habitats and have different behaviors and diets, they occasionally interbreed when they come in contact with each other, producing a hybrid called a "grolar bear," "grizzlar," or "nanulak." Genetic analysis shows that hybridization has also occurred in the past.




The carrion crow, *Corvus corone*, and the hooded crow, *Corvus cornix*, are classified as separate species but hybridize freely where their geographical ranges overlap.



Species in the Asteraceae, or daisy, family hybridize easily. Members of the *Helianthus* genus (sunflowers) are especially prone to hybridizing. If five wild sunflower species live in the same area, there will likely be hybrids between all of them!



If DNA from a prehistoric dragonfly was recovered and that dragonfly was cloned, would it be the same species as a modern dragonfly? Why or why not?

The appearance of dragonflies has remained similar over millions of years, but even so, it is unlikely that the genes will be compatible. More junk DNA will have been added to modern dragonflies and other mutations and changes would have occurred. Most likely, the modern genome would not be similar enough for ancient and modern dragonflies to breed.



Explain why a cheetah and a wolf belong to gene pools that will never naturally hybridize.

Cheetahs and wolves do not live in overlapping regions. However, even if they did, they wouldn't be able to breed. They have different numbers of chromosomes, and their most recent common ancestor lived millions of years ago.



Consider two populations of dogs, each with 1,000 individuals. The first contains pure-bred miniature poodles, and the second contains dogs from animal shelters, most of mixed or unknown breeding. Which population of dogs will have more diversity in their gene pool and why?




Which population would be more likely to survive if placed in a new location such as an island?

The animal shelter will have a wide variety of genes among the mixed breeds, while the poodles will share many genes, especially genes related to physical characteristics.


If they are transferred to an island to form a new population, the dogs from the animal shelter would have a better chance of surviving because their population would have more genetic diversity.

Taxonomy

Grouping living organisms according to shared attributes is called *taxonomy*. This field of study is useful in the study of biodiversity and conservation biology. There have been several different classification systems in the past. Systems change as new information is gathered.



Within each domain there are smaller categories: phylum, class, order, family, genus, and species. Notice how much more similar the animals become in each category in the diagram below. Can you circle the correct animal type from each category to show the classification of the domestic dog? The first two have been filled in as examples.




About Shared Traits




COMMON Nighthawk
Chordeiles minor

MOBILITY: Flying, walking, hopping
BEHAVIOR: Crepuscular/Nocturnal
DIET: Insectivore
ANATOMY: Bones are hollow and porous.
Lungs have air sacs.
REPRODUCTION: Lays eggs



BROWN RAT
Rattus norvegicus

MOBILITY: Walking, running, jumping
BEHAVIOR: Often nocturnal
DIET: Omnivore
ANATOMY: Bones are not hollow. Lungs do not have air sacs.
REPRODUCTION: Live birth



HOARY BAT
Aeorestes cinereus

MOBILITY: Flying, walking
BEHAVIOR: Crepuscular/Nocturnal
DIET: Insectivore
ANATOMY: Bones are not hollow. Lungs do not have air sacs.
REPRODUCTION: Live birth


Compare the nighthawk and rat. Which of these animals is more closely related to a bat?
Support your opinion by including similar and dissimilar traits.

The nighthawk and bat both fly and have similar diets. However, the brown rat and bat have more in common with their anatomy and reproduction. Bats are more closely related to rats than birds because bats and rats are both mammals. While the wings of a bird and bat both provide the ability to fly, their structures are quite different.

Homologous Traits

FILL IN THE BLANKS USING THESE WORDS:
similar homologous bone common

A homologous trait is one that is inherited from a common ancestor. The bones in the limbs of tetrapods are one example of homologous traits. Dogs, whales, bats, and humans all have front limbs that look and function differently from each other. But the bone structures of those limbs are remarkably similar. Each limb has a humerus, radius and ulna, carpals, metacarpals, and phalanges.



Convergent Traits

FILL IN THE BLANKS USING THESE WORDS:

environment diets convergent closely streamlined

Sharks and dolphins look a lot alike. They both have streamlined bodies with two fins on their sides and a dorsal fin on top of their body. They also have similar diets. But these similarities do not mean that sharks and dolphins are closely related. They each have a streamlined body shape, dorsal fin, and flippers because they live in the same environment and these traits help them to swim faster to catch prey. Their shared traits are an example of convergent evolution.

SHARK vs DOLPHIN

Draw lines to match each trait to the correct animal

Skeleton made of cartilage

Breathes air through blowhole

Skeleton made of bone


Uses gills to get oxygen from water

Do not nurse young with milk

Has hair

Nurses young with milk

Does not have hair




ARE THESE FACT OR FICTION? Write your verdict below each statement.

Bats and birds share a common ancestor.




True. All vertebrates share a common ancestor.

Today's elephants have evolved to be less likely to have tusks.



True. By poaching elephants with tusks, humans have artificially selected for a lack of tusks by making those genes more likely to be passed on.

Hippos are the ancestors of whales.



False. Hippos are the most closely related land mammals to whales, but not their ancestor. Both hippos and whales descended from a common ancestor that is now extinct.

The Fossil Record

FILL IN THE BLANKS USING THESE WORDS:

pressure survive fossils complete circumstances geological

Very few organisms leave behind **fossils**. Fossils can only form when an organism dies at the right time, in the right place, and under the right **circumstances**. Over a long period of time, intense heat and **pressure** are needed to encase the organism within a sedimentary rock that will eventually become the fossil. Usually, only the hard parts of the body like bones and teeth can **survive** this process to become a fossil. Even when a fossil is made, it may not survive **geological** changes over time as rocks break or melt. Fortunately, the fossil record is sufficiently **complete** for us to learn about how life evolved throughout history.


What can be inferred about the ages of the fossils in this picture? Why do you think one layer doesn't contain any fossils?

The oldest fossils (the trilobites on bottom) are on the lowest layer. We see the dinosaurs came later as they are in higher layers. The layer without fossils may indicate that earlier organisms died out.


What is the best explanation for a fossil being found in an older layer of rock but not any newer layers? Draw a picture of an example.

Likely, the organism died out so there was no chance for new fossils. It's also possible the new layer is not a type of rock that forms fossils.


Draw lines to match the name of each fossil type with its corresponding description:




BONY NOSTRILS MIGRATE TO TOP OF HEAD



Artiocetus - 47 million years



Dorudon - 36 million years



Tiktaalik - 375 million years



Pederpes - 348 million years

LEGS EVOLVE TO ASSIST MOVING ON LAND




Panderichthys - 380 million years


LEGS DISAPPEAR AND BODY BECOMES LONGER



Megachirella wachtleri - 240 million years



Tetrapodophis amplectus - 120 million years



Snake - living

Transition Fossils

FILL IN THE BLANKS USING THESE WORDS:

snakes transitional tetrapods whales discoveries document

The fossil record has gaps where organisms and characteristics have arisen with little fossil evidence to show how the new attributes came about. Fossils that fill in a perceived gap are called **transitional** fossils. Darwin didn't expect to be able to fill in these holes, but many surprising **discoveries** have filled in the "missing links" in the fossil record. **Snakes** evolved from lizards with legs. Four-legged creatures (**tetrapods**) evolved from fish and emerged from the water to live on the land. **Whales** evolved from land mammals as the nostrils gradually moved to the back of the skull, the front arms were replaced with flippers, and the hind legs diminished as the tail widened. We now have fossils to **document** each of these changes.

ARE THESE FACT OR FICTION? Write your verdict below each statement.

Soft tissue animals cannot form fossils.

IT SAYS HERE THAT JELLYFISH WILL NEVER BE FOSSILIZED.

BUT THAT'S NOT FAIR!

False. They can leave behind imprint fossils.

Bones can become fossils after being buried for 1,000 years.

HAPPY 1,000TH BIRTHDAY.

THANKS! I'M A FOSSIL NOW!

False. Fossils are at least 10,000 years old by definition.


Sea creature fossils have been found on Mt. Everest.

Fact


Phylogeny

FILL IN THE BLANKS USING THESE WORDS:


higher phylogeny common recent ancestor descendant speciation





A phylogenetic tree or a phylogeny is a diagram that depicts the lines of evolutionary descent of different species, organisms, or genes from a common ancestor. The bottom of a phylogenetic tree represents the past and the higher branches represent more recent times. As we travel from the bottom to the top of the tree we are tracing the history from ancestor to descendant. Each fork or branch represents the splitting of a lineage or speciation.



Highlight the unique or shared portions of the diagrams below. The first is completed as an example.




Each species on earth has ancestors that are unique to them and ancestors that they share with other species. Study the example of unique and common ancestors on the small phylogenetic tree for species A, B, and C. Then use the key to mark the unique and common ancestors in the larger phylogenetic tree.



△ Unique ancestor of C
□ Common ancestor of B and C
○ Common ancestor of A, B, and C

① Common ancestor of C - H
② Common ancestor of G and H
③ Common ancestor of F, G, and H
④ Common ancestor of C and D
⑤ Unique ancestor of E
⑥ Common ancestor of A and B



According to the phylogenetic chart above, is the hyena more closely related to cats or dogs?

Hyenas are more closely related to cats than dogs because hyenas and cats have a more recent common ancestor (marked ☆) than hyenas and dogs (marked △).


Are skunks more closely related to raccoons or mongooses? How can you tell?

Skunks are more closely related to raccoons than mongooses because skunks and mongooses have a more recent common ancestor (marked ○) than skunks and mongooses ?(marked △).


Hands-on Science Project

BUILD YOUR OWN PHYLOGENY

MATERIALS:



paper




crayons, colored pencils, or markers

Invent your own phylogeny by considering an organism (it can be real or invented) and the different environmental conditions it could face. Draw out different descendants and evolutionary branches that could result. Your conditions could be realistic and based on actual fossils or living organisms, or you could invent something fanciful like Math Dad did with the example phylogeny for shapes below.

Once you have your idea figured out, draw the complete phylogenetic tree and explain why each branch formed.

Math Dad's Example



2D shapes live in a flat plane and experience gravity. Circles are great for rolling and moving around, but they are not very good at stopping or standing still. In steep regions, rolling is a disadvantage and flat sides are a desirable trait.


In the left branch, we see descendants of the circle that lived in steeper regions where flat edges provided an advantage. Over time, the shapes evolved flat edges which led to the square. Triangles and rectangles had an advantage when minimal motion was desirable because they were unlikely to roll unintentionally, so they could live in extra steep regions. A hexagon and octagon were able to roll much more easily while still having straight edges for resting, so they were able to live in moderately-steep regions. But they could also compete in flatter regions with other more mobile or rounder shapes.

In the middle branch, the terrain was gently sloped with narrow pathways. Roundness was a useful trait but the shapes got longer over time because added length helped them avoid rolling too far. A thinner middle also made the shapes more flexible so they could navigate the narrow pathways.

In the right branch, we see a shape called a Reuleaux triangle that has a constant diameter like a circle. The Reuleaux triangles are able to climb in canyons of constant width just like circles. But they have an advantage over circles in that they have a smaller area, so they don't have to find as much food as a circle to survive.

Australia vs New Zealand

In ecosystems around the world, we find organisms adopting the same roles. There are producers (plants), animals that eat plants (herbivores), animals that eat insects (insectivores), and predators (carnivores). At the top of the food chain is a predator that has no natural predators (an apex predator). Consider these ecological niches in Africa, New Zealand, and Australia. Then answer the following questions:



Before humans arrived, bats and seals were the only mammals living in New Zealand. Why didn't New Zealand have other mammal species such as canines, felines, ungulates, or rodents?

New Zealand broke away from the Southern supercontinent Gondwana 80 million years ago.

Dinosaurs still roamed the Earth, and few mammals were around at that point. As an isolated

island, only flying or swimming mammals were able to get to New Zealand.

Why do you think birds were the dominant animal in New Zealand, while marsupial mammals dominated Australia?

Both Australia and New Zealand are examples of the founder effect. Placental mammals were not

living in either location. Marsupials evolved to occupy the main ecological niches (apex predators,


large grazers, small foragers, etc) in Australia while birds did the same in New Zealand.

Timeline of Life on Earth

Extinctions occur periodically as a normal part of Earth's natural history. Two billion years ago, cyanobacteria evolved photosynthesis and the ability to produce oxygen. Earth's atmosphere changed from one with virtually no oxygen to one with abundant oxygen, which was toxic to many forms of cellular life. This is called the Great Oxidation Event or the Oxygen Catastrophe. It is not known how many species went extinct because all life on Earth at this point was single celled, and the fossil record is sparse.

In normal conditions, we expect about 10% of species to be lost every million years while new species are arising.

A mass extinction occurs when 75% of the species die in a relatively short period of time. The fossil record shows evidence of five major mass extinction events in Earth's history.




Match each extinction event with the cause

Ordovician-Silurian (O-S) extinction - 450 MYA

85% of marine species

After this event, many trilobites, echinoderms, crinoids, bivalves, and brachiopods disappeared from the fossil record.

A massive asteroid collided with Earth.



Late Devonian (Late D) extinction - 375 MYA

75% of marine species

Placoderms (armored fish) went extinct as well as many species of coral, trilobites, and brachiopods.

The exact cause is unknown. It could have been a gamma-ray burst from a nearby supernova, glaciation, or volcanic eruptions.




Permian-Triassic (P-T) extinction - 250 MYA

96% marine life; 70% terrestrial life

Also known as the Great Dying. Trilobites became entirely extinct. This may be the only mass extinction to significantly affect insect diversity.

This extinction may have been caused by volcanic eruptions, plants colonizing land, an asteroid impact, climate change, or a combination of those factors.



Triassic-Jurassic (Tr-J) extinction - 200 MYA

70% of marine/terrestrial life

Large amphibians and crocodiles disappeared, making room for dinosaurs to diversify and fill other ecological niches.

Enormous volcanic eruptions in modern-day Siberia, the release of methane into the atmosphere, and climate change all contributed to this extinction.



Cretaceous-Paleogene (K-Pg) extinction - 65 MYA

75-80% of marine/terrestrial life


Large dinosaurs and pterosaurs became extinct. No tetrapods weighing more than 25 kg (55 lbs) survived.

Volcanic eruptions caused climate change. An asteroid impact could have also played a role.



Color the timeline to match each corresponding age or era.


4.6 Billion Years Ago



Precambrian


4.6 billion to 542 Million Years Ago (MYA)

The Precambrian era is the name for the first 4 billion years of Earth's existence. The fossil record is less complete because most life was in the ocean, and few organisms had hard bodies or shells.



Cambrian Explosion

542 MYA




Paleozoic Era

542 Million to 250 Million Years Ago

The Cambrian explosion started a period of incredible growth where the first land animals and vertebrates arose. The Paleozoic Era ended with the

Earth's largest mass extinction event.



Cenozoic Era

65 Million Years Ago to the present

NOW


Our current era is the Cenozoic. This era is characterized by alternating ice ages and warmer periods. The first humans appeared 2 million years ago.

Mesozoic Era

250 Million to 65 Million Years Ago

The Mesozoic Era saw the rise of the dinosaurs and the first mammals. Flowering plants developed for the first time during the Cretaceous Period. The Mesozoic Era ended when a giant meteorite crashed into the Gulf of Mexico, causing nearly 80% of all species to go extinct.

Evolution Crossword Puzzle



VERTICAL Words

- Characteristics of organisms that enhance survival and reproduction chances
- Darwin described evolution as "Descent with _____"
- A taxonomic group whose members can interbreed
- A change in the structure of a gene
- The immediate descendants of an organism

HORIZONTAL Words


- The preserved remains of an ancient organism
- The author of *On the Origin of Species*
- The process of changing over time
- Selective breeding for traits is _____ selection.
- Survival of the _____
- The process through which populations adapt and change is _____ selection
- Mapping the evolutionary development and diversification of species

Evolution Word Search

There are a lot of new words to learn when studying biology. Repetition is the best way to learn them, and word games can be part of that! Find each of the hidden words in the word-search. The words can run in any direction: horizontal or diagonal. The letters might go left to right or right to left!

ADAPTATION	REPRODUCTION	TRAIT
FITTEST	INHERITANCE	DARWIN
SPECIES	EVOLUTION	GENERATION
EXTINCTION	SELECTION	FOSSIL
OFFSPRING	POPULATION	MUTATION
CHARACTERISTIC	DRIFT	PHYLOGENY

Bonus words in blue -



Quiz Time!

ANSWER THE QUESTIONS TO SEE WHAT YOU LEARNED ABOUT EVOLUTION!

- 1 Any variation that can help an organism survive in its environment is called a(n)
 - A. adaptation
 - B. characteristic
 - C. competition
 - D. natural selection

- 2 Which of the following is NOT a component of the Theory of Evolution by Natural Selection?
 - A. variation among individuals
 - B. inheritance of acquired characteristics
 - C. competition for limited resources
 - D. survival and reproduction *Acquired traits are not passed on.*

- 3 The dog breeds we have today developed by
 - A. Sexual selection
 - B. Disruptive selection
 - C. Artificial selection (selective breeding)
 - D. Natural selection

- 4 What is the correct definition of evolution?
 - A. Changes in a population of organisms over a period of time
 - B. Changes in individual organisms during their lifetime
 - C. Organisms coming into existence at a defined moment in time and not changing
 - D. One organism transforming into another organism

- 5 Which of the following provide evidence for the evolution of species? Choose all that apply.
 - A. The fossil record
 - B. Biogeography (locations of organisms on the globe)
 - C. DNA
 - D. Comparative anatomy (between species)

- 6 In evolutionary biology, what does fitness mean?
 - A. The ratio of body fat to muscle weight
 - B. The ability of that individual to survive to reproductive age
 - C. The likelihood that that individual will have 2 or more offspring
 - D. The likelihood that an individual will contribute its genes to the next generation

- 7 Which of the following is the best example of natural selection?
 - A. A population of jackrabbits undergoes a seasonal coat color change from brown to white as soon as snow covers their habitat.
 - B. Field mice born during a year with excess wheat will gain muscle mass and thus have offspring that have an increased muscle mass.
 - C. Bass that are born with darker scales are better able to survive in the shadows among the river rocks.

- 8 Traits that perform a similar function but arise from different ancestral traits (such as bat and bird wings) are called
 - A. Anomalous
 - B. Homologous
 - C. Divergent
 - D. Convergent

- 9 The term species is defined as
 - A. A grouping of morphologically similar organisms
 - B. A grouping of organisms that do not interbreed
 - C. A group of populations that can interbreed to produce viable offspring
 - D. A group of populations that live in the same area

- 10 How many mass extinctions has the Earth experienced in the past?
 - A. 2
 - B. 5
 - C. 8
 - D. 9

- 11 According to the fossil record, how long ago did life appear on earth?
 - A. 3.5 million years
 - B. 35 million years ago
 - C. 350 million years ago
 - D. 3.5 billion years ago

- 12 Number each period below using the numbers 1-4 to put them in chronological order.
 - 3 Mesozoic
 - 1 Precambrian
 - 2 Paleozoic
 - 4 Cenozoic

Use the feline phylogeny to the right to answer the next four questions.


- 13 Of the options below, which species is most closely related to a lynx?
 - A. Ocelot
 - B. Cheetah
 - C. Tiger
 - D. Leopard

Look for the most recent common ancestor!

- 14 Of the options below, which species is most closely related to a Ocelot?
 - A. House cat
 - B. Leopard
 - C. Lion

- 15 An ancestor of both a cougar and a lynx must also be an ancestor of a cheetah.
 - A. True
 - B. False

- 16 Ocelots and Lions have no common ancestors.
 - A. True
 - B. False



Below are the mechanisms by which evolution takes place. Match the mechanism on the left with the corresponding example on the right.


Genetic Drift
A change in the frequency of a genotype due to chance

Parents with genes for light coloring could have an offspring with dark coloring.




Mutation
A change in the structure of a gene

A snake with light coloring might move to a region inhabited by dark colored snakes.




Migration
Movement from one region to another

Through random chance, the dark coloring gene might disappear from the population.



Natural Selection
The survival of a species that is better adapted to an environment

The environment might make it easier for predators to spot the light colored snakes.



APPENDIX



- 113 - The Codon Wheel
- 114-115 – The Math of Punnett Squares
- 116-117 - Tri-hybrid Cross
- 118-119 - Tetra-hybrid Cross
- 120-121 - Gummy Worm Karyotype Game
Don't print double-sided.
- 122- World Map for the Journey of the HMS Beagle Project
Don't print double-sided.
- 123 - Labels for the Journey of the HMS Beagle Project
Don't print double-sided.
- 124-end - Timeline of Earth's History
Don't print double-sided.

THE CODON WHEEL

A tool for deciphering the genetic code!

How to use: Start in the center and follow the letters outward!

For example, if the codon you are deciphering reads "AAT," select A in the inner circle, then A again in the second ring, and T in the third ring. This shows that AAT codes for the amino acid **asparagine**, which has a 3-letter abbreviation of **Asn** and a 1-letter abbreviation of **N**. Be aware that multiple codons can code for the same amino acid! Notice that the codon AAC also codes for asparagine.



The Math of Punnett Squares

The number of allele combinations quickly rises when the number of traits rise. For a genotype, the order of the alleles doesn't matter. For example, **Aa** is the same as **aA**. To keep things simple, we usually put the capital letter first. For example, when there is just one trait with alleles **A** and **a**, then there will be 3 possible genotypes (**AA**, **Aa**, and **aa**) but only 2 possible phenotypes: the dominant phenotype **A** (from **AA** or **Aa**) and the recessive phenotype **a** (from **aa**).

With 2 traits we must count the possible combinations for alleles **A** and **a** together with **B** and **b**.

Complete the table below by identifying how many distinct combinations arise for a given number of traits? You will probably want to write out the possibilities on a piece of scratch paper.

Number of traits	Number of distinct phenotypes	Number of distinct genotypes	Number of cells in a Punnett Square
1	2	3	4
2	4	9	16
3	8	27	64
4	16	81	256
5	32	243	1024
n	2^n	3^n	4^n

Now you have a formula that will let you find the number of distinct phenotypes or genotypes for any number of traits. Suppose we considered 25 specific traits that a person could have determined by their genotypes. For each of the possible phenotypes of the 25 traits, do you think there is someone in the world that exhibits that phenotype? Explain.

The total number of possible phenotypes is $2^{25} = 33,554,432$. There are 200 times as many people on Earth as that number, so it's very possible that each of the different phenotype combinations could exist for those 25 traits.

For each of the possible genotypes of the 25 traits, do you think there is someone in the world that has that genotype? Explain.

The total number of possible genotypes is $3^{25} = 847,288,609,443$. This is way bigger than the population of the Earth, so most of the genotype combinations will not occur.

Identical twins share the same alleles. Do you think it is possible for non-twins to have the same alleles? Explain.

Not unless you limit the scope of the question. Siblings are most likely to share many alleles, but there are so many different genotypes possible that two non-twins would only have the same alleles if you limit the inquiry to a small number of genes. For example, if you look at just 4 traits, there are only 81 possible genotypes, so many pairs of people would match genotypes on those 4 traits.

Calculating Probabilities

For a single trait, we can find the probability of passing a specific phenotype or genotype along by filling in a simple Punnett square.

For example, in the cross **Aa** × **Aa**, there are 4 equally likely outcomes: **AA**, **Aa**, **aA**, and **aa**. That means:

- $\frac{1}{4}$ of the time we get the homozygous dominant genotype **AA**.
- $\frac{1}{2}$ of the time we get the heterozygous genotype **Aa**.
- $\frac{1}{4}$ of the time we get the homozygous recessive genotype **aa**.

In the cross **Aa** × **Aa**, the probability (P) of each **genotype** is:

$$P(\text{genotype } \mathbf{AA}) = \frac{1}{4}$$

$$P(\text{genotype } \mathbf{Aa}) = \frac{1}{2}$$

$$P(\text{genotype } \mathbf{aa}) = \frac{1}{4}$$

In the cross **Aa** × **Aa**, the probability of each **phenotype** is:

$$P(\text{phenotype } \mathbf{A}) = \frac{3}{4}$$

$$P(\text{phenotype } \mathbf{a}) = \frac{1}{4}.$$

Punnett square for the cross **Aa** × **Aa**

	A	a
A	AA	Aa
a	aA	aa

To calculate the probabilities for multiple traits, we find the probabilities of the outcomes for each individual trait and then multiply the probabilities.

For example: Find the probability of getting the genotype **AaBB** in the cross **AABb** × **AaBB**.

1. First find the probability of the genotype **Aa** from the cross **Aa** × **Aa**.

$$P(\text{genotype } \mathbf{Aa}) = \frac{1}{2}$$

2. Find the probability of the genotype **BB** from the cross **Bb** × **BB**. $P(\text{genotype } \mathbf{BB}) = \frac{1}{2}$.

$$P(\text{genotype } \mathbf{BB}) = \frac{1}{2}$$

3. Multiply the probabilities.

$$P(\text{genotype } \mathbf{AaBB}) = P(\text{genotype } \mathbf{Aa}) \times P(\text{genotype } \mathbf{BB}) = \frac{1}{2} \times \frac{1}{2} = \frac{1}{4}$$

Challenge 1. Find the probability of getting the genotype **AaBb** in the cross **AABb** × **aaBB**.

The chance of getting **Aa** is $\frac{1}{2}$, and the chance of getting **Bb** is $\frac{1}{2}$, so the chance of getting **AaBb** from this cross is $\frac{1}{2} \times \frac{1}{2} = \frac{1}{4}$.

Challenge 2. Find the probability of getting the recessive phenotype **ab** in the cross **aabb** × **aaBb**.

The chance of getting phenotype **a** is $\frac{1}{2}$, and the chance of getting phenotype **b** is $\frac{1}{2}$, so the chance of getting both **a** and **b** is $\frac{1}{2} \times \frac{1}{2} = \frac{1}{4}$.

Challenge 3. Find the probability of getting the phenotype **AbC** in the cross **aabbCC** × **AaBbcc**.

The chance of getting phenotype **A** is $\frac{1}{2}$, phenotype **b** is $\frac{1}{2}$, and phenotype **C** is $\frac{1}{2}$, so the chance of getting phenotype **AbC** is $\frac{1}{2} \times \frac{1}{2} \times \frac{1}{2} = \frac{1}{8}$.

Challenge 4. Find the probability of getting the genotype **AaBbcc** in the cross **aaBbCc** × **AaBbcc**.

The chance of getting **Aa** is $\frac{1}{2}$, **Bb** is $\frac{1}{2}$, and **cc** is $\frac{1}{2}$, so the chance of getting phenotype **AaBbcc** is $\frac{1}{2} \times \frac{1}{2} \times \frac{1}{2} = \frac{1}{8}$.


Tri-hybrid Cross!

Make up a critter and supply phenotypes and genotypes for three different traits that can be determined by alleles.

Critter Name: _____

Trait 1: _____ vs _____

Phenotypes




Genotypes

_____ vs _____

Trait 2: _____ vs _____

Phenotypes




Genotypes

_____ vs _____

Trait 3: _____ vs _____

Phenotypes



Genotypes

_____ vs _____

What will the genotype and phenotype for your critter be if it is heterozygous for all three traits?

Genotype: _____

Phenotype: _____

Now cross two of your critters that are heterozygous for all three traits. Include both the genotype and phenotype in each cell of your Punnett square.

Tetrahybrid Cross!

Make up a critter and supply phenotypes and genotypes for four different traits that can be determined by alleles.

Critter Name: _____

What will the genotype and phenotype for your critter be if it is heterozygous for all four traits?

Genotype: _____

Phenotype:

A simple black rectangular outline, likely a placeholder for a drawing or a short response.

Now cross two of your critters that are heterozygous for all four traits.

<p>Trait 1: _____ vs _____</p> <p>Details: _____ _____</p>	<p>Phenotypes</p> <div style="display: flex; justify-content: space-around;"> <div style="text-align: center;">_____</div> <div style="text-align: center;">vs</div> <div style="text-align: center;">_____</div> </div> <p>Genotypes</p> <div style="display: flex; justify-content: space-around;"> <div style="text-align: center;">_____</div> <div style="text-align: center;">vs</div> <div style="text-align: center;">_____</div> </div>
<hr/>	
<p>Trait 2: _____ vs _____</p> <p>Details: _____ _____</p>	<p>Phenotypes</p> <div style="display: flex; justify-content: space-around;"> <div style="text-align: center;">_____</div> <div style="text-align: center;">vs</div> <div style="text-align: center;">_____</div> </div> <p>Genotypes</p> <div style="display: flex; justify-content: space-around;"> <div style="text-align: center;">_____</div> <div style="text-align: center;">vs</div> <div style="text-align: center;">_____</div> </div>
<hr/>	
<p>Trait 3: _____ vs _____</p> <p>Details: _____ _____</p>	<p>Phenotypes</p> <div style="display: flex; justify-content: space-around;"> <div style="text-align: center;">_____</div> <div style="text-align: center;">vs</div> <div style="text-align: center;">_____</div> </div> <p>Genotypes</p> <div style="display: flex; justify-content: space-around;"> <div style="text-align: center;">_____</div> <div style="text-align: center;">vs</div> <div style="text-align: center;">_____</div> </div>
<hr/>	
<p>Trait 4: _____ vs _____</p> <p>Details: _____ _____</p>	<p>Phenotypes</p> <div style="display: flex; justify-content: space-around;"> <div style="text-align: center;">_____</div> <div style="text-align: center;">vs</div> <div style="text-align: center;">_____</div> </div> <p>Genotypes</p> <div style="display: flex; justify-content: space-around;"> <div style="text-align: center;">_____</div> <div style="text-align: center;">vs</div> <div style="text-align: center;">_____</div> </div>

A blank 10x10 grid of squares, suitable for various applications such as graphing or tracking data.

Gummy Worm Karyotype Game

Instructions:

1. Prepare gummy worms according to instructions for the gummy worm karyotype activity.
2. Print off the Trait Key and Fruit Fly Phenotype Cards.
3. Cut out the 16 fruit fly phenotypes. Color them to match their description.
4. Take turns arranging karyotypes and matching the cards by playing one of the following versions of the Karyotype game!

*Note that although the Y and number 4 chromosome are colored in this template, there are no traits being tracked for these chromosomes in the karyotype game.

It's a race!

All 16 phenotype cards are placed face up. One person arranges a karyotype from either gummy worms or paper chromosomes. The karyotype is revealed to the players, who then race to find the correct phenotype card. The first person to choose the correct card earns a point! Play to 10 points.

Reverse it

Shuffle the phenotype cards and draw one at random. The player then uses gummy worms or paper chromosomes to arrange a karyotype that would produce that phenotype.

For a more advanced version, don't look at the trait key while building the phenotype! Just use it after the karyotype is made to check and see if it's correct. If it is, the player earns a point! Play to 10 points.

Double Trouble

Shuffle the phenotype cards and draw one at random. Each player should have a piece of paper and pencil and will draw a karyotype to match the phenotype card, making sure that their answer is not visible to the other player.



The karyotypes are revealed. If they match, neither player gets a point. If the karyotypes are different but both produce the same phenotype, each player receives 2 points. If either player creates a karyotype that does not match the card, both players lose 1 point. Play until 10 points.

Trait Key

- XX chromosomes = female
- XY chromosome pair = male
- Dominant trait in X = red eyes
- Recessive trait in X = white eyes
- Dominant trait in 2 = functional wings
- Recessive trait in 2 = miniature wings (flightless)
- Dominant trait in 3 = yellow body color
- Recessive trait in 3 = ebony or black body color


Chromosome Templates

Can be used in place of gummy worms for either gummy worm karyotypes or meiosis. Print this page twice, so there are 4 copies of each chromosome.










Fruit Fly Phenotype Cards

♂ Yellow, winged, red-eyed male	♂ Yellow, flightless, red-eyed male	♂ Yellow, winged, white-eyed male	♂ Yellow, flightless, white-eyed male
♀ Yellow, winged, + red-eyed female	♀ Yellow, flightless, + red-eyed female	♀ Yellow, winged, white-eyed	♀ Yellow, flightless, white-eyed
♂ Ebony, winged, red-eyed male	♂ Ebony, flightless, red-eyed male	♂ Ebony, winged, white-eyed male	♂ Ebony, flightless, white-eyed male
♀ Ebony, winged, red-eyed female	♀ Ebony, flightless, red-eyed female	♀ Ebony, winged, white-eyed	♀ Ebony, flightless, white-eyed




Journey of the H.M.S. Beagle



Hands-on Science Project

BUILD A TIMELINE OF LIFE ON EARTH

MATERIALS:













Print each of the 10 pages that follow. Print single-sided. Then choose a color for each of the circles below. Color the corresponding portion of the timeline to match. (Note that the Pangea circle overlaps two periods of time, the Carboniferous and Permian). Then cut along the dotted lines to create the narrow strips of paper for your timeline. Tape the strips together so that they form a continuous timeline of life on Earth!

*Note: The entire timeline will be approximately 80 ft (25 m) long. Each inch represents 5 million years of time! If you do not have room to lay out the entire timeline, you can put together 10 foot segments and line them up on a wall or floor.

As an optional extension, cut strips of colored paper to represent the four eons of Earth's history, and use these as a colorful backing to the timeline.

EON	TIME	NOTABLE EVENT(S)
Hadean	4.6 - 4 billion years ago	Earth formed and the surface cooled.
Archaean	4 - 2.5 billion years ago	First bacteria evolved. Atmosphere lacked oxygen.
Proterozoic	2.5 billion - 541 MYA	Oxygen entered the atmosphere.
Phanerozoic	541 MYA - present	This era includes everything from the Cambrian explosion onward.














For example:













Eon	Era	Period	Years (millions of years ago or MYA)
PHANEROZOIC	Cenozoic	Quaternary (Pleistocene/Holocene)	2.588 to 0
		Neogene (Miocene/Pliocene)	23.03 to 2.588
		Paleogene (Paleocene/Eocene/Oligocene)	66.0 to 23.03
	Mesozoic	Cretaceous	145.0 to 66.0
		Jurassic	201.3 to 145.0
		Triassic	252.17 to 201.3
	Paleozoic	Permian	298.9 to 252.17
		Carboniferous (Mississippian/Pennsylvanian)	358.9 to 298.9
		Devonian	419.2 to 358.9
		Silurian	443.4 to 419.2
		Ordovician	485.4 to 443.4
		Cambrian	541.0 to 485.4
PROTEROZOIC	Neoproterozoic	Ediacaran	635 to 541
		Cryogenian	720 to 635
		Tonian	1,000 to 720
	Mesoproterozoic	Stenian	1,200 to 1,000
		Ectasian	1,400 to 1,200
		Calymmian	1,600 to 1,400
	Paleoproterozoic	Statherian	1,800 to 1,600
		Orosirian	2,050 to 1,800
		Rhyacian	2,300 to 2,050
		Siderian	2,500 to 2,300
ARCHEAN	Neoarchean, Mesoarchean, Paleoarchean, Eoarchean	<i>Not divided into periods</i>	2,500 to 4,000
HADEAN	<i>Not divided into eras</i>		Formation of Earth to 4,000

*Some sources have slightly different start and end dates for different periods and eras, and that's okay! Remember that people invented periods, eras, and eons to help them study geologic history. Often the event that defines the end of one period and the beginning of another is an extinction event that spans hundreds of thousands or even millions of years.