

Name: Jenny Petrova

Problem Set 0

(P.1) *Prove Chebyshev's inequality: Consider a random variable X with finite expectation $\mathbb{E}X$ and variance $\mathbb{V}X$. Then*

$$\mathbb{P}(|X - \mathbb{E}X| \geq t) \leq \frac{\mathbb{V}X}{t^2},$$

for all $t > 0$.

First, recall Markov's inequality which provides, for a nonnegative random variable Y and positive real number $t > 0$,

$$\mathbb{P}(Y \geq t) \leq \frac{\mathbb{E}Y}{t}.$$

Since $|X - \mathbb{E}X| \geq 0$ and $t > 0$, the transformation $h(z) = z^2$ on $z \in [0, \infty)$ is a monotone transformation, implying

$$\mathbb{P}(|X - \mathbb{E}X| \geq t) = \mathbb{P}((X - \mathbb{E}X)^2 \geq t^2). \quad (1)$$

Applying Markov's inequality to the right-hand side of (1), we obtain

$$\mathbb{P}((X - \mathbb{E}X)^2 \geq t^2) \leq \frac{\mathbb{E}(X - \mathbb{E}X)^2}{t^2} = \frac{\mathbb{V}X}{t^2},$$

noting the definition of variance $\mathbb{V}X := \mathbb{E}(X - \mathbb{E}X)^2$, establishing the inequality

$$\mathbb{P}(|X - \mathbb{E}X| \geq t) \leq \frac{\mathbb{V}X}{t^2},$$

for all $t > 0$.

(P.2) *Prove the expected value of X when $X \sim \text{Poisson}(\lambda)$ for $\lambda \in (0, \infty)$.*

By definition,

$$\mathbb{E}X = \sum_{x=0}^{\infty} x \mathbb{P}(X = x) = \sum_{x=0}^{\infty} x \frac{\lambda^x \exp(-\lambda)}{x!}, \quad (2)$$

substituting in the form of the probability mass function for a Poisson random variable with rate parameter $\lambda \in (0, \infty)$. Next, we manipulate and re-arrange terms in (2):

$$\sum_{x=0}^{\infty} x \frac{\lambda^x \exp(-\lambda)}{x!} = \sum_{x=1}^{\infty} x \frac{\lambda^x \exp(-\lambda)}{x!} = \sum_{x=1}^{\infty} \frac{\lambda^x \exp(-\lambda)}{(x-1)!} = \lambda \sum_{x=1}^{\infty} \frac{\lambda^{x-1} \exp(-\lambda)}{(x-1)!}.$$

A change of index allows us to write

$$\sum_{x=1}^{\infty} \frac{\lambda^{x-1} \exp(-\lambda)}{(x-1)!} = \sum_{x=0}^{\infty} \frac{\lambda^x \exp(-\lambda)}{x!} = 1, \quad (3)$$

recognizing the right-hand side of (3) to be $\mathbb{P}(X \in \{0, 1, 2, \dots\})$, which has probability 1 by the axioms of probability because the support of a Poisson random variable is the set $\{0, 1, 2, \dots\}$. Altogether, we have shown

$$\mathbb{E}X = \lambda \sum_{x=1}^{\infty} \frac{\lambda^{x-1} \exp(-\lambda)}{(x-1)!} = \lambda,$$

establishing λ as the expected value of X .

(P.3) Let $\{X_n\}_{n \geq 1}$ be a sequence of Bernoulli random variables such that $\mathbb{P}(X_n = 1) = n^{-1}$. Prove that $X_n \xrightarrow{P} 0$, i.e., prove that X_n converges in probability to 0.

The sequence $\{X_n\}_{n \geq 1}$ converges in probability to a random variable X^* if

$$\lim_{n \rightarrow \infty} \mathbb{P}(|X_n - X^*| \geq \epsilon) = 0, \quad \text{for all } \epsilon > 0. \quad (4)$$

At least this is one definition. Let $\epsilon > 0$ and $n \in \{1, 2, \dots\}$ be arbitrary and consider the probability

$$\mathbb{P}(|X_n - 0| \geq \epsilon) = \mathbb{P}(|X_n| \geq \epsilon) = \mathbb{P}(X_n \geq \epsilon) = \mathbb{P}(X_n = 1),$$

which follows because $X_n \in \{0, 1\}$ so that $|X_n| = X_n$ and

$$\{x \in \{0, 1\} : x \geq \epsilon\} = \begin{cases} \{1\} & \epsilon \in (0, 1] \\ \{\} & \epsilon > 1 \end{cases}.$$

Thus, for any $\epsilon > 0$,

$$\lim_{n \rightarrow \infty} \mathbb{P}(|X_n - 0| \geq \epsilon) = \lim_{n \rightarrow \infty} \mathbb{P}(X_n = 1) = \lim_{n \rightarrow \infty} \frac{1}{n} = 0,$$

showing by (4) that $X_n \xrightarrow{P} 0$ as $n \rightarrow \infty$.