

# Galactoglucomannan fibres promote a beneficial porcine gut microbiome

Jenny Merkesvik<sup>a</sup>, Lars J. Lindstad<sup>a</sup>, Antton Alberdi<sup>b</sup>, Anders Miki Bojesen<sup>c</sup>, Marta Contreras-Serrano<sup>d</sup>, Jorge E. Langa<sup>be</sup>, Carlotta Pietroni<sup>b</sup>, Louise L. Poulsen<sup>c</sup>, Torgeir R. Hvidsten<sup>a</sup>, Phillip B. Pope<sup>af</sup>, and Bjørge Westereng<sup>a</sup>.

<sup>a</sup> Faculty of Chemistry, Biotechnology, and Food Science, Norwegian University of Life Sciences  
<sup>b</sup> Centre for Evolutionary Hologenomics, University of Copenhagen  
<sup>c</sup> Department of Veterinary and Animal Science, University of Copenhagen


<sup>d</sup> Centre for Volatile Interactions, University of Copenhagen  
<sup>e</sup> Faculty of Science and Technology, University of the Basque Country  
<sup>f</sup> Faculty of Biosciences, Norwegian University of Life Sciences

## 1 MOTIVATION

Most mammals and their microbiomes are codependent, forming a functional unit known as a **holobiont**. Exchanging metabolites, regulating gene expression, and combating pathogens are vital to the **health** and **performance** of the holobiont<sup>1-2</sup>. By understanding the **interactions** occurring within these complex systems, we can more effectively improve **animal** and **feed production**, favouring both animal welfare, production efficiency, and the growing needs of the increasing **human population**<sup>3-4</sup>.

**Mannan fibres** derived from spruce can be metabolised into **host-accessible compounds** by microbes with **carbohydrate-active enzymes**<sup>5-6</sup>. Microbes like these also ease piglets' transition from milk to solid feed<sup>7</sup>. Hence we ask: can we **jump-start** the young porcine microbiome by adding mannan fibres to their pre-weaning diet? **spoiler!**


## 2 THE TRIAL



The datasets were analysed both as **individual omic** layers (Sec. 3 and 4), and jointly through a **holo-omic** approach (Sec. 5).


## 3 MICROBIAL GRADIENTS

Principal component (PC) analyses<sup>8</sup> of **individual omic** layers in Fig. 1 show **gradients** that correspond with **mangan exposure** duration.




**Figure 1.** Principal component analyses of each omic layer. Points represent samples from piglets introduced to dietary mannan fibres at different developmental stages. Relative placements indicate similarities between samples' omic profiles.

**REFERENCES** 1 Rosenberg et al. The hologenome concept of evolution after 10 years. *Microbiome*. 2018;6(1):78. 2 Roughgarden et al. Holobionts as Units of Selection and a Model of Their Population Dynamics and Evolution. *Biol Theory*. 2018;13(1):44–65. 3 Simon et al. Host-microbiota interactions: from holobiont theory to analysis. *Microbiome*. 2019;7(1):5. 4 Alberdi et al. Disentangling host-microbiota complexity through hologenomics. *Nat Rev Genet*. 2022;23(5):281–97. 5 La Rosa et al. The human gut Firmicute Roseburia intestinalis is a primary degrader of dietary  $\beta$ -mannans. *Nat Commun*. 2019;10(1):905. 6 Michalak et al. Microbiota-directed fibre activates both targeted and secondary metabolic shifts in the distal gut. *Nat Commun*. 2020;11(1):5773. 7 Vasquez et al. Gut microbiome-produced metabolites in pigs: a review on their biological functions and the influence of probiotics. *J Anim Sci Technol*. 2022;64(4):671–95. 8 stats [base R, v.4.2.1]. 9 DESeq2 [R package, v.1.38.3], doi: 10.18129/B9.bioc.DESeq2. 10 MCFA [Python package v. 1.0.1], doi: 10.5281/zenodo.8128339.




read me later

follow  
3D'omics




## 4 SELECTIVE PROMOTION

Among the **differentially abundant**<sup>9</sup> populations between control and mannan-fed piglets are bacteria often associated with **host health** benefits: *Bifidobacterium longum* (log<sub>2</sub> fold change 8.3), *Lactobacillus johnsonii* (LFC 5.9), and *Butyribacter* sp. (LFC 5.7) (Fig. 2).



**Figure 2.** Three microbial populations with differential abundance between piglet groups. Significance by false discovery rate-adjusted p-values are indicated by asterisks: \* < 0.05, \*\* < 0.01, \*\*\* < 0.001.

Many of these populations' differentially expressed genes yield enzymes for **mangan degradation**, like glycoside **hydrolases** and **transferases**, and carbohydrate **esterases** (Fig. 3).




**Figure 3.** Differentially expressed features in the 'populations of interest containing genes for mangan degradation.

Thus it seems **yes, we can** jump-start the porcine microbiome by mangan supplementation!

## 5 HOLO-OMIC MODEL

A multiset correlation and factor analysis (MCFA)<sup>10</sup> model reconstructs the full dataset in three **feature spaces**: one **shared** by all omic layers; one **private** to each layer; and an omic-specific residual. Inferred shared components are similar to PCs, but comprises features from all omic layers. The model fit with mangan exposure (Fig. 4), hence it will be used to learn more about the **interactions** across the porcine **holo-omic boundary**.



**Figure 4.** Samples represented by MCFA components in two model spaces. The shared space contains features from all layers. Aspects private to each omic layer add to this shared space, together reconstructing the full holo-omic dataset.