

Galactoglucomannan fibres promote a beneficial porcine gut microbiome

Jenny Merkesvik ^a, Lars J. Lindstad ^a, Antton Alberdi ^b, Anders Miki Bojesen ^c, Marta Contreras-Serrano ^d, Jorge E. Langa ^{be}, Carlotta Pietroni ^b, Louise L. Poulsen ^c, Torgeir R. Hvidsten ^a, Phillip B. Pope ^{af}, and Bjørge Westereng ^a

^a Faculty of Chemistry, Biotechnology, and Food Science, Norwegian University of Life Sciences

^b Centre for Evolutionary Hologenomics, University of Copenhagen

^c Department of Veterinary and Animal Science, University of Copenhagen

^d Centre for Volatile Interactions, University of Copenhagen

^e Faculty of Science and Technology, University of the Basque Country

^f Faculty of Biosciences, Norwegian University of Life Sciences

1 MOTIVATION

Most mammals and their microbiomes are codependent, forming a functional unit known as a **holobiont**. Exchanging metabolites, regulating gene expression, and combating pathogens are vital to the **health** and **performance** of the holobiont ¹⁻². By understanding the interactions occurring within these systems, we can more effectively improve **animal** and **feed production**, favouring both animal welfare, production efficiency, and the needs of the World's **growing human population** ³⁻⁴.

Mannan fibres made from spruce can be metabolised into **host-accessible compounds** by microbes with carbohydrate-active **enzymes** ⁵⁻⁶. These microbes also ease piglets' transition from milk to **solid feed** ⁷. Hence we ask; can we **jump-start the young porcine microbiome by dietary mannan fibre supplementation?** spoiler!

2 THE TRIAL

We used **three groups** of 10-day old piglets and gave fibres to two groups; one starting **before** and another **after weaning**.



6 x control
no mannan




6 x late
two weeks with mannan




6 x early
four weeks with mannan

After one month, we sampled their caeca and generated **Four omic data layers**: metagenomics, meta-transcriptomics, metabolomics, and host transcriptomics. The resulting data were analysed both as **individual omic** layers and jointly through a **holo-omic** approach. Specifically, utilised methods were R-implemented tools for principal component ⁸ and differential abundance ⁹ analyses, and a Python package for holo-omic modelling through multiset correlation and factor analysis (MCFA) ¹⁰.

3 SELECTIVE PROMOTION




Individual principal component analyses show **gradients** that correspond to **duration of mannan exposure**. Among the populations with different abundance in control and mannan piglets are ***Bifidobacterium longum*** (log₂ fold change 8.3), ***Lactobacillus johnsonii*** (LFC 5.9), and ***Butyribacter sp.*** (LFC 5.7). Many of their differentially expressed genes yield enzymes for mannan degradation, like glycoside **hydrolases** and **transferases**, and carbohydrate **esterases**.




Thus **yes, we can** jump-start their microbiome!

4 HOLO-OMIC MODEL

The model reconstructs the full dataset using **three feature spaces**: one **shared** for all omic layers; one **private** to each layer; and lastly, an omic-specific **residual**. The inferred features fit well to mannan exposure; hence this model will be used to learn about the **interactions across this holo-omic boundary**.



REFERENCES ¹ Rosenberg *et al.* The hologenome concept of evolution after 10 years. *Microbiome*. 2018;6(1):78. ² Roughgarden *et al.* Holobionts as Units of Selection and a Model of Their Population Dynamics and Evolution. *Biol Theory*. 2018;13(1):44–65. ³ Simon *et al.* Host-microbiota interactions: from holobiont theory to analysis. *Microbiome*. 2019;7(1):5. ⁴ Alberdi *et al.* Disentangling host-microbiota complexity through hologenomics. *Nat Rev Genet*. 2022;23(5):281–97. ⁵ La Rosa *et al.* The human gut Firmicute *Roseburia intestinalis* is a primary degrader of dietary β-mannans. *Nat Commun*. 2019;10(1):905. ⁶ Michalak *et al.* Microbiota-directed fibre activates both targeted and secondary metabolic shifts in the distal gut. *Nat Commun*. 2020;11(1):5773. ⁷ Vasquez *et al.* Gut microbiome-produced metabolites in pigs: a review on their biological functions and the influence of probiotics. *J Anim Sci Technol*. 2022;64(4):671–95. ⁸ stats [base R, v.4.2.1]. ⁹ DESeq2 [R package, v. 1.38.3]. doi: 10.18129/B9.bioc.DESeq2. ¹⁰ MCFA [Python package v. 1.0.1] doi: 10.5281/zenodo.8128339.



read
me later

follow
3D'omics

