

Galactoglucomannan fibres promote a beneficial porcine gut microbiome

Jenny Merkesvik ^a, Lars J. Lindstad ^a, Antton Alberdi ^b, Anders Miki Bojesen ^c, Marta Contreras-Serrano ^d, Jorge E. Langa ^{be}, Carlotta Pietroni ^b, Louise L. Poulsen ^c, Torgeir R. Hvidsten ^a, Phillip B. Pope ^{af}, and Bjørge Westereng ^a

^a Faculty of Chemistry, Biotechnology, and Food Science, Norwegian University of Life Sciences
^b Centre for Evolutionary Hologenomics, University of Copenhagen
^c Department of Veterinary and Animal Science, University of Copenhagen

^d Centre for Volatile Interactions, University of Copenhagen
^e Faculty of Science and Technology, University of the Basque Country
^f Faculty of Biosciences, Norwegian University of Life Sciences

1 MOTIVATION

Most mammals and their microbiomes are codependent, forming a functional unit known as a **holobiont**. Exchanging metabolites, regulating gene expression, and combating pathogens are vital to the **health** and **performance** of the holobiont. By understanding the interactions occurring within this system, we can more effectively improve **animal** and **feed production**, favouring both animal welfare, production efficiency, and the world's **growing human population**.

Mannan fibres made from spruce can be broken down into **host-accessible compounds** by microbes with carbohydrate-active **enzymes**. These microbes can also ease piglets' **transition** from milk to **solid feed**. Can we **jump-start the porcine microbiome by mannan fibre supplementation?**


2 THE TRIAL

We used **three groups** of 10-day old piglets and gave fibres in addition to the basal feed to two groups; one starting **before** and one **after weaning**.




After one month, we sampled their caeca and generated **four omic data layers**: metagenomics, meta-transcriptomics, metabolomics, and host transcriptomics. The resulting data were analysed both as **individual omics** layers and jointly through a **holo-omic** approach.

3 SELECTIVE PROMOTION



Individual principal component analyses show **gradients** corresponding to **mannan exposure** duration. Among the microbial populations of different abundance in control and mannan-fed piglets, we find ***Bifidobacterium longum*** (LFC 8.3), ***Lactobacillus johnsonii*** (LFC 5.9), and ***Butyrivacter*** (LFC 5.7).




Their metaT activity and metaB connection, hostT impacts...

4 HOLO-OMICS TITLE

Holo-omic model, shared and private dimensions through MCCA...

REFERENCES 1 ... 2 ... 3 ... 4 ... 5 ... 6 ...



read
me later

follow
3D'omics

