

Galactoglucomannan fibres promote a beneficial porcine gut microbiome

Jenny Merkesvik ^a, Lars J. Lindstad ^a, Antton Alberdi ^b, Anders Miki Bojesen ^c, Marta Contreras-Serrano ^d, Jorge E. Langa ^{b,e}, Carlotta Pietroni ^b, Louise L. Poulsen ^c, Torgeir R. Hvidsten ^a, Phillip B. Pope ^{a,f}, and Bjørge Westereng ^a.

^a Faculty of Chemistry, Biotechnology, and Food Science, Norwegian University of Life Sciences
^b Centre for Evolutionary Hologenomics, University of Copenhagen
^c Department of Veterinary and Animal Science, University of Copenhagen

^d Centre for Volatile Interactions, University of Copenhagen
^e Faculty of Science and Technology, University of the Basque Country
^f Faculty of Biosciences, Norwegian University of Life Sciences


1 MOTIVATION

Most mammals and their microbiomes are codependent, forming a functional unit known as a **holobiont**. Exchanging metabolites, regulating gene expression, and combating pathogens are vital to the **health** and **performance** of the holobiont ^{1,2}. By understanding the **interactions** occurring within these complex systems, we can more effectively improve **animal** and **feed production**, favouring both animal welfare, production efficiency, and the growing needs of the World's increasing **human population** ^{3,4}.


Mannan fibres made from spruce can be metabolised into **host-accessible compounds** by microbes with carbohydrate-active **enzymes** ^{5,6}. These microbes also ease piglets' transition from milk to **solid feed** ⁷. Hence we ask: can we **jump-start the young porcine microbiome by adding mannan to their pre-weaning diet?** *spoiler!*

2 THE TRIAL

We used **three groups** of 10-day old piglets and gave fibres to two groups: one starting **before** and one **after weaning**.



6 x control
no mannan




3 x late
two weeks with mannan




3 x early
two weeks with mannan

After one month, we sampled their caeca and generated **four omic data layers**: metagenomics, metatranscriptomics, metabolomics, and host transcriptomics. The datasets were analysed both as **individual omic** layers, and jointly through a **holo-omic** approach. Utilised methods were R-implemented tools for principal component ⁸ and differential abundance ⁹ analyses, and a Python-based package for holo-omic modelling through multiset correlation and factor analysis (MCFA) ¹⁰.

3 SELECTIVE PROMOTION




Individual principal component analyses show **gradients** that correspond with duration of **mannan exposure**. Among the differentially abundant populations in control and mannan-fed piglets are ***Bifidobacterium longum*** (log₂ fold change 8.3), ***Lactobacillus johnsonii*** (LFC 5.9), and ***Butyrivacter* sp.** (LFC 5.7). Many of their differentially expressed genes yield enzymes for mannan **degradation**, like glycoside **hydrolases** and **transferases**, and carbohydrate **esterases**.




Thus **yes, we can** jump-start their microbiome!

4 HOLO-OMIC MODEL

The fitted MCFA model reconstructs the full dataset as **matrices** across **three feature spaces**: one **shared** by all omic layers; one **private** to each dataset; and an omic-specific **residual**. The inferred features fit well with mannan exposure, hence the model will be used to learn more about the **interactions** across the porcine **holo-omic boundary**.



REFERENCES **1** Rosenberg *et al.* The hologenome concept of evolution after 10 years. *Microbiome*. 2018;6(1):78. **2** Roughgarden *et al.* Holobionts as Units of Selection and a Model of Their Population Dynamics and Evolution. *Biol Theory*. 2018;13(1):44–65. **3** Simon *et al.* Host-microbiota interactions: from holobiont theory to analysis. *Microbiome*. 2019;7(1):5. **4** Alberdi *et al.* Disentangling host-microbiota complexity through hologenomics. *Nat Rev Genet*. 2022;23(5):281–97. **5** La Rosa *et al.* The human gut Firmicute *Roseburia intestinalis* is a primary degrader of dietary β -mannans. *Nat Commun*. 2019;10(1):905. **6** Michalak *et al.* Microbiota-directed fibre activates both targeted and secondary metabolic shifts in the distal gut. *Nat Commun*. 2020;11(1):5773. **7** Vasquez *et al.* Gut microbiome-produced metabolites in pigs: a review on their biological functions and the influence of probiotics. *J Anim Sci Technol*. 2022;64(4):671–95. **8** stats [base R, v.4.2.1]. **9** DESeq2 [R package, v. 1.38.3], doi: 10.18129/B9.bioc.DESeq2. **10** MCFA [Python package v. 1.0.1], doi: 10.5281/zenodo.8128339.



read me
later

follow
3D'omics

