

# Galactoglucomannan fibres promote a beneficial porcine gut microbiome

Jenny Merkesvik <sup>a</sup>, Lars J. Lindstad <sup>a</sup>, Antton Alberdi <sup>b</sup>, Anders Miki Bojesen <sup>c</sup>, Marta Contreras-Serrano <sup>d</sup>, Jorge E. Langa <sup>be</sup>, Carlotta Pietroni <sup>b</sup>, Louise L. Poulsen <sup>c</sup>, Torgeir R. Hvidsten <sup>a</sup>, Phillip B. Pope <sup>af</sup>, and Bjørge Westereng <sup>a</sup>

<sup>a</sup> Faculty of Chemistry, Biotechnology, and Food Science, Norwegian University of Life Sciences  
<sup>b</sup> Centre for Evolutionary Hologenomics, University of Copenhagen  
<sup>c</sup> Department of Veterinary and Animal Science, University of Copenhagen

<sup>d</sup> Centre for Volatile Interactions, University of Copenhagen  
<sup>e</sup> Faculty of Science and Technology, University of the Basque Country  
<sup>f</sup> Faculty of Biosciences, Norwegian University of Life Sciences


## 1 MOTIVATION

Most mammals and their microbiomes are codependent, forming a functional unit known as a **holobiont**. Exchanging metabolites, regulating gene expression, and combating pathogens are vital to the **health** and **performance** of the holobiont. By understanding the interactions occurring within this system, we can more effectively improve **animal** and **feed production**, favouring both animal welfare, production efficiency, and the **growing human population**.

**Mannan fibres** made from spruce can be broken down into **host-accessible compounds** by microbes with carbohydrate-active **enzymes**. These microbes can also ease piglets' **transition** from milk to **solid feed**. Can we **jump-start the porcine microbiome by mannan fibre supplementation?**



## SELECTIVE PROMOTION




Individual principal component analyses show **gradients** corresponding to **mannan exposure** duration. Among the populations ...

## 2 THE TRIAL


We used **three groups** of 10-day old piglets and gave **fibres** in addition to the basal feed to two groups; one starting **before** and one **after weaning**.



6 x control  
no mannan




3 x late  
two weeks with mannan



3 x early  
four weeks with mannan

After **one month**, we sampled their caeca and generated **four omic data layers**: metagenomics, metatranscriptomics, metabolomics, and host transcriptomics. The resulting data were analysed both as **individual omics** layers and jointly through a **holo-omic** approach.

## REFERENCES 1 ... 2 ... 3 ... 4 ... 5 ... 6 ...



read  
me later

follow  
3D'omics

