

Sparsity-inducing Bayesian Causal Forest with Instrumental Variable

Jens Klenke^{1, $\bar{\wedge}$} , Lennard Maßmann^{1,2,*}

¹Chair of Econometrics, University of Duisburg-Essen

²Ruhr Graduate School in Economics

*E-mail: lennard.massmann@uni-due.de

$\bar{\wedge}$ E-mail: jens.klenke@vwl.uni-due.de

February 21, 2024

Abstract

Your text:Suspendisse vitae elit. Aliquam arcu neque, ornare in, ullamcorper quis, commodo eu, libero. Fusce sagittis erat at erat tristique mollis. Maecenas sapien libero, molestie et, lobortis in, sodales eget, dui. Morbi ultrices rutrum lorem. Nam elementum ullamcorper leo. Morbi dui. Aliquam sagittis. Nunc placerat. Pellentesque tristique sodales est. Maecenas imperdiet lacinia velit. Cras non urna. Morbi eros pede, suscipit ac, varius vel, egestas non, eros. Praesent malesuada, diam id pretium elementum, eros sem dictum tortor, vel consectetuer odio sem sed wisi.

Contents

1	Introduction	1
1.1	Literature	1
1.2	Contribution	1
2	Potential Outcomes and IV	3
2.1	Potential Outcomes and ITT	3
2.2	IV assumptions	3
2.3	Estimation	4
3	Sparsity-inducing Bayesian Causal Forest with Instrumental Variable	5
3.1	Sparse Bayesian Causal Forest	5
3.2	Sparse BCF-IV	5
4	Simulation study	6
5	Empirical application	7
6	Discussion	8

1 Introduction

Hahn et al. (2020)

1.1 Literature

- Using machine learning to infer heterogeneous effects in observational studies often focuses on CATE estimation under regular assignment mechanisms
- Focus in this work: Methods to discover heterogeneous effects in the presence of imperfect compliance
- Methods
 - tree-based
 - ensemble-of-trees
 - deep-learning-based methods
- BCF-IV: Discovers and estimates HTE in an interpretable way
 - BCF: BART-based semi-parametric Bayesian regression model, able to estimate HTE in regular assignment mechanisms, even with strong confounding
 - Use BCF to estimate $\hat{\tau}_C(x)$ and $\widehat{ITT}_Y(x)$ such that the conditional Complier Average Causal Effect $\hat{\tau}^{cace}(x) = \frac{\widehat{ITT}_Y(x)}{\hat{\tau}_C(x)}$

1.2 Contribution

- - BART benefits
 - good performance in high-noise settings
 - shrinkage to/emphasize on low-order interactions
 - established software implementations (‘BayesTree’, ‘bartMachine’, ‘dbarts’)
- BART shortcomings

- non-smooth predictions as BART prior produces stepwise-continuous functions
 - BART prior is overconfident in regions with weak common support
- Research proposal: Rewrite the BCF-IV model with SoftBART instead of BART prior to account for sparsity

2 Potential Outcomes and IV

2.1 Potential Outcomes and ITT

- Y_i : outcome variable; W_i : treatment variable; Z_i : instrumental variable
- \mathbb{X} : $N \times P$ matrix of control variables
- G_i : sub-populations of units
 - $G_i = C : W_i(Z_i = 0) = 0, W_i(Z_i = 1) = 1$
 - $G_i = D : W_i(Z_i = 0) = 1, W_i(Z_i = 1) = 0$
 - $G_i = AT : W_i(Z_i = 0) = 1, W_i(Z_i = 1) = 1$
 - $G_i = NT : W_i(Z_i = 0) = 0, W_i(Z_i = 1) = 0$

Given the Stable Unit Treatment Value Assumption (SUTVA), one can postulate the existence of potential outcomes $Y_i(W_i)$ such that $Y_i^{obs} = Y_i(1)W_i + Y_i(0)(1 - W_i)$.

One can directly get from the data the effect of the assignment to treatment:

$$\begin{aligned} ITT_Y &= \mathbb{E}[Y_i|Z_i = 1] - \mathbb{E}[Y_i|Z_i = 0] \\ &= \pi_C ITT_{Y,C} + \pi_D ITT_{Y,D} + \pi_{AT} ITT_{Y,AT} + \pi_{NT} ITT_{Y,NT}. \end{aligned}$$

2.2 IV assumptions

Assumptions to infer Complier Average Causal Effect (CACE),

$$\tau^{cace} = ITT_{Y,C} = \frac{\mathbb{E}[Y_i|Z_i = 1] - \mathbb{E}[Y_i|Z_i = 0]}{\mathbb{E}[W_i|Z_i = 1] - \mathbb{E}[W_i|Z_i = 0]} = \frac{ITT_Y}{\pi_C},$$

and its conditional version (cCACE),

$$\tau^{cace}(x) = \frac{\mathbb{E}[Y_i|Z_i = 1, \mathbb{X}_i = x] - \mathbb{E}[Y_i|Z_i = 0, \mathbb{X}_i = x]}{\mathbb{E}[W_i|Z_i = 1, \mathbb{X}_i = x] - \mathbb{E}[W_i|Z_i = 0, \mathbb{X}_i = x]} = \frac{ITT_Y(x)}{\pi_C(x)}.$$

1. exclusion restriction: $Y_i(0) = Y_i(1)$, for $G_i \in \{AT, NT\}$.
2. monotonicity: $W_i(1) \geq W_i(0) \rightarrow \pi_D = 0$.
3. existence of compliers: $P(W_i(0) < W_i(1)) > 0 \rightarrow \pi_C \neq 0$.
4. unconfoundedness of the instrument: $Z_i \perp (Y_i(0, 0), Y_i(0, 1), Y_i(1, 0), Y_i(1, 1), W_i(0), W_i(1))$.

2.3 Estimation

The conditional CACE can be estimated in a generic sub-sample (i.e., for each $\mathbf{X}_i \in \mathbb{X}_j$, where \mathbb{X}_j is a generic node of the discovered tree, like a non-terminal node or a leaf) as:

3 Sparsity-inducing Bayesian Causal Forest with Instrumental Variable

3.1 Sparse Bayesian Causal Forest

3.2 Sparse BCF-IV

4 Simulation study

5 Empirical application

6 Discussion

Bibliography

Hahn, P. R., Murray, J. S. & Carvalho, C. M. (2020), ‘Bayesian Regression Tree Models for Causal Inference: Regularization, Confounding, and Heterogeneous Effects (with Discussion)’, *Bayesian Analysis* **15**(3), 965–1056. Publisher: International Society for Bayesian Analysis.

URL: <https://projecteuclid.org/journals/bayesian-analysis/volume-15/issue-3/Bayesian-Regression-Tree-Models-for-Causal-Inference-Regularization-Confounding/10.1214/19-BA1195.full>