Imatissma posticata -	[1.4, 5.1]	-	[-0.23, 1.5]		0.59
Imatissma sackeni -	[-2.4, 6.3]	-	[-0.97, 1.6]		0.51
Laetodon laetus -	[-4.5, 1.2]	-	[-0.65, 1.5]		0.5
Lejops bilinearis -	[-0.53, 5]	-	[–0.26, 1.8]		0.55
Lejops chrysostomus -	[-3.7, 3.2]	-	[–0.51, 1.5]		0.55
Lejops lineatus -	[-1.8, 0.63]	-	[-1.1, 0.38]		0.8
Melanostoma mellina -	[0.88, 5.9]	-	[-0.91, 1.4]		0.61
Meliscaeva cinctella -	[-1.8, 3.1]	-	[–1, 1.3]		0.54
Merodon equestris -	[2.1, 3.9]	-	[-1.4, -0.22]		0.94
Meromacrus acutus -	[0.62, 2.2]	-	[-0.11, 1]		0.87
Meromacrus gloriosus -	[-2.9, 6]	-	[-0.93, 1.6]		0.3
Milesia virginiensis -	[1.4, 3.4]	-	0.082, 1.4		0.72
Monoceromyia floridensis -	[-3.5, 5.9]	-	[–0.95, 1.6]		0.29
Myáthropa florea -	[2.1, 7.6]	-	[–2.3, 1.1]		0.73
Myólepta strigilata -	[-4.3, 1.4]	-	[-0.98, 1.1]		0.5
Nausigaster unimaculata -	[-3.4, 5.9]	-	[–0.87, 1.6]		0.44
Ňuntianus cubanus -	[-4.1, 5.7]	-	[-0.92, 1.6]		0.38
Ocyptamus antiphates -	[-2.8, 6]	-	[-0.97, 1.6]		0.42
Ócyptamus costatus -	[-1.2, 0.89]	-	[-0.23, 0.98]		0.76
Ocyptamus dimidiatus -	[-3.3, 2.1]	-	[-0.76, 1.5]		0.49
Ocyptamus fuscipennis -	[1, 2.4]	-	[-0.018, 0.89]		0.96
Ornidia obesa -	[-5.1, -3.3]	-	[-0.88, 0.27]		0.85
Orphnabaccha jactator -	[-4.8, 5.6]	-	[-0.91, 1.6]		0.35
· Orthonevra flukei -	[–2.3, 6.1]	-	[–0.87, 1.5]		0.5
Orthonevra nitida -	[1.5, 4]	-	[0.25, 1.9]		0.7
Palpada agrorum -	[-2.2, -0.18]	-	[-0.51, 0.74]		0.68
Palpada albifrons -	[-4.4, -0.84]	-	[–1.1, 0.82]		0.51
Palpada alhambra -	[-5.8, -3.3]	-	[-0.65, 0.4]		0.84
Palpada furcata -	[-0.45, 1.1]	-	[-0.29, 0.77]		0.8
Palpada mexicana -	[-5.3, -3.3]	-	[-0.38, 0.7]		0.9
				nativity	'
	(5)		(5)	\mathcal{C}_{i} ;	KR 55
	cil [®]		cile .	Viji.	
				100°	× ·
	(5)		(S)		
	pecies species 1		aditat species		
	ciles		Oito		
. (&	Źĸ	1	(o-		
M ₂		Mar.			
		11.			