Spilomyia fusca -	[-0.55, 1.4]
Spilomyia interrupta -	[-0.98, 1.6]
Spilomyia liturata -	[-0.91, 1.5]
Spilomyia lóngicornis -	[-0.55, 0.57]
. Spilomyia sayi-	[-0.86, 1.3]
Syritta flaviventris -	[-0.33, 1.2]
Syritta pipiens -	[-0.68, 0.43]
Syrphus knabi-	[-0.35, 1.4]
Syrphus opinator -	[-0.67, 1.9]
Syrphus rectus -	[-0.8, 0.75]
Syrphus ribesii -	[-0.55, 1.7]
Syrphus torvus -	[-0.56, 1.3]
Temnostoma daochum -	[–0.81, 1.2]
Teuchocnemis lituratus -	[-0.16, 1.7]
Toxomerus boscii -	[-0.29, 0.96]
Toxomerus corbis -	[-0.74, 1.6]
_ Toxomerus floralis -	[–0.84, 1.6]
Toxomerus geminatus -	[0.23, 2.1]
_Toxomerus jussiaeae -	[-0.64, 1.7]
Toxomerus marginatus -	[-0.35, 0.97]
Toxomerus occidentalis -	[-0.83, 1.5]
Toxomerus parvulus -	[-0.89, 1.6]
_ Toxomerus politus -	[-0.24, 0.91]
Toxomerus verticalis -	[-0.89, 1.6]
Tropidia albistylum -	[-0.28, 1.4]
Tropidia quadrata -	[-0.63, 1.6]
Victoriana attenuata -	[-0.89, 1.6]
Victoriana parvicornis -	[-0.83, 1.6]
Volucella bombylans -	[-0.59, 1.8]
	:(0)
	o'C'
المراجع	
Y	
Sal.	
	Joseph. Species

ativity