Helophilus fasciatus -	[-0.95, 1.3]	
Helophilus latifrons -	[-1.3, 0.96]	
Hypocritanus fascipennis -	[-0.81, 0.25]	
Hypocritanus lemur -	[–1, 1.5]	
Imatissma albipilis -	[-0.55, 1.8]	
Imatissma bautias -	[-0.72, 0.98]	
Imatissma posticata -	[-0.34, 1.5]	
Imatissma sackeni -	[-0.82, 1.7]	
Laetodon laetus -	[-0.82, 1.6]	
Lejops bilinearis -	[-0.28, 1.8]	
Lejops lineatus -	[-1.2, 0.43]	
Melanostoma mellina -	[-0.098, 2.1]	
Meliscaeva cinctella -	[–0.81, 1.6]	
Merodon equestris -	[-0.73, 0.39]	
Meromacrus acutus -	[–0.21, 1.1]	
Milesia virginiensis -	[0.12, 1.5]	
Myathropa florea -	[–1.3, 1.5]	
Nausigaster unimaculata -	[–0.4, 1.8]	
Nuntianus cubanus -	[-0.69, 1.7]	
Ocyptamus antiphates -	[–0.91, 1.7]	
Ocyptamus costatus -	[-0.39, 1]	
Ocyptamus dimidiatus -	[–0.74, 1.6]	
Ocyptamus fuscipennis -	[-0.26, 0.73]	
Ornidia obesa -	[-0.76, 0.6]	
Orphnabaccha jactator -	[–0.8, 1.7]	
Orthonevra flukei -	[–0.83, 1.6]	
Orthonevra nitida -	[0.31, 1.9]	
Palpada agrorum -	[-0.67, 0.8]	
Palpada albifrons -	[-0.87, 1]	
	1	
	· 65)	
	- Cile	
	200	4
	19 8	
	·	
	[-0.87, 1]	
	, O,	
- 1	No.	
	X	