

Danaus plexippus -	[-1.8, -1.1]	-	[-1.0, 1.9]
Nymphalis antiopa -	[-2.0, -1.2]	-	[-2.7, 3.9]
Lon melane -	[-2.1, -1.4]	-	[-4.1, 3.0]
Pieris rapae -	[-2.4, -1.5]	-	[-2.1, 2.9]
Hylephila phyleus -	[-3.0, -1.6]	-	[-4.2, 2.9]
Dione vanillae -	[-2.9, -1.7]	-	[-4.3, 1.4]
Vanessa cardui -	[-2.8, -1.9]	-	[-6.0, -1.7]
Junonia grisea -	[-3.4, -1.6]	-	[-4.2, 2.7]
Vanessa atalanta -	[-3.3, -2.0]	-	[-2.1, 3.1]
Anthocharis sara -	[-3.4, -1.9]	-	[-2.7, 3.8]
Autographa californica -	[-3.7, -1.7]	-	[-5.5, 0.7]
Euphydryas chalcedona -	[-3.6, -1.9]	-	[-4.0, 3.6]
Phoebis sennae -	[-3.8, -1.8]	-	[-4.4, 0.4]
Pontia protodice -	[-3.6, -2.0]	-	[-3.4, 3.3]
Apodemia virgulti -	[-3.7, -2.0]	-	[-3.6, 3.8]
Hyles lineata -	[-3.5, -2.2]	-	[0.2, 4.9]
Spodoptera ornithogalli -	[-4.7, -2.4]	-	[-4.4, 2.2]
Mythimna unipuncta -	[-4.7, -2.5]	-	[-3.9, 3.5]
Burnsius albezens -	[-4.6, -2.6]	-	[-2.8, 3.9]
Feltia subterranea -	[-4.7, -2.6]	-	[-3.9, 2.9]
Limenitis lorquini -	[-4.6, -2.6]	-	[-3.4, 4.5]
Noctua pronuba -	[-4.8, -2.6]	-	[-4.7, 2.1]
Hemieuxoa rudens -	[-4.8, -2.6]	-	[-4.8, 1.8]
Emmelina monodactyla -	[-4.8, -2.6]	-	[-4.6, 2.1]
Coenonympha californica -	[-4.9, -2.6]	-	[-4.6, 2.6]
Neoterpes edwardsata -	[-4.9, -2.6]	-	[-4.3, 3.2]
Euxoa auxiliaris -	[-4.8, -2.7]	-	[-4.2, 3.4]
Trichoplusia ni -	[-4.7, -2.6]	-	[-5.4, 0.3]
Uresiphita reversalis -	[-5.0, -2.7]	-	[-4.6, 3.1]
Pyrausta volupialis -	[-5.0, -2.7]	-	[-4.6, 1.6]
Coptodisca powellella -	[-5.0, -2.7]	-	[-4.0, 3.4]
Pyrausta unifascialis -	[-5.4, -2.7]	-	[-5.2, 2.2]

detection
rate

persistence
rate