

INF5620 Exam, Problem 5

Jens Kristoffer Reitan Markussen

December 15, 2013

Nonlinear diffusion

$$u_t = \nabla \cdot (\alpha(u) \nabla u), \quad x \in \Omega, t > 0 \quad (1)$$

$$u(x, 0) = I(x), \quad x \in \Omega \quad (2)$$

$$u(x, t) = g(x), \quad x \in \partial\Omega_D, t > 0 \quad (3)$$

$$-\alpha(u) \frac{\partial}{\partial n} u(x, t) = h(u - T_s), \quad x \in \partial\Omega_R \quad (4)$$

$$\alpha(u) = (1 + \alpha_0 u^4) \quad (5)$$

a) Compact difference notation CN,

$$[D_t u = \nabla \cdot (\overline{\alpha(u)}^t \nabla u)]^{n+\frac{1}{2}} \quad (6)$$

Written out

$$\frac{u^{n+1} - u^n}{\Delta t} = \nabla \cdot (\overline{\alpha(u^{n+\frac{1}{2}})}^t \nabla u^{n+\frac{1}{2}}) \quad (7)$$

$$\frac{u^{n+1} - u^n}{\Delta t} = \frac{1}{4} (\nabla \cdot (\alpha(u^n) + \alpha(u^{n+1}) \nabla (u^n + u^{n+1}))) \quad (8)$$

Picard

$$F(u^{n+1}) = u^{n+1} - u^n - \Delta t \frac{1}{4} (\nabla \cdot (\alpha(u^n) + \alpha(u^{n+1}) \nabla (u^n + u^{n+1}))) \quad (9)$$

Replace u^{n+1} with $u^{n+1,k}$, where k is the picard iteration index, in all the non-linear terms

$$F(u^{n+1}) = u^{n+1} - u^n - \Delta t \frac{1}{4} (\nabla \cdot (\alpha(u^n) + \alpha(u^{n+1,k}) \nabla (u^n + u^{n+1,k}))) \quad (10)$$

b) Perform a Backward-Euler time discretization and derive the variational form for the resulting spatial problems. Picard iteration method to linearize

..

c) Newton's method to the nonlinear variational form $F = 0$ in b).

$$J_{i,j}\delta u = -F_i \tag{11}$$

$$J_{i,j} = \frac{\partial F_i}{\partial u_j}$$