Hypergamy Is Solved

Feasibility, Deficit, and Equilibrium in Status-Ordered Matching Markets

October 23, 2026

Abstract

Hypergamy, the tendency to form unions with a partner of higher status, is usually argued about as a behavioral claim. We instead model the marriage market as a coupling problem between Borel probability measures on Polish type spaces and show that, absent taste heterogeneity, every aggregate question about hypergamy reduces to a question about two marginal distributions. We prove: (i) universal hypergamy is feasible if and only if one status distribution first-order stochastically dominates the other (Strassen); (ii) otherwise the minimal unmatched mass is the *Hypergamy Deficit* $\mathsf{D} = \sup_t (G_{\mathfrak{W}} - G_{\mathfrak{M}})^+$ (a continuum Hall marriage theorem); (iii) no behavioral strategy changes any of this (*Cope Invariance*), and status gains are zero-sum in rank (*Conservation of Rank*); (iv) under status-ranked preferences the unique Gale–Shapley stable matching is positive assortative, and under Becker–Shapley–Shubik transferable utility it is the comonotone optimal transport plan, with the Choo–Siow entropic relaxation tending to random matching as taste shocks grow. For equal height variances we derive a closed-form deficit for required height gaps, $\mathsf{D}(\delta) = (2\Phi(\frac{\delta - \Delta m}{2\sigma}) - 1)^+$, which vanishes up to $\Delta m \approx 14$ cm and, at the pooled US standard deviation $\sigma \approx 7.35$ cm, reaches 0.74 at one foot. With several traits, checklist requirements shrink the eligible pool multiplicatively when traits are independent, trade-offs reduce to a scalar index, and a required rank gap δ leaves exactly δ of one side unmatched. We model partner history as a comparison effect, which applies to both sexes.

1 The Marriage Market as a Measure Space

Definition 1.1 (Market)**.** A *marriage market* is a tuple $(\mathcal{X}, \mathcal{Y}, \mu, \nu, s)$ where $\mathcal{X}$ and $\mathcal{Y}$ are Polish spaces of agent types for the two sides $\mathfrak{M}$ and $\mathfrak{W}$, $\mu \in \mathcal{P}(\mathcal{X})$ and $\nu \in \mathcal{P}(\mathcal{Y})$ are Borel probability measures describing the populations, and $s : \mathcal{X} \sqcup \mathcal{Y} \to \mathbb{R}$ is a measurable *status functional* (height, income, education, follower count, or any scalar index thereof). We write $\mu_s = s_{\#}\mu$ and $\nu_s = s_{\#}\nu$ for the pushforward status distributions, with CDFs $F_{\mathfrak{M}}, F_{\mathfrak{W}}$ and survival functions $G_{\mathfrak{M}}(t) = \mu(s \geq t)$, $G_{\mathfrak{W}}(t) = \nu(s \geq t)$.

Definition 1.2 (Matching)**.** A *matching* is a coupling $\pi \in \Pi(\mu, \nu)$, i.e. a Borel probability measure on $\mathcal{X} \times \mathcal{Y}$ with marginals μ and ν. A *partial matching* is a sub-coupling: a measure π with $(p_1)_{\#}\pi \leq \mu$ and $(p_2)_{\#}\pi \leq \nu$. Its *mass* $|\pi| = \pi(\mathcal{X} \times \mathcal{Y})$ is the amount of each side that is matched.

Definition 1.3 (Hypergamy)**.** A partial matching π is *hypergamous for* $\mathfrak{W}$ if $\pi(\{s(x) \geq s(y)\}) = |\pi|$, and *strictly hypergamous* if the inequality is strict π-a.e. The *hypergamy index* is $\mathsf{H}(\pi) = \pi(\{s(x) > s(y)\})$ and the *hypogamy index* is $\mathsf{H}^{\downarrow}(\pi) = \pi(\{s(x) < s(y)\})$.

Remark 1.4. Every definition is symmetric under exchanging $(\mathcal{X}, \mu)$ with $(\mathcal{Y}, \nu)$. Conventional usage applies hypergamy to $\mathfrak{W}$, and we follow that convention for notation only. No result depends on which side is labeled which, and the reader is free to swap the letters if it makes them feel better.

2 Who Can Marry Up? Feasibility and the Deficit

Theorem 2.1 (Hypergamy Feasibility)**.** *A hypergamous matching $\pi \in \Pi(\mu, \nu)$ exists if and only if μ_s first-order stochastically dominates ν_s:*

$$G_{\mathfrak{M}}(t) \geq G_{\mathfrak{W}}(t) \qquad \text{for all } t \in \mathbb{R}. \tag{1}$$

Proof. ($\Rightarrow$) If π is hypergamous then $\{s(y) \geq t\} \subseteq \{s(x) \geq t\}$ up to a π-null set, so $G_{\mathfrak{W}}(t) = \pi(s(y) \geq t) \leq \pi(s(x) \geq t) = G_{\mathfrak{M}}(t)$. ($\Leftarrow$) With generalized inverses $F^{-1}(u) = \inf\{t : F(t) \geq u\}$, condition (1) gives $F^{-1}_{\mathfrak{M}}(u) \geq F^{-1}_{\mathfrak{W}}(u)$ for all u, so for $U \sim \mathrm{Unif}(0,1)$ the pair $(F^{-1}_{\mathfrak{M}}(U), F^{-1}_{\mathfrak{W}}(U))$ couples (μ_s, ν_s) on $\{a \geq b\}$. Lift to $\mathcal{X} \times \mathcal{Y}$ by disintegrating μ, ν along the fibers of s (possible since the spaces are Polish). This is the one-dimensional case of Strassen's theorem [10]. □

When (1) fails, someone has to marry down or stay single. We compute exactly how many. Allow general masses $\mu(\mathcal{X}) = m$, $\nu(\mathcal{Y}) = w$ and let $\mathcal{C} = \{(x, y) : s(x) \geq s(y)\}$.

Theorem 2.2 (Hypergamy Deficit Formula)**.** *The maximal mass of a partial matching supported on $\mathcal{C}$ is*

$$\max_{\pi(\mathcal{C}^c)=0} |\pi| = w - \mathsf{D}, \qquad \mathsf{D} := \sup_{t \in \mathbb{R}} \big(\nu(s \geq t) - \mu(s \geq t)\big)^+. \tag{2}$$

D *is the minimal mass of $\mathfrak{W}$ left unmatched if every union is hypergamous.*

Proof. Upper bound. Agents in $\mathfrak{W}$ with $s(y) \geq t$ are compatible only with $\{x : s(x) \geq t\}$, so at least $\nu(s \geq t) - \mu(s \geq t)$ of them are unmatched; take the supremum. *Achievability.* Leave the top D of $\mathfrak{W}$ by status unmatched, splitting an atom if necessary, and use only the top $w - \mathsf{D} \leq m$ of $\mathfrak{M}$. These women and men have survival functions $(G_{\mathfrak{W}} - \mathsf{D})^+$ and $\min(G_{\mathfrak{M}}, w - \mathsf{D})$, and the second dominates the first because $G_{\mathfrak{W}} - G_{\mathfrak{M}} \leq \mathsf{D}$ and $G_{\mathfrak{W}} \leq w$, so Theorem 2.1, rescaled to mass $w - \mathsf{D}$, matches them hypergamously. For finitely supported measures this is the deficiency form of Hall's theorem: compatibility is *nested*, so Hall's condition need only be checked on upper sets. □

Corollary 2.3. $\mathsf{D} = 0$ *iff* (1) *holds (when $m = w$), and $\mathsf{D} \geq (w - m)^+$. A skewed sex ratio is a lower bound on the deficit; everything above it is caused by the shape of the distributions.*

Example 2.4. Let $\mu_s = \mathrm{Unif}[0,1]$ and $\nu_s = \mathrm{Beta}(2,1)$, so $G_{\mathfrak{M}}(t) = 1 - t$ and $G_{\mathfrak{W}}(t) = 1 - t^2$. Then $\mathsf{D} = \sup_t (t - t^2) = \frac{1}{4}$ at $t^* = \frac{1}{2}$. With a perfectly balanced sex ratio, one quarter of $\mathfrak{W}$ still cannot be matched hypergamously, and the bottleneck sits exactly at the median of $\mathfrak{M}$.

3 Application: The Six-Foot Rule

Height is the cleanest test case: scalar, observable, approximately Gaussian within each sex, and the subject of strongly held opinions. Take $s =$ height in cm with $\mu_s = N(m_{\mathfrak{M}}, \sigma^2_{\mathfrak{M}})$ and $\nu_s = N(m_{\mathfrak{W}}, \sigma^2_{\mathfrak{W}})$. We calibrate to US adult means from NHANES 2015–2018 [5], $m_{\mathfrak{M}} \approx 175.3$ and $m_{\mathfrak{W}} \approx 161.3$, write $\Delta m \approx 14.0$, and take $\sigma_{\mathfrak{M}} = 7.6$ and $\sigma_{\mathfrak{W}} = 7.1$. The report gives percentiles rather than standard deviations, and its percentiles imply 7.4–7.5 and 6.9–7.1.

Proposition 3.1 (The -37.5 cm Theorem)**.** *For Gaussian marginals with $\sigma_{\mathfrak{M}} \neq \sigma_{\mathfrak{W}}$, the survival functions cross exactly once, at*

$$t_\times = \frac{\sigma_{\mathfrak{M}} m_{\mathfrak{W}} - \sigma_{\mathfrak{W}} m_{\mathfrak{M}}}{\sigma_{\mathfrak{M}} - \sigma_{\mathfrak{W}}} \approx -37.5\,\mathrm{cm}. \tag{3}$$

Hence (1) *fails only on $(-\infty, t_\times)$ and $\mathsf{D}(0) < 10^{-150}$.*

Proof. $G_{\mathfrak{M}}(t) = G_{\mathfrak{W}}(t)$ iff $(t - m_{\mathfrak{M}})/\sigma_{\mathfrak{M}} = (t - m_{\mathfrak{W}})/\sigma_{\mathfrak{W}}$, a linear equation with unique root $t_\times$. Above it, $\mathfrak{M}$ has the smaller standardized score, so $G_{\mathfrak{M}} > G_{\mathfrak{W}}$. Below it the gap is at most Φ of a standardized value near -28. □

Remark 3.2. "He must be taller than me" is therefore feasible for every human of positive height. The only obstruction occurs below -37.5 cm, a region the author was unable to sample. Other readings of the report's percentiles move the crossing anywhere from -6 to -108 cm, which the author did not sample either.

Now require a gap. A union satisfies the *δ-rule* if $s(x) \geq s(y) + \delta$. Shifting ν_s by δ reduces this to Theorem 2.2, so $\mathsf{D}(\delta) = \sup_t (G_{\mathfrak{W}}(t) - G_{\mathfrak{M}}(t + \delta))^+$.

Theorem 3.3 (Closed-Form Height Deficit)**.** *If $\sigma_{\mathfrak{M}} = \sigma_{\mathfrak{W}} = \sigma$, then*

$$\mathsf{D}(\delta) = \left(2\,\Phi\!\left(\frac{\delta - \Delta m}{2\sigma} \right) - 1 \right)^+ . \tag{4}$$

D *vanishes iff $\delta \leq \Delta m$, and is concave increasing beyond it with initial slope $(\sigma\sqrt{2\pi})^{-1}$, or* 5.4% *per* cm *at the pooled $\sigma = 7.35$* cm*.*

Proof. With $z = (t - m_{\mathfrak{W}})/\sigma$ and $c = (\delta - \Delta m)/\sigma$, the gap is $\Phi(z + c) - \Phi(z)$, the Gaussian mass of an interval of length c. For $c > 0$ it is maximized by centering the interval at the mode, $z = -c/2$, giving $2\Phi(c/2) - 1$. Its δ-derivative $\varphi(c/2)/\sigma$ is decreasing in c. □




Figure 1: Minimal fraction of $\mathfrak{W}$ left unmatched as a function of the required height gap, US calibration with unequal variances. Essentially zero until about 13 cm, then roughly 5% per centimeter.

Theorem 3.4 (The Six-Foot Theorem)**.** *Suppose a fraction α of $\mathfrak{W}$ requires $s(x) \geq \tau$, and everyone else imposes any constraint whatsoever. Then at least $(\alpha - G_{\mathfrak{M}}(\tau))^+$ of $\mathfrak{W}$ is unmatched in every feasible matching. For $\tau = 182.9$* cm *(six feet), $G_{\mathfrak{M}}(\tau) \approx 0.16$.*

Proof. The constrained set A has neighborhood inside $\{s \geq \tau\}$, of mass $G_{\mathfrak{M}}(\tau)$. Apply the Hall bound from Theorem 2.2. □

Remark 3.5. Theorem 3.4 holds regardless of personality, effort, or app. It is also generous: it assumes every six-footer is available and interested.

4 The Blackpill Theorems

Lemma 4.1 (Conservation of Status)**.** *If $s \in L^1(\mu) \cap L^1(\nu)$, then for every $\pi \in \Pi(\mu,\nu)$, $\int (s(x) - s(y))\, d\pi = \int t\, d\mu_s - \int t\, d\nu_s$. The average status gap across all couples does not depend on who marries whom.*

Proof. Integrate against π and use its marginals. □

Definition 4.2 (Strategy)**.** A *strategy* is any map $\mathfrak{S} : \Pi(\mu,\nu) \to \Pi(\mu,\nu)$: every intervention that changes who matches with whom without changing the population. This includes openers, dating apps, "just be confident," and every other object the literature calls *cope*.

Theorem 4.3 (Cope Invariance)**.** *If $\Psi : \Pi(\mu,\nu) \to \mathbb{R}$ depends on π only through its marginals, then $\Psi \circ \mathfrak{S} = \Psi$ for every strategy $\mathfrak{S}$. In particular the mean status gap, the feasibility of universal hypergamy, the deficit D, and the equilibrium index H^* of (8) are invariant under every strategy.*

Proof. Strategies preserve marginals, and each listed quantity is a functional of (μ_s, ν_s) alone. □

Corollary 4.4 (It's Over)**.** *If $\mu_s = \nu_s$, no strictly hypergamous matching exists and no strategy produces one: by Lemma 4.1 applied to the bounded status* $\arctan s$*, the integrand* $\arctan s(x) - \arctan s(y)$ *would be positive π-a.e. with zero integral.*

The only escape is to change the marginals, replacing s by $s + h$ with $h \geq 0$ on a subset of agents. We call this *maxxing*.

Theorem 4.5 (Conservation of Rank)**.** *Let $F_{\mathfrak{M}}, F_{\mathfrak{W}}$ be continuous and let the market clear at the comonotone coupling. If $U = F_{\mathfrak{M}}(s(x))$ is an agent's own quantile and $V = F_{\mathfrak{W}}(s(y))$ their partner's, then $V = U$ a.s. and $U \sim \mathrm{Unif}(0,1)$. Hence for any perturbation $s \mapsto s + h$ that keeps $F_{\mathfrak{M}}$ and $F_{\mathfrak{W}}$ continuous, maxxing included,*

$$\mathbb{E}[V] = \tfrac{1}{2}, \qquad \mathrm{Var}(V) = \tfrac{1}{12} \tag{5}$$

before and after. Maxxing by a subpopulation raises its partners' quantiles by exactly the amount it lowers everyone else's.

Proof. Under the comonotone coupling $s(y) = F_{\mathfrak{W}}^{-1}(F_{\mathfrak{M}}(s(x)))$, so $V = U$. The probability integral transform gives $U \sim \mathrm{Unif}(0,1)$ for any continuous $F_{\mathfrak{M}}$, perturbed or not. □

> **Remark (The Blackpill, Stated Precisely).** Theorems 4.3 and 4.5 are the rigorous content of "it's over": aggregate outcomes are pinned by the marginals, strategies act within a fiber of fixed marginals, and status gains are transfers of rank, not creation of it. But both are statements about the *population*. For an individual at quantile u the comonotone map is strictly increasing, so raising your own status raises your partner's quantile one-for-one, and for $\theta > 0$ (Section 7) outcomes are random rather than determined.
>
> **It is over for the population. It is not over for you.**

5 Everyone Wants Up, Almost Everyone Goes Down

Take finite sides $|\mathfrak{M}| = |\mathfrak{W}| = n$ with distinct statuses and *status-ranked* preferences: every agent strictly prefers a higher-status partner, and any partner to none. A matching is *stable* if no pair would both rather be with each other than with their assigned partners.

Theorem 5.1 (Uniqueness of the Assortative Equilibrium)**.** *Under status-ranked preferences the unique stable matching pairs the k-th highest of $\mathfrak{M}$ with the k-th highest of $\mathfrak{W}$ for every k, and Gale–Shapley deferred acceptance [6] returns it whichever side proposes.*

Proof. Induct on n. If the two top agents are not matched to each other, they form a blocking pair. Remove them; the reduced market has the same structure. □

Corollary 5.2 (Everyone Wants Up, Almost Everyone Goes Down)**.** *In Example 2.4 the assortative matching is the quantile map $T = F_{\mathfrak{M}}^{-1} \circ F_{\mathfrak{W}}$, $T(b) = b^2 < b$ on $(0,1)$, and no pair blocks it: a man at quantile u prefers a woman at quantile v only if $v > u$, and she prefers him only if $u > v$. Hence $\mathsf{H} = 0$ and $\mathsf{H}^{\downarrow} = 1$: every member of $\mathfrak{W}$ prefers to marry up, the market is stable, and almost every member of $\mathfrak{W}$ marries down, by an average of $\int_0^1 (b - b^2)\, 2b\, db = \frac{1}{6}$.*

6 Transferable Utility and Optimal Transport

Following Becker [1] and Shapley–Shubik [8], a couple with statuses a and b generates surplus $f(a,b)$, which partners may split. The equilibrium solves the Monge–Kantorovich problem

$$\sup_{\pi \in \Pi(\mu,\nu)} \int f(s(x), s(y))\, d\pi = \inf_{u(x)+v(y) \geq f(s(x),s(y))} \left\{ \int u\, d\mu + \int v\, d\nu \right\}, \tag{6}$$

where equality is Kantorovich duality and (u, v) are the equilibrium utilities [11].

Theorem 6.1 (Becker via Cyclical Monotonicity)**.** *If f is strictly supermodular ($\partial_{ab} f > 0$), every optimal plan has status projection equal to the comonotone coupling $\gamma^* = (F_{\mathfrak{M}}^{-1}, F_{\mathfrak{W}}^{-1})_{\#}\,\mathrm{Unif}(0,1)$.*

Proof. Optimal plans are cyclically monotone. If $a_1 < a_2$ and $b_1 > b_2$ were both in the support, then

$$f(a_2,b_1) + f(a_1,b_2) - f(a_1,b_1) - f(a_2,b_2) = \int_{a_1}^{a_2} \int_{b_2}^{b_1} \partial_{ab} f\, d\beta\, d\alpha > 0, \tag{7}$$

so swapping partners strictly improves total surplus, a contradiction. □

So bargaining or not, the realized hypergamy index is the same:

$$\mathsf{H}^* := \mathsf{H}(\gamma^*) = \mathrm{Leb}\{u \in (0,1) : F_{\mathfrak{M}}^{-1}(u) > F_{\mathfrak{W}}^{-1}(u)\}. \tag{8}$$

7 Taste Heterogeneity: The Entropic Relaxation

Real agents are not perfectly status-ranked. Following Choo and Siow [3] and Galichon and Salanié [7], add i.i.d. Gumbel taste shocks of scale $\theta > 0$. With finitely many types, write f_{ij} for the surplus of a type-i man with a type-j woman. Equilibrium couples π_{ij} and singles π_{i0}, π_{0j} satisfy

$$\pi_{ij} = \sqrt{\pi_{i0}\pi_{0j}}\, e^{f_{ij}/(2\theta)}, \tag{9}$$

the first-order conditions of the entropy-regularized problem

$$\max_{\pi \geq 0} \Big\{ \sum_{i,j} f_{ij}\pi_{ij} - \theta\, \mathrm{Ent}(\pi) \Big\}, \qquad \mathrm{Ent}(\pi) = 2\sum_{i,j} \pi_{ij} \log \pi_{ij} + \sum_i \pi_{i0} \log \pi_{i0} + \sum_j \pi_{0j} \log \pi_{0j}, \tag{10}$$

subject to the population constraints. Writing $a_i = \sqrt{\pi_{i0}}$, $b_j = \sqrt{\pi_{0j}}$, $K_{ij} = e^{f_{ij}/(2\theta)}$, these become $a_i^2 + a_i (Kb)_i = \mu_i$ and $b_j^2 + b_j (K^\top a)_j = \nu_j$, solved alternately in closed form by a generalized iterative proportional fitting procedure.

Theorem 7.1 (Vanishing and Exploding Taste)**.** *Let the market be balanced and f strictly supermodular. If every $f_{ij} > 0$, then as $\theta \downarrow 0$ the equilibrium converges to an optimal plan of (6) and $\mathsf{H} \to \mathsf{H}^*$. In the variant without singles this holds for any sign of f_{ij}, and as $\theta \to \infty$ the equilibrium converges to $\mu \otimes \nu$ and $\mathsf{H} \to \mathbb{P}(S_{\mathfrak{M}} > S_{\mathfrak{W}})$ for independent $S_{\mathfrak{M}} \sim \mu_s$, $S_{\mathfrak{W}} \sim \nu_s$.*

Sketch. Γ-convergence of the regularized objectives on a compact feasible set. If every $f_{ij} > 0$, a man and a woman left single can be matched for a gain, so limit plans have no singles, and Theorem 6.1 fixes their status projection, on which H depends; for $\theta \to \infty$ the entropy term dominates and its unique minimizer over $\Pi(\mu, \nu)$ is the product measure. □

In Example 2.4 the endpoints are $\mathsf{H}^* = 0$ and $\mathbb{P}(S_{\mathfrak{M}} > S_{\mathfrak{W}}) = \frac{1}{3}$. For height, they are $\mathsf{H}^* = 1$ and $\Phi(\Delta m/\sqrt{\sigma_{\mathfrak{M}}^2 + \sigma_{\mathfrak{W}}^2}) \approx 0.91$. Caring *less* about status produces *more* marrying up in the first case. Preferences and outcomes are different objects.

8 Main Result

Theorem 8.1 (Hypergamy Is Solved)**.** *For any marriage market:*

(a) universal hypergamy is feasible iff $\mu_s \succeq_{\mathrm{FSD}} \nu_s$;

(b) otherwise the minimal unmatched mass is $\mathsf{D} = \sup_t (G_{\mathfrak{W}} - G_{\mathfrak{M}})^+$;

(c) no strategy changes (a) or (b), and for continuous $F_{\mathfrak{M}}, F_{\mathfrak{W}}$ maxxing is zero-sum in rank;

(d) the equilibrium is the comonotone coupling, with index $\mathsf{H}^* = \mathrm{Leb}\{F_{\mathfrak{M}}^{-1} > F_{\mathfrak{W}}^{-1}\}$, under strictly supermodular transferable utility and under status-ranked preferences in the finite markets of Theorem 5.1;

(e) when taste shocks of scale θ are added to the transferable utility of (d), with finitely many types and no singles, the realized index tends to H^* as $\theta \downarrow 0$ and to $\mathbb{P}(S_{\mathfrak{M}} > S_{\mathfrak{W}})$ as $\theta \to \infty$.

Every quantity in (a)–(d) is a functional of (μ_s, ν_s) alone, and (e) adds only θ and the surplus f. No other behavioral input is required.

Proof. Theorems 2.1, 2.2, 4.3, 4.5, 5.1, 6.1, and 7.1. □

Remark 8.2 (Interpretation). When one side's status distribution overtakes the other's, measured hypergamy falls even if no one's preferences change. This matches the reversal of the gender gap in education documented by Esteve et al. [4], who report a global shift from educational hypergamy toward hypogamy.

9 Beyond Height: Multidimensional Status

Height is one coordinate of a longer list. Let each man carry a trait vector $x = (x_1, \ldots, x_k)$ (height, face, confidence, charisma, and so on), each coordinate a within-sex z-score, so that μ is now a distribution on $\mathbb{R}^k$. A requirement can combine traits in two ways: as a checklist, where every trait must clear its own bar, or as a trade-off, where strength in one trait offsets weakness in another.

Theorem 9.1 (The Checklist Theorem)**.** *Suppose a fraction α of $\mathfrak{W}$ requires every trait to clear a bar, $x \in A = \{x_1 \geq \tau_1, \ldots, x_k \geq \tau_k\}$. Then at least $(\alpha - \mu(A))^+$ of $\mathfrak{W}$ is unmatched in every feasible matching, and for independent traits $\mu(A) = \prod_i \mu(x_i \geq \tau_i)$.*

Proof. The constrained women's neighborhood lies inside A, of mass $\mu(A)$. Apply the Hall bound from Theorem 2.2, exactly as in Theorem 3.4. □

Example 9.2. In the calibration of Section 3, six feet is $x_1 \geq 1$, so $\mu(x_1 \geq 1) \approx 0.16$. Adding a top-20% face and top-30% confidence leaves $\mu(A) \approx 0.16 \times 0.2 \times 0.3 \approx 0.01$ for independent traits. With a common pairwise correlation ρ between traits (a one-factor Gaussian model), $\mu(A)$ rises to 0.032 at $\rho = 0.3$ and 0.054 at $\rho = 0.5$. For independent traits, bars multiply: every requirement added shrinks the eligible pool by its own pass rate. Correlation softens this without removing it: at $\rho = 0.5$ the face bar keeps 46% of the six-footers, and the confidence bar 73% of those left.

Proposition 9.3 (Compensation)**.** *If instead $\mathfrak{W}$ ranks men by an index $s(x) = \beta \cdot x$, then s is a scalar status functional, and Theorem 8.1 applies to it once s is also defined on $\mathfrak{W}$ on the same scale. For independent standard Gaussian traits, the fraction of the top $\Phi(-c)$ of men by index who fall below c on trait i is*

$$1 - \frac{\Phi_2(-c, -c; r_i)}{\Phi(-c)}, \qquad r_i = \frac{\beta_i}{\|\beta\|}, \tag{11}$$

where $\Phi_2(\cdot, \cdot; r)$ is the standard bivariate normal CDF with correlation r.

Proof. $(x_i, s/\|\beta\|)$ is standard bivariate normal with correlation r_i, and the top $\Phi(-c)$ of men by index is the event $s/\|\beta\| \geq c$. □

Remark 9.4. With equal weights on height, face and confidence ($r_i = 1/\sqrt{3}$) and $c = 1$, take the top 16% of men by index, as many men as are six feet tall: 56% of them are under six feet. Under a checklist six feet is a gate; under a trade-off it is one input among several. The Six-Foot Theorem is a statement about the checklist.

Nothing requires the two sides to be scored on the same traits. Let $\mathfrak{W}$ rank men by an index $s_{\mathfrak{M}}$ and $\mathfrak{M}$ rank women by a different index $s_{\mathfrak{W}}$. Scores on different traits share no unit, so compare positions instead: write $U = F_{\mathfrak{M}}(s_{\mathfrak{M}}(x))$ and $V = F_{\mathfrak{W}}(s_{\mathfrak{W}}(y))$ for each person's quantile within their own side.

Theorem 9.5 (Quantile Hypergamy)**.** *In a balanced market with continuous score distributions, if every member of $\mathfrak{W}$ requires $U \geq V + \delta$, where $0 \leq \delta \leq 1$, the minimal unmatched mass of $\mathfrak{W}$ is exactly δ, whatever traits define $s_{\mathfrak{M}}$ and $s_{\mathfrak{W}}$.*

Proof. By the probability integral transform U and V are both $\mathrm{Unif}(0, 1)$. Shifting by δ as in Section 3, Theorem 2.2 gives $\mathsf{D}(\delta) = \sup_{t \in [0,1]} \big((1 - t) - (1 - t - \delta)^+\big)^+ = \delta$, attained at every $t \in [0, 1 - \delta]$. □

Remark 9.6. At $\delta = 0$ the two quantile distributions coincide, so by Corollary 4.4 no strictly hypergamous matching exists. Once status means rank, a market in which everyone marries strictly up is not rare but impossible: by Lemma 4.1, every rank gained is a rank lost elsewhere.

10 Partner History

A common claim holds that fewer previous partners make for better relationships, either because the capacity to pair-bond gets used up or because a partner with no history has no one to compare you to. The data support an association, but not one confined to women. Among 2,654 married individuals, Busby, Willoughby and Carroll [2] find that more lifetime partners go with lower sexual quality, poorer communication and lower stability, with no significant difference between men and women. Using longitudinal data on both sexes, Smith and Wolfinger [9] find divorce risk highest with six or more premarital partners and next highest with one or two, while three to five do not differ

significantly from none besides the eventual spouse; the link survives controls for religion, values, family relationships, personality and mental health, shows no gender difference, and selection is not ruled out.

Wolfinger [12] reports the lowest divorce rates for women with zero or one premarital partners, an irregular pattern above that, and religiosity as part of the explanation for the low end; that analysis covers only women because the survey lacks full data on men's premarital sex. The neurobiology of pair bonding comes from studies of monogamous rodents [13], and its human version, in which bonding capacity is used up by previous partners, has not been established. We therefore treat partner history as a hypothesis, and treat it symmetrically.

Definition 10.1 (Comparison)**.** A person whose previous partners had qualities $Q_1, \dots, Q_n$, drawn i.i.d. from a distribution F with finite mean, values a new partner of quality q at $q - \lambda(M_n - q)^+$, where $M_n = \max_{i \le n} Q_i$, $M_0 = -\infty$, and $\lambda \ge 0$ measures how much comparison matters. Only an ex who was better than the new partner costs anything.

Proposition 10.2 (Diminishing Comparisons)**.** *The expected regret $R_n(q) = \mathbb{E}[(M_n - q)^+]$ is increasing and concave in n, with $R_0 = 0$ and*

$$R_{n+1}(q) - R_n(q) = \int_q^\infty F(t)^n \big(1 - F(t)\big)\, dt. \tag{12}$$

For standard Gaussian quality and an average new partner ($q = 0$), $R_n = 0,\ 0.40,\ 0.68,\ 0.89,\ 1.17,\ 1.54,\ 1.87$ at $n = 0,\ 1,\ 2,\ 3,\ 5,\ 10,\ 20$.

Proof. $R_n(q) = \int_q^\infty \mathbb{P}(M_n > t)\, dt = \int_q^\infty \big(1 - F(t)^n\big)\, dt$, which is finite because F has a finite mean. Subtract consecutive terms; the integrand $F^n(1 - F)$ is nonnegative and decreasing in n because $0 \le F \le 1$. □

Remark 10.3. The first partner sets the bar and each later one raises it by less, so the largest single step is from none to one, though later partners still add most of the total: the first supplies 45% of R_3 and 21% of R_{20}. This fits Wolfinger's jump from one premarital partner, usually the husband, to two [12], but not his higher divorce rate at two than at three to nine, nor Smith and Wolfinger [9], whose results suggest a link that grows stronger, not weaker, as partners accumulate. The model cannot tell comparison apart from selection, and nothing in it refers to sex: a man with many previous partners compares too. If partner history enters a status index, Theorem 4.5 applies unchanged: it re-sorts who ranks where, and every rank one person gains, someone else loses.

11 Conclusion

Hypergamy reduces to optimal transport on the real line. Its feasibility is stochastic dominance, its failure is the Hypergamy Deficit, its equilibrium is the quantile coupling, and its realized rate is the measure of the set where one quantile function exceeds another. Section 9 extends the deficit to several traits, where checklist requirements multiply when traits are independent and trade-offs reduce to a scalar index, and Section 10 treats partner history as a comparison effect that applies to both sexes. Open problems include general multidimensional status (Brenier maps and upper sets in a partial order), search frictions, and the case in which the status functional is itself endogenous.

Acknowledgments. The author thanks Zhang et al. for contributions that were substantial but unspecified. *Conflict of interest:* the author is a participant in the market under study.

References

[1] G. S. Becker, A theory of marriage: Part I, *Journal of Political Economy* 81(4), 1973.

[2] D. M. Busby, B. J. Willoughby, and J. S. Carroll, Sowing wild oats: Valuable experience or a field full of weeds? *Personal Relationships* 20(4), 2013.

[3] E. Choo and A. Siow, Who marries whom and why, *Journal of Political Economy* 114(1), 2006.

[4] A. Esteve, C. R. Schwartz, J. Van Bavel, I. Permanyer, M. Klesment, and J. García-Román, The end of hypergamy: Global trends and implications, *Population and Development Review* 42(4), 2016.

[5] C. D. Fryar, M. D. Carroll, Q. Gu, J. Afful, and C. L. Ogden, Anthropometric reference data for children and adults: United States, 2015–2018, *Vital and Health Statistics* Series 3, No. 46, National Center for Health Statistics, 2021.

[6] D. Gale and L. S. Shapley, College admissions and the stability of marriage, *American Mathematical Monthly* 69(1), 1962.

[7] A. Galichon and B. Salanié, Cupid's invisible hand: Social surplus and identification in matching models, *Review of Economic Studies* 89(5), 2022.

[8] L. S. Shapley and M. Shubik, The assignment game I: The core, *International Journal of Game Theory* 1(1), 1971.

[9] J. Smith and N. H. Wolfinger, Re-examining the link between premarital sex and divorce, *Journal of Family Issues* 45(3), 2024.

[10] V. Strassen, The existence of probability measures with given marginals, *Annals of Mathematical Statistics* 36(2), 1965.

[11] C. Villani, *Optimal Transport: Old and New*, Springer, 2009.

[12] N. H. Wolfinger, Counterintuitive trends in the link between premarital sex and marital stability, Institute for Family Studies, June 6, 2016.

[13] L. J. Young and Z. Wang, The neurobiology of pair bonding, *Nature Neuroscience* 7(10), 2004.