How BARCS Finds Important Genes

A student guide with worked examples and a quiz

Prepared for Hyun-Hwan Jeong and Kyu-Won Lee

BARCS version 0.2.1 · Reading level: grades 7 to 9

About this guide. This guide explains how the BARCS computer program works. You need only arithmetic, fractions, percents, powers and square roots. The guide teaches logarithms when it uses them. Each section uses the same small set of numbers, so you can follow every step. The guide uses the writing rules of ASD-STE100 Simplified Technical English.

Contents

1 The experiment: a CRISPR screen **4**

2 The data **4**

3 Step 1: Turn counts into proportions **5**

4 Step 2: Compare the groups with a fold change **5**

4.1 Logarithms make changes symmetric 6

5 Step 3: Odds and log-odds **7**

6 Step 4: Two kinds of noise **7**

7 Step 5: The design matrix **8**

7.1 What a design matrix is 8

7.2 From the design matrix to a prediction 9

7.3 From log-odds back to counts 10

7.4 The reference group 10

7.5 Paired replicates: add a column 10

7.6 Numbers instead of groups . . . 11
7.7 Two factors and an interaction . . . 11
7.8 Comparing two treatments: a contrast . . . 12
7.9 Rules for a good design matrix . . . 12
7.10 The design matrix in R . . . 13

8 Step 6: How BARCS finds the best β values 13
8.1 The loop . . . 13
8.2 The formulas . . . 14
8.3 Iteration 1 in full . . . 15
8.4 The next iterations . . . 15

9 Step 7: How BARCS measures the biological noise 16
9.1 Pearson residuals . . . 16
9.2 The rule for ρ . . . 16

10 Step 8: Is the effect real? 17
10.1 The standard error . . . 17
10.2 The t value . . . 18
10.3 Degrees of freedom . . . 18
10.4 The p-value . . . 18

11 Step 9: Borrow information from other guides 19
11.1 What moderation does to the test . . . 20

12 Step 10: Check the results with control guides 20

13 Step 11: From guides to genes 21
13.1 When guides share noise . . . 22

14 Step 12: Many genes and false discoveries 22

15 Two special cases 23
15.1 A guide with zero reads . . . 23
15.2 A very large change . . . 23

16 Summary: the full path **23**

17 Quiz **25**

18 Answers and calculations **30**

1 The experiment: a CRISPR screen

Each cell in your body has about 20,000 genes. A gene is a set of instructions. Scientists want to know which genes a cell needs, for example to survive a cancer drug.

A **CRISPR screen** tests thousands of genes at the same time. It works in four steps:

1. Scientists make a **library** of short pieces of RNA, called **guides**. Each guide switches off one gene.
2. They put one guide into each cell. Now each cell has one gene switched off.
3. They grow the cells. Some cells grow under normal conditions (**control**). Other cells grow with a drug (**drug**).
4. A sequencing machine reads the guides in each group. The machine counts how many times it reads each guide. This number is a **read count**.

If cells need a gene to survive the drug, the cells that lose that gene die. Their guide becomes rare in the drug group. So the guide count goes down.

Remember

BARCS does not look at cells directly. It looks at read counts. A guide that becomes rare tells us that its gene is important.

Scientists repeat each group more than one time. Each repeat is a **replicate**. Each sample that the machine reads is a **library**. In this guide, we use four libraries: two control replicates and two drug replicates.

2 The data

The machine gives two numbers for each library:

- the read count of our guide, and
- the total read count of all guides in that library.

Table 1 shows the data for one guide. We use these numbers in all sections of this guide.

What the symbols mean

K	The read count of one guide in one library. Library 1 has $K = 180$.
N	The total read count of all guides in one library. Library 1 has $N = 1{,}000{,}000$.
i	The library number (1, 2, 3 or 4). We write K_i for the count in library i. So $K_2 = 240$.

Table 1: Read counts for one guide in four libraries.

Library	Group	Guide count K	Total count N
1	control	180	1,000,000
2	control	240	1,200,000
3	drug	120	1,000,000
4	drug	80	800,000

Remember

Always record N before you remove any guides from the table. If you remove guides first, the totals become wrong. Then all the later numbers become wrong too.

3 Step 1: Turn counts into proportions

Library 2 has more reads than library 1 because the machine read it more. So we cannot compare raw counts. We divide each count by its library total. The result is a **proportion**:

$$p = \frac{K}{N}.$$

Proportions are very small numbers. So we multiply by one million. The result is the count **per million** reads.

Worked example 1: proportions

Library 1: $p = 180 \div 1{,}000{,}000 = 0.000180$ = 180 per million
Library 2: $p = 240 \div 1{,}200{,}000 = 0.000200$ = 200 per million
Library 3: $p = 120 \div 1{,}000{,}000 = 0.000120$ = 120 per million
Library 4: $p = 80 \div 800{,}000 = 0.000100$ = 100 per million

The two control libraries give 180 and 200 per million. The two drug libraries give 120 and 100 per million. The guide is less common with the drug.

What the symbols mean

p The proportion of all reads that belong to the guide. It is always between 0 and 1.

4 Step 2: Compare the groups with a fold change

We want one number that tells how much the guide changes. First, we combine the replicates of each group. We add their counts and add their totals:

$$p_{\text{control}} = \frac{180 + 240}{1{,}000{,}000 + 1{,}200{,}000} = \frac{420}{2{,}200{,}000} = 0.000191 \text{ (191 per million)},$$

$$p_{\text{drug}} = \frac{120 + 80}{1{,}000{,}000 + 800{,}000} = \frac{200}{1{,}800{,}000} = 0.000111 \text{ (111 per million)}.$$

The **fold change** is the drug proportion divided by the control proportion:

$$\text{fold change} = \frac{p_{\text{drug}}}{p_{\text{control}}} = \frac{0.000111}{0.000191} = 0.58.$$

A fold change of 0.58 means that the guide keeps only 58% of its control level. It loses 42%.

Remember

- Fold change $= 1$: no change.
- Fold change < 1: the guide becomes rarer. Cells may need this gene.
- Fold change > 1: the guide becomes more common. Losing this gene may help the cells.

4.1 Logarithms make changes symmetric

A fold change of 2 means "double". A fold change of 0.5 means "half". These changes have the same size, but the numbers 2 and 0.5 do not look equal. A **logarithm** fixes this problem.

The **base-2 logarithm**, $\log_2$, tells how many times you double.

Fold change	0.25	0.5	1	2	4
$\log_2$(fold change)	-2	-1	0	1	2

Now "double" is $+1$ and "half" is -1. The sizes look equal.

Worked example 2: log fold change

Our fold change is 0.58. It is between 0.5 and 1. So $\log_2(0.58)$ is between -1 and 0.
A calculator gives $\log_2(0.58) = -0.78$.
The guide drops by 0.78 "halvings".

BARCS uses the **natural logarithm**, ln. It counts powers of the number $e = 2.718\ldots$ instead of powers of 2. The idea is the same. To change back from ln, you raise e to that power: if $\ln(x) = b$, then $x = e^b$. For our guide, $\ln(0.58) = -0.54$.

5 Step 3: Odds and log-odds

BARCS works with **odds** instead of proportions. The odds compare the reads that belong to the guide with the reads that do not:

$$\text{odds} = \frac{p}{1-p}.$$

The **log-odds** is the natural logarithm of the odds:

$$\text{log-odds} = \ln\left(\frac{p}{1-p}\right).$$

Worked example 3: odds and log-odds

Control: $p = 0.000191$. So odds $= 0.000191 \div 0.999809 = 0.000191$.
The odds and the proportion are almost equal. This is true for all small proportions, because $1 - p$ is almost 1.
Log-odds for control: $\ln(0.000191) = -8.56$.
Log-odds for drug: $\ln(0.000111) = -9.10$.
Difference: $-9.10 - (-8.56) = -0.54$.
The difference of the log-odds is -0.54. This is the same as ln(fold change) from Step 2.

Remember

For guides, a difference in log-odds is almost exactly the natural log of the fold change. So you can read BARCS results as log fold changes.

Why use log-odds at all? A proportion must stay between 0 and 1. A log-odds can be any number, positive or negative. This property lets BARCS add effects together safely, as Step 5 shows.

6 Step 4: Two kinds of noise

The two control libraries gave 180 and 200 per million. They are not equal. Replicates always differ a little. We call this difference **noise**. BARCS must know how big the noise is. Then it can tell a real change from a chance change.

Guide counts have two kinds of noise:

1. **Counting noise.** The machine reads a random sample of all the guides. Think of a bag with a million marbles. You take out a handful. You do not get exactly the same number of red marbles each time.
2. **Biological noise.** Each replicate is a different group of living cells. The cells grow a little differently each time.

BARCS measures the size of the noise with the **variance**. The variance is the average squared distance of a value from its expected value. The **standard deviation** is the square root of the variance. It is the typical distance in the same units as the counts.

BARCS uses this formula for the variance of a guide count:

$$\text{variance} = \underbrace{N\,\mu\,(1-\mu)}_{\text{counting noise}} \times \underbrace{\big(1+(N-1)\,\rho\big)}_{\text{extra biological noise}}.$$

What the symbols mean

μ	"mu". The true proportion of the guide that we expect. For the control group, our best value is $\mu = 0.000191$.
N	The library total, as before.
ρ	"rho". A number that measures the biological noise. If $\rho = 0$, only counting noise exists. A bigger ρ means more biological noise.

Worked example 4: variance

Use N = 1,000,000 and μ = 0.000191. For our guide, BARCS estimates ρ = 0.000000386.
Counting noise: $N\mu(1-\mu) = 1{,}000{,}000 \times 0.000191 \times 0.999809 = 191$.
Extra factor: $1 + (N-1)\rho = 1 + 999{,}999 \times 0.000000386 = 1.39$.
Total variance: $191 \times 1.39 = 265$.
Standard deviation: $\sqrt{265} = 16.3$ reads.
So a count of about 191 normally moves up or down by about 16. Biological noise adds 39% to the counting noise.

Remember

Look at the extra factor $1 + (N-1)\rho$. It contains N. When the machine reads more, N grows, and the factor grows too. So more reading does not remove biological noise. Only more replicates help with biological noise.

7 Step 5: The design matrix

A **model** is a rule that predicts a value. BARCS predicts the log-odds of a guide in each library. The model needs to know which group each library belongs to. BARCS gets this information from the **design matrix**.

7.1 What a design matrix is

A design matrix is a table of numbers. It has:

- one **row** for each library, and
- one **column** for each β value in the model.

We write the design matrix with the letter X.

For our four libraries, the model has two β values. So X has four rows and two columns:

Library	Group	Column 1: "intercept"	Column 2: "drug"
1	control	1	0
2	control	1	0
3	drug	1	1
4	drug	1	1

What the symbols mean

X	The design matrix. One row for each library, one column for each β.
Intercept	A column of 1s. It belongs to β_0, the starting level. Every library gets the starting level.
Drug	A column that is 1 for a drug library and 0 for a control library. It belongs to β_1, the effect of the drug.
β_0	"beta zero". The log-odds of the guide in the control group.
β_1	"beta one". The change in log-odds from control to drug. This is the **effect**. It is the most important number.

7.2 From the design matrix to a prediction

To get the prediction for one library, take the row of that library. Multiply each number in the row by its β. Then add the products:

$$\text{predicted log-odds} = (\text{column 1}) \times \beta_0 + (\text{column 2}) \times \beta_1.$$

This calculation is a **matrix multiplication**. You do it one row at a time. We call the predicted log-odds η ("eta").

Worked example 5: predictions from the design matrix

BARCS finds $\beta_0 = -8.565$ and $\beta_1 = -0.542$ (Step 6 shows how).

Library 1 (row 1, 0): $1 \times (-8.565) + 0 \times (-0.542) = -8.565$
Library 2 (row 1, 0): $1 \times (-8.565) + 0 \times (-0.542) = -8.565$
Library 3 (row 1, 1): $1 \times (-8.565) + 1 \times (-0.542) = -9.107$
Library 4 (row 1, 1): $1 \times (-8.565) + 1 \times (-0.542) = -9.107$

The control libraries get β_0 only. The drug libraries get $\beta_0 + \beta_1$. So β_1 is the difference between the groups.

7.3 From log-odds back to counts

A prediction in log-odds is hard to read. So we change it back into a proportion, and then into a count. The rule is:

$$\mu = \frac{e^{\eta}}{1 + e^{\eta}}, \qquad \text{expected count} = N \times \mu.$$

This rule undoes the log-odds of Step 3.

Worked example 6: expected counts

Control libraries ($\eta = -8.565$): $e^{-8.565} = 0.000190676$. Then $\mu = 0.000190676 \div 1.000190676 = 0.000190640$, which is 190.6 per million.
Drug libraries ($\eta = -9.107$): $e^{-9.107} = 0.000110840$. Then $\mu = 0.000110827$, which is 110.8 per million.

Library	N	μ (per million)	Expected count $N\mu$	Real count K
1	1,000,000	190.6	190.6	180
2	1,200,000	190.6	228.8	240
3	1,000,000	110.8	110.8	120
4	800,000	110.8	88.7	80

The expected counts are near the real counts. The differences are -10.6, $+11.2$, $+9.2$ and -8.7 reads. These differences are the noise.

7.4 The reference group

The control group has 0 in the drug column. So control is the **reference group**. The effect β_1 always compares a group with the reference group. If you make drug the reference group, β_1 changes sign. It becomes $+0.542$. The meaning stays the same.

7.5 Paired replicates: add a column

Sometimes the replicates are pairs. For example, replicate 1 of control and replicate 1 of drug come from the same batch of cells. Then BARCS can remove the batch difference. You add a column for the batch:

Library	Group	Replicate	Intercept	Replicate 2	Drug
1	control	1	1	0	0
2	control	2	1	1	0
3	drug	1	1	0	1
4	drug	2	1	1	1

Now the model has three β values: the starting level, the batch difference and the drug effect. Each new column costs one degree of freedom (Step 8). With four libraries and three columns, only one degree of freedom stays.

Worked example 7: the paired model for our guide

BARCS gives the drug effect $\beta = -0.544$ with standard error 0.143. The effect is almost the same as before (-0.542). But the degrees of freedom drop from 2 to 1. The p-value becomes 0.164 instead of 0.033.
Our two replicates have almost no batch difference ($\beta = 0.005$). So the extra column costs information and gives nothing back. Add a batch column only when the replicates really are pairs.

7.6 Numbers instead of groups

A column can also hold a number, for example the day of the experiment. Suppose that you collect cells on day 0, day 7 and day 14, two libraries each. You put the day divided by 14 into a column:

Library	Day	Intercept	Day ÷ 14
1, 2	0	1	0
3, 4	7	1	0.5
5, 6	14	1	1

Now β_1 is the change in log-odds over 14 days. The model draws a straight line through the three days.

Worked example 8: a time course

Suppose that $\beta_1 = -0.70$.
After 14 days: change $= 1 \times (-0.70) = -0.70$. Fold change $= e^{-0.70} = 0.50$. The guide drops to half.
After 7 days: change $= 0.5 \times (-0.70) = -0.35$. Fold change $= e^{-0.35} = 0.70$. The guide drops to 70%.
The model has 6 libraries and 2 columns. So it has $6 - 2 = 4$ degrees of freedom.

7.7 Two factors and an interaction

Some experiments change two things. For example, cells can be normal or can have a second gene switched off (“knockout”). And cells can get no drug or the drug. This experiment has four groups. The design matrix has four columns:

Group	Intercept	Knockout	Drug	Knockout × Drug
normal, no drug	1	0	0	0
knockout, no drug	1	1	0	0
normal, drug	1	0	1	0
knockout, drug	1	1	1	1

The last column is the product of the knockout column and the drug column. It is 1 only when both are 1. Its β is the **interaction**. The interaction measures the extra change that happens only when both changes happen together.

Worked example 9: reading an interaction

Suppose that the four β values are -8.5, -0.2, $+0.1$ and -0.7. Multiply each row by the β values and add:

normal, no drug: -8.5

knockout, no drug: $-8.5 + (-0.2) = -8.7$

normal, drug: $-8.5 + 0.1 = -8.4$

knockout, drug: $-8.5 + (-0.2) + 0.1 + (-0.7) = -9.3$

The knockout alone changes the log-odds by -0.2. The drug alone changes it by $+0.1$. If the two effects only added up, the last group would be $-8.5 - 0.2 + 0.1 = -8.6$. The real value is -9.3. The extra -0.7 is the interaction. The guide drops strongly only when both changes happen.

7.8 Comparing two treatments: a contrast

Suppose an experiment has three groups: control, drug A and drug B. The design matrix has the columns intercept, drug A and drug B. Then β_A compares drug A with control, and β_B compares drug B with control. To compare drug A with drug B, you subtract: $\beta_A - \beta_B$. This calculation is a **contrast**. In BARCS, the function `bb_contrast()` calculates it and its test.

For example, if $\beta_A = -0.9$ and $\beta_B = -0.4$, then $\beta_A - \beta_B = -0.5$. Drug A lowers the guide more than drug B.

7.9 Rules for a good design matrix

1. Make one row for each library, in the same order as the count table.
2. Make one column for each β that you want.
3. Keep more rows than columns. The difference is the degrees of freedom. Two or more is better than one.
4. Do not let two columns carry the same information. For example, suppose that all drug libraries come from batch 2. Suppose also that all control libraries come from batch 1.

Then the batch column and the drug column are equal. BARCS cannot separate the drug effect from the batch effect.

7.10 The design matrix in R

In R, you describe the design matrix with a **formula**. The formula `~ group` means "an intercept and a column for each group except the reference group".

The function `model.matrix()` builds X from the formula:

```
> data <- data.frame(group = c("control", "control", "drug", "drug"))
> model.matrix(~ group, data)
  (Intercept) groupdrug
1           1         0
2           1         0
3           1         1
4           1         1
```

Then you give the counts, the totals, the formula and the column name to BARCS:

```
fit <- bb_screen(counts, data, formula = ~ group,
                 term = "groupdrug", totals = totals)
```

For the paired design, use the formula `~ replicate + group`. For the interaction, use `~ knockout * drug`.

8 Step 6: How BARCS finds the best β values

BARCS cannot find β_0 and β_1 in one calculation. It uses a loop of "guess, check, improve". The name of this method is **iteratively reweighted least squares** (IRLS). Each pass through the loop is an **iteration**.

8.1 The loop

1. **Guess.** Start with the simple log-odds of each group (Step 3).
2. **Predict.** Use the design matrix to predict η and μ for each library (Step 5).
3. **Measure the noise.** Find ρ from the differences between the real and the expected counts (Step 7).
4. **Weigh.** Give each library a **weight**. A library with less noise gets a larger weight.
5. **Correct.** Calculate a corrected value, the **working value**, for each library.
6. **Improve.** Take a weighted average of the working values in each group. These averages give the new β values.
7. **Repeat** from part 2 until the β values stop changing.

8.2 The formulas

The weight of library i is

$$w_i = \frac{N_i \, \mu_i \, (1 - \mu_i)}{1 + (N_i - 1) \, \rho}.$$

The top is the counting noise turned upside down. The bottom is the extra biological noise from Step 4. So more noise gives a smaller weight.

The working value of library i is

$$\text{working value}_i = \eta_i + \frac{K_i/N_i - \mu_i}{\mu_i \, (1 - \mu_i)}.$$

It is the current prediction η_i plus a correction. The correction is large when the real proportion K_i/N_i is far from the expected proportion μ_i.

For our design matrix, the new β values are weighted averages:

$$\beta_0 = \frac{w_1 \, v_1 + w_2 \, v_2}{w_1 + w_2}, \qquad \beta_0 + \beta_1 = \frac{w_3 \, v_3 + w_4 \, v_4}{w_3 + w_4},$$

where v_i is the working value of library i. (For other design matrices, BARCS solves a small set of equations instead. The idea is the same.)

8.3 Iteration 1 in full

Worked example 10: iteration 1

Guess. From Step 3, the control log-odds is -8.563522 and the drug log-odds is -9.104869. So $\beta_0 = -8.563522$ and $\beta_1 = -9.104869 - (-8.563522) = -0.541347$.
Predict. The control libraries get $\mu = 0.000190909$. The drug libraries get $\mu = 0.000111111$.
Measure the noise. Step 7 gives $\rho = 0.0000003841$ for these predictions.
Weigh library 1. Top: $1{,}000{,}000 \times 0.000190909 \times 0.999809 = 190.87$.
Bottom: $1 + 999{,}999 \times 0.0000003841 = 1.3841$.
Weight: $w_1 = 190.87 \div 1.3841 = 137.90$.
Correct library 1. $K_1/N_1 - \mu_1 = 0.000180 - 0.000190909 = -0.0000109$.
$\mu_1(1 - \mu_1) = 0.000190873$.
Correction: $-0.0000109 \div 0.000190873 = -0.0572$.
Working value: $v_1 = -8.5635 + (-0.0572) = -8.6207$.
The same calculations for all four libraries give:

Library	Weight w_i	Working value v_i	$w_i \times v_i$
1 (control)	137.905	−8.62068	−1188.84
2 (control)	156.785	−8.51589	−1335.16
3 (drug)	80.269	−9.02486	−724.42
4 (drug)	67.988	−9.20488	−625.82

Improve.
Control: $\beta_0 = (-1188.84 - 1335.16) \div (137.905 + 156.785) = -2524.00 \div 294.69 = -8.564929$.
Drug: $\beta_0 + \beta_1 = (-724.42 - 625.82) \div (80.269 + 67.988) = -1350.24 \div 148.257 = -9.107414$.
So $\beta_1 = -9.107414 - (-8.564929) = -0.542486$.

8.4 The next iterations

BARCS repeats the loop with the new values. The changes become very small:

After iteration	β_0	β_1
0 (the guess)	−8.563522	−0.541347
1	−8.564929	−0.542486
2	−8.564934	−0.542492
3	−8.564934	−0.542492

After iteration 3, the values do not change in the first six decimal places. BARCS stops. The final values are $\beta_0 = -8.565$ and $\beta_1 = -0.542$.

Remember

The weighted average gives a slightly different answer from the simple log-odds of Step 3 (-0.5425 instead of -0.5413). The difference comes from the weights. Library 2 has the most reads and the least relative noise. So it counts the most.

9 Step 7: How BARCS measures the biological noise

Step 4 used $\rho = 0.000000386$. This section shows where the number comes from.

9.1 Pearson residuals

For each library, BARCS subtracts the expected count from the real count. It divides the result by the standard deviation from Step 4. The result is a **Pearson residual**:

$$\text{Pearson residual}_i = \frac{K_i - N_i\mu_i}{\sqrt{N_i\mu_i(1-\mu_i)\big(1+(N_i-1)\rho\big)}}.$$

A Pearson residual tells how many standard deviations a library is from its prediction.

9.2 The rule for ρ

If the noise model is correct, the squares of the Pearson residuals add up to about the degrees of freedom. So BARCS chooses ρ to make this statement exactly true:

$$\text{sum of the squared Pearson residuals} = \text{df}.$$

Worked example 11: start with $\rho = 0$

Use the expected counts from Step 5. With $\rho = 0$, the bottom of each residual is the counting noise only.

Library	$(K - N\mu)^2$	$N\mu(1-\mu)$	Squared residual
1	$(-10.64)^2 = 113.21$	190.60	0.594
2	$(+11.23)^2 = 126.16$	228.72	0.552
3	$(+9.17)^2 = 84.14$	110.82	0.759
4	$(-8.66)^2 = 75.03$	88.65	0.846
		Sum	2.751

The sum is 2.751. The degrees of freedom are 2. The sum is too big. So there is more noise than counting noise alone. We need $\rho > 0$.

Worked example 12: guess and check for ρ

A larger ρ makes the bottom of each residual larger. So the sum goes down. BARCS tries values until the sum equals 2:

ρ	Sum of squared residuals
0	2.751
0.0000001	2.506
0.0000002	2.302
0.0000003	2.129
0.000000386	2.000
0.0000005	1.851

The answer is $\rho = 0.000000386$. If the sum is already 2 or less at $\rho = 0$, BARCS uses $\rho = 0$.

The ratio $2.751 \div 2 = 1.376$ is also useful. It is the **variance inflation**. It tells that the guide has about 1.38 times more noise than counting noise alone. Step 9 uses this number.

10 Step 8: Is the effect real?

The effect is $\beta_1 = -0.542$. But replicates are noisy. Maybe the true effect is 0 and the noise made the drop. BARCS tests this question in four parts.

10.1 The standard error

The **standard error** (SE) tells how much the estimate of β_1 can change because of noise. A small SE means a precise estimate.

For our design matrix, the standard error comes from the weights of Step 6. Add the weights of each group:

$$W_{\text{control}} = w_1 + w_2, \qquad W_{\text{drug}} = w_3 + w_4.$$

Then

$$\text{SE}(\beta_1) = \sqrt{\frac{1}{W_{\text{control}}} + \frac{1}{W_{\text{drug}}}}.$$

More weight means less noise, so the SE becomes smaller. (For other design matrices, BARCS uses the inverse of the matrix $X^\top W X$. For our design, that calculation gives exactly this formula.)

Worked example 13: the standard error

The final weights are 137.524, 156.322, 79.955 and 67.737.
$W_{\text{control}} = 137.524 + 156.322 = 293.846$.
$W_{\text{drug}} = 79.955 + 67.737 = 147.692$.
$1 \div 293.846 = 0.003403$.
$1 \div 147.692 = 0.006771$.
Sum: $0.003403 + 0.006771 = 0.010174$.
SE $= \sqrt{0.010174} = 0.101$.
The control group has about twice the weight of the drug group. So most of the uncertainty comes from the drug group.

10.2 The t value

The **t value** compares the effect with its standard error:

$$t = \frac{\beta_1}{\text{SE}}.$$

A t value far from 0 means that the effect is large compared with the noise.

10.3 Degrees of freedom

The **degrees of freedom** (df) tell how much information we have about the noise:

$$\text{df} = (\text{rows of } X) - (\text{columns of } X) = (\text{number of libraries}) - (\text{number of } \beta \text{ values}).$$

Fewer degrees of freedom mean less information. Then BARCS needs a bigger t value before it accepts an effect as real.

Degrees of freedom	1	2	14	very many
$\lvert t \rvert$ needed for $p < 0.05$	12.71	4.30	2.14	1.96

10.4 The p-value

The **p-value** answers this question: "If the true effect is 0, how often does noise alone give a t value this far from 0?" A small p-value means that noise alone seldom gives such a result. Scientists often use 0.05 as a limit.

Worked example 14: testing the effect

1. $t = -0.542 \div 0.101 = -5.38$.
2. X has 4 rows and 2 columns. So $\text{df} = 4 - 2 = 2$.
3. The table shows that 2 degrees of freedom need $|t| > 4.30$. Our $|t| = 5.38$. It is larger than 4.30. So $p < 0.05$.
4. A computer gives the exact p-value: $p = 0.033$.

The p-value is less than 0.05. So the drop is probably not only noise.

Remember

The degrees of freedom come from the number of libraries, not from the number of reads. A library with 100 million reads still counts as one row of X.

11 Step 9: Borrow information from other guides

With only 2 degrees of freedom, the noise estimate of one guide is not reliable. But a screen has thousands of guides. Most guides have a similar amount of noise. So BARCS combines two estimates:

- the variance inflation of this guide (Step 7), and
- the typical variance inflation of guides with similar counts.

This method is **moderation**. BARCS takes a weighted average. Each estimate gets a number of "votes". The guide's own estimate gets its degrees of freedom as votes. The typical estimate gets the **prior degrees of freedom**, d_0. BARCS calculates d_0 and the typical value from all the guides in the screen.

$$\text{moderated inflation} = \frac{\text{df} \times \text{own inflation} + d_0 \times \text{typical inflation}}{\text{df} + d_0}.$$

Worked example 15: moderation with simple numbers

A guide has 3 degrees of freedom. Its own inflation is 4.0. The typical inflation is 1.5, with $d_0 = 12$ votes.

$$\text{moderated inflation} = \frac{3 \times 4.0 + 12 \times 1.5}{3 + 12} = \frac{12 + 18}{15} = \frac{30}{15} = 2.0.$$

The guide's value moves from 4.0 toward 1.5. It stops at 2.0.

11.1 What moderation does to the test

BARCS uses the moderated inflation in three places:

1. It multiplies the standard error by $\sqrt{\text{moderated inflation} \div \text{own inflation}}$.
2. It recalculates t with the new standard error.
3. It adds the d_0 votes to the degrees of freedom. So the test uses $\text{df} + d_0$.

Worked example 16: moderation for our guide

Our guide has own inflation 1.376 (Step 7) and 2 degrees of freedom. Suppose that the screen gives typical inflation 1.5 with $d_0 = 12$.
Moderated inflation: $(2 \times 1.376 + 12 \times 1.5) \div (2 + 12) = (2.751 + 18) \div 14 = 1.482$.
Factor for the SE: $\sqrt{1.482 \div 1.376} = \sqrt{1.077} = 1.038$.
New SE: $0.101 \times 1.038 = 0.105$.
New t: $-0.542 \div 0.105 = -5.18$.
New degrees of freedom: $2 + 12 = 14$.
With 14 degrees of freedom, $|t|$ must be larger than 2.14 for $p < 0.05$. Our $|t| = 5.18$.
The new p-value is 0.00014.
The SE became slightly larger, because the typical guide is a little noisier than our guide. But the degrees of freedom grew from 2 to 14. So the test became much more powerful.

BARCS uses moderation when it has 50 or more usable guides.

12 Step 10: Check the results with control guides

A good screen contains **non-targeting control guides**. These guides switch off no gene. So their true effect is 0.

If the noise model is correct, about 5% of control guides get $p < 0.05$ by chance. If many more control guides get $p < 0.05$, the noise model is too small. Then BARCS makes all the tests wider.

Worked example 17: calibration with control guides

A screen has 100 control guides. We expect about 5 of them to get $p < 0.05$. But 15 of them get $p < 0.05$. The noise is too small.

BARCS finds a **scale factor** that makes the control guides behave correctly. It sorts the 100 control guides by the size of $|t|$ (the value without its sign). It takes the value that 95 of the 100 guides stay below. Suppose that this value is 2.47. With many degrees of freedom, the t table says that this value should be 1.96. So the scale factor is $2.47 \div 1.96 = 1.26$. BARCS divides every t value by 1.26.

A gene with $t = 3.5$ now gets $t = 3.5 \div 1.26 = 2.78$. Its p-value becomes larger. This change protects you from false results.

The scale factor is never less than 1. So calibration can only make the results more careful.

13 Step 11: From guides to genes

Each gene usually has several guides, often 3 to 6. Each guide gives its own result. BARCS combines them into one result for the gene. It uses the **Stouffer method** in three parts.

1. Change each guide's p-value into a **z score**. A z score tells how many standard deviations a result is from 0. The sign of the z score follows the sign of the effect.
2. Add the z scores.
3. Divide the sum by the square root of the number of guides, m.

$$Z_{\text{gene}} = \frac{z_1 + z_2 + \cdots + z_m}{\sqrt{m}}.$$

To change a p-value into a z score, use the table of the normal distribution. A two-sided p-value of 0.05 belongs to $z = 1.96$. Some other pairs:

Two-sided p-value	0.317	0.134	0.046	0.012	0.0027
z score	1.0	1.5	2.0	2.5	3.0

If the effect is negative (the guide drops), the z score gets a minus sign.

Worked example 18: combining three guides

A gene has $m = 3$ guides. Their p-values are 0.046, 0.134 and 0.012. Their z scores are 2.0, 1.5 and 2.5.

$$Z_{\text{gene}} = \frac{2.0 + 1.5 + 2.5}{\sqrt{3}} = \frac{6.0}{1.732} = 3.46.$$

This Z gives $p = 0.00053$ for the gene. No single guide is that strong. Together, the three guides give strong evidence.

13.1 When guides share noise

Sometimes the guides of one gene share noise. Then they agree more than they should by chance. BARCS measures this sharing with the **guide correlation**, r. It makes the bottom of the fraction bigger:

$$Z_{\text{gene}} = \frac{z_1 + z_2 + \cdots + z_m}{\sqrt{m + m(m-1)\,r}}.$$

Worked example 19: shared noise

Use the same three guides and $r = 0.1$. The bottom becomes $\sqrt{3 + 3 \times 2 \times 0.1} = \sqrt{3.6} = 1.897$. So $Z_{\text{gene}} = 6.0 \div 1.897 = 3.16$, and $p = 0.0016$. The evidence is still strong, but less strong than before.

14 Step 12: Many genes and false discoveries

A screen tests about 20,000 genes. Suppose that no gene has an effect. If we use the limit $p < 0.05$, about 5% of the genes still pass by chance. That is about 1,000 false results.

So BARCS uses the **false discovery rate** (FDR). The FDR is the expected fraction of false results in your list of hits. BARCS calculates it with the **Benjamini–Hochberg method**:

1. Sort the genes from the smallest p-value to the largest.
2. Give each gene a rank: 1, 2, 3, and so on.
3. For each gene, calculate $p \times$ (number of genes) $\div$ rank.
4. Go from the bottom of the list to the top. If a value is larger than the value below it, replace it with the value below it.

Worked example 20: false discovery rate for five genes

Gene	Rank	p	$p \times 5 \div$ rank	FDR (after part 4)
A	1	0.001	$0.001 \times 5 \div 1 = 0.005$	0.005
B	2	0.008	$0.008 \times 5 \div 2 = 0.020$	0.020
C	3	0.039	$0.039 \times 5 \div 3 = 0.065$	0.051 (from D)
D	4	0.041	$0.041 \times 5 \div 4 = 0.051$	0.051
E	5	0.600	$0.600 \times 5 \div 5 = 0.600$	0.600

With the limit FDR < 0.05, genes A and B are hits. With the limit FDR < 0.10, genes A, B, C and D are hits.

15 Two special cases

15.1 A guide with zero reads


Sometimes a guide has 0 reads in all control libraries. Then the log-odds of the control group is $\ln(0)$. This value does not exist, because no power of e equals 0. Older versions of BARCS could not fit such a guide. So they removed it.

BARCS version 0.2.1 adds a very small number to the counts first. This number is less than one read. The method is the **Firth correction**. Now the guide gets a value, and BARCS keeps it.

15.2 A very large change

Sometimes a guide grows by hundreds of times. The usual test from Step 8 can then fail. The standard error grows faster than the effect, so the t value becomes small. BARCS version 0.2.1 has a second test, the **likelihood-ratio test**. It compares how well the model fits with the effect and without the effect. The option `test = "auto"` uses this test only for guides with very large changes.

16 Summary: the full path



All symbols in one table

Symbol	Meaning	Value in our example
K	read count of the guide in one library	180, 240, 120, 80
N	total read count of one library	1.0, 1.2, 1.0, 0.8 million
p	proportion, K/N	191 and 111 per million
fold change	$p_{\text{drug}} \div p_{\text{control}}$	0.58
$\log_2$	number of doublings	$\log_2(0.58) = -0.78$
ln	natural logarithm (powers of $e = 2.718$)	$\ln(0.58) = -0.54$
log-odds	$\ln(p/(1-p))$	-8.56 (control)
μ	expected proportion	0.000191 (control)
ρ	biological noise	0.000000386
X	design matrix: a row per library, a column per β	4 rows, 2 columns
β_0	control log-odds in the model	-8.565
β_1	effect of the drug	-0.542
η	predicted log-odds of a library (row of X times β)	-8.565, -9.107
w	weight of a library in the fit	137.5 (library 1)
working value	prediction plus correction (Example 10)	-8.621 (library 1)
Pearson residual	distance from the prediction in SDs	-0.65 (library 1)
inflation	noise compared with counting noise	1.376
SE	standard error of the effect	0.101
t	effect divided by SE	-5.38
df	libraries minus β values	$4 - 2 = 2$
p-value	chance of this result if the effect is 0	0.033
d_0	prior degrees of freedom (votes)	12 (Example 15)
z, Z	z score of a guide, of a gene	3.46 (Example 18)
m	number of guides for a gene	3
r	guide correlation (shared noise)	0.1 (Example 19)
FDR	expected fraction of false hits	0.005 for gene A

Note: the letter p has two uses in this guide. Alone, p is a proportion. In the word "p-value", it is a probability. The text always tells you which one it is.

17 Quiz

Choose the best answer. The answers and calculations are in the next section.

Q1. What does one guide do in a CRISPR screen?

(A) It counts the cells.

(B) It adds a drug to the cells.

(C) It reads the DNA.

(D) It switches off one gene.

Q2. A library has $K = 300$ and $N = 1{,}500{,}000$. What is the count per million?

(A) 200

(B) 300

(C) 450

(D) 500

Q3. Why do we divide each count by its library total N?

(A) To make the numbers bigger.

(B) Because the machine reads some libraries more than others.

(C) To remove the biological noise.

(D) Because BARCS cannot use whole numbers.

Q4. A guide has 400 per million in control and 100 per million with a drug. What is the fold change?

(A) 4

(B) 0.4

(C) 0.25

(D) 300

Q5. What is $\log_2(8)$?

(A) 2

(B) 3

(C) 4

(D) 8

Q6. A gene has $\log_2(\text{fold change}) = -1$. What happened to its guide?

(A) It doubled.

(B) It did not change.

(C) It dropped to half.

(D) It dropped to zero.

Q7. For a very small proportion p, which statement is true?

(A) The odds are almost equal to p.

(B) The odds are almost equal to 1.

(C) The odds are twice p.

(D) The odds are always 0.

Q8. In the variance formula, what does $\rho = 0$ mean?

(A) There is no noise at all.

(B) There is only counting noise.

(C) There is only biological noise.

(D) The guide has zero reads.

Q9. Use $N\mu(1-\mu)$ with $N = 100$ and $\mu = 0.5$. What is the counting variance?

(A) 25

(B) 50

(C) 0.25

(D) 100

Q10. The machine reads each library ten times more deeply. What happens to the biological noise?

(A) It disappears.

(B) It becomes ten times smaller.

(C) It does not go away. Only more replicates help.

(D) It becomes zero for control guides.

Q11. In the design matrix of our example, which β belongs to the column of 1s?

(A) β_1

(B) the interaction

(C) β_0

(D) ρ

Q12. BARCS gives $\beta_1 = 0.69$ for a guide. Use $e^{0.69} = 2.0$. What does this mean?

(A) The guide halves with the drug.

(B) The guide does not change.

(C) The guide disappears.

(D) The guide doubles with the drug.

Q13. An experiment has 6 libraries. The model has 3 β values. How many degrees of freedom does it have?

(A) 3
(B) 2
(C) 6
(D) 9

Q14. An effect is $\beta_1 = -0.8$ with SE $= 0.2$. What is the t value?

(A) -0.16
(B) -1.0
(C) -0.6
(D) -4.0

Q15. A guide has df $= 2$ and its own noise 5.0. The typical noise is 2.0 with $d_0 = 8$. What is the moderated noise?

(A) 2.0
(B) 3.5
(C) 5.0
(D) 2.6

Q16. A screen has 200 control guides. About how many should get $p < 0.05$ by chance?

(A) 0
(B) 5
(C) 10
(D) 50

Q17. A gene has 4 guides, each with $z = 1.0$. What is Z_{gene} (no shared noise)?

(A) 1.0
(B) 2.0
(C) 4.0
(D) 0.5

Q18. Three genes have p-values 0.01, 0.02 and 0.30. What is the Benjamini–Hochberg value $p \times 3 \div$ rank for the second gene?

(A) 0.02
(B) 0.06
(C) 0.30
(D) 0.03

Q19. Why does BARCS 0.2.1 add a very small number to the counts of some guides?

(A) Because $\ln(0)$ does not exist when a group has zero reads.

(B) To make all guides look important.

(C) To increase the library totals.

(D) To remove control guides.

Q20. Which action gives BARCS more information about biological noise?

(A) Read each library more deeply.

(B) Use fewer guides.

(C) Add more replicate libraries.

(D) Use a larger p-value limit.

Q21. An experiment has 3 control libraries and 3 drug libraries. The formula is `~ group`. What is the size of the design matrix?

(A) 3 rows and 2 columns

(B) 6 rows and 2 columns

(C) 2 rows and 6 columns

(D) 6 rows and 3 columns

Q22. A library has the row $(1, 1)$ in the design matrix. The values are $\beta_0 = -8.0$ and $\beta_1 = -0.5$. What is the predicted log-odds?

(A) -8.5

(B) -8.0

(C) -7.5

(D) 4.0

Q23. In the knockout-and-drug design, which group has a 1 in the interaction column?

(A) normal, no drug

(B) knockout, no drug

(C) normal, drug

(D) knockout, drug

Q24. Two control libraries have weights 100 and 300. Their working values are -8.0 and -9.0. What is the weighted average?

(A) -8.5

(B) -8.25

(C) -8.75

(D) -9.0

Q25. The control weights add up to $W_{\text{control}} = 100$. The drug weights add up to $W_{\text{drug}} = 25$. What is $\text{SE}(\beta_1)$?

(A) 0.05

(B) 0.224

(C) 0.5

(D) 1.25

Q26. At $\rho = 0$, the squared Pearson residuals of a guide add up to 1.5. The model has 2 degrees of freedom. Which ρ does BARCS use?

(A) $\rho = 0$

(B) $\rho = 1.5$

(C) $\rho = 0.75$

(D) $\rho = 2$

Q27. You add a paired-replicate column to the model of our four libraries. How many degrees of freedom stay?

(A) 3

(B) 2

(C) 4

(D) 1

Q28. Of 100 control guides, 95 have $|t|$ below 2.94. The t table says 1.96. What is the scale factor?

(A) 0.67

(B) 1.0

(C) 1.5

(D) 4.9

18 Answers and calculations

Q1. D. Each guide switches off one gene (Section 1).

Q2. A. $300 \div 1{,}500{,}000 = 0.0002$. Then $0.0002 \times 1{,}000{,}000 = 200$ per million.

Q3. B. Libraries get different numbers of reads. Division by N makes them comparable (Step 1).

Q4. C. $100 \div 400 = 0.25$. The guide keeps one quarter of its control level.

Q5. B. $2 \times 2 \times 2 = 8$. You double three times, so $\log_2(8) = 3$.

Q6. C. -1 means one halving (Step 2).

Q7. A. $1 - p$ is almost 1, so $p/(1-p)$ is almost p (Step 3).

Q8. B. The extra factor becomes $1 + (N-1) \times 0 = 1$. Only counting noise stays (Step 4).

Q9. A. $100 \times 0.5 \times (1 - 0.5) = 100 \times 0.5 \times 0.5 = 25$.

Q10. C. The factor $1 + (N-1)\rho$ grows with N. Deeper reading does not remove biological noise (Step 4).

Q11. C. The column of 1s is the intercept. It belongs to β_0, the starting level (Step 5).

Q12. D. The fold change is $e^{0.69} = 2.0$. The guide doubles.

Q13. A. df $= 6 - 3 = 3$ (Step 8).

Q14. D. $t = -0.8 \div 0.2 = -4.0$.

Q15. D. $(2 \times 5.0 + 8 \times 2.0) \div (2 + 8) = (10 + 16) \div 10 = 26 \div 10 = 2.6$ (Step 9).

Q16. C. 5% of $200 = 0.05 \times 200 = 10$ (Step 10).

Q17. B. $(1.0 + 1.0 + 1.0 + 1.0) \div \sqrt{4} = 4.0 \div 2 = 2.0$ (Step 11).

Q18. D. The second gene has rank 2. So $0.02 \times 3 \div 2 = 0.03$ (Step 12).

Q19. A. With zero reads, the log-odds is $\ln(0)$, which does not exist. The Firth correction fixes this problem.

Q20. C. Degrees of freedom come from libraries. Only more replicates add information about biological noise (Steps 4 and 8).

Q21. B. One row for each library ($3 + 3 = 6$). One column for the intercept and one for drug (Step 5).

Q22. A. $1 \times (-8.0) + 1 \times (-0.5) = -8.5$ (Step 5).

Q23. D. The interaction column is knockout times drug. Only $1 \times 1 = 1$ (Step 5).

Q24. C. $(100 \times (-8.0) + 300 \times (-9.0)) \div (100 + 300) = (-800 - 2700) \div 400 = -3500 \div 400 = -8.75$ (Step 6).

Q25. B. $\sqrt{1/100 + 1/25} = \sqrt{0.01 + 0.04} = \sqrt{0.05} = 0.224$ (Step 8).

Q26. A. The sum (1.5) is already 2 or less. So BARCS uses $\rho = 0$ (Step 7).

Q27. D. The model now has 3 columns. So $\text{df} = 4 - 3 = 1$ (Step 5).

Q28. C. $2.94 \div 1.96 = 1.5$ (Step 10).