

Physical Chemistry (II)

Lecture 19

Multi-electron Atoms



Last class

Orbital Approximation

- Each electron's spatial part is represented by an one-electron orbital $\phi_i(\vec{r})$
- Hartree-Fock method provides the “best” orbitals
- Functional forms of orbitals?



Slater-Type Orbitals (STOs)

$$S_{nlm_l}(r, \theta, \phi; \zeta) = \frac{(2\zeta/a_0)^{3/2}}{\sqrt{(2n)!}} \left(\frac{2\zeta r}{a_0} \right)^{n-1} e^{-\zeta r/a_0} Y_l^{m_l}(\theta, \phi)$$

- ζ : orbital exponent
- Linear combination of STOs can be used to model orbitals
 - e.g. double-zeta orbital

$$\phi_{1s}(\vec{r}) = c_1 S_{1s}(\vec{r}; \zeta_1) + c_2 S_{1s}(\vec{r}; \zeta_2)$$

- Determination of ζ takes most of computational time
 - We can “preset” the values of ζ and optimize

Basis Set

- Atomic orbitals ψ can be represented by linear combinations of some functions ϕ_j

$$\psi(\vec{r}) = \sum_{j=1}^K c_j \phi_j = c_1 \phi_1 + c_2 \phi_2 + \cdots + c_K \phi_K$$

- Basis functions: ϕ_j (e.g. STOs)
- Basis set: $\{\phi_j\}$

Last class

Gaussian-Type Orbitals (GTOs)

$$G_{ijk}(r, \theta, \phi; \alpha) = Nr^{n-1}e^{-\alpha r^2}Y_l^m(\theta, \phi)$$

Computationally efficient

Can be combined to model orbitals

N-MPG Basis Sets

- *N*: number of Gaussian functions for each inner-shell electron orbital
- *M*, *P*: numbers of Gaussian functions for each valence-shell electron orbital

e.g. 3-21G for Lithium (d_i and α_i are preset)

$$\phi_{1s}(\vec{r}) = d_1 G_{1s}(\alpha_1) + d_2 G_{1s}(\alpha_2) + d_3 G_{1s}(\alpha_3)$$


$$\phi_{2s}(\vec{r}) = c_1 \phi_{2s,1}(\vec{r}) + c_2 \phi_{2s,2}(\vec{r})$$

$$\phi_{2s,1}(\vec{r}) = d_4 G_{2s}(\alpha_4) + d_5 G_{2s}(\alpha_5)$$

$$\phi_{2s,2}(\vec{r}) = G_{2s}(\alpha_6)$$

Last class

Size of Basis Set



Extension of HF

- Start from one-electron wavefunctions (“orbitals”) u_i

$$\phi = |u_a \quad u_b \quad \cdots \quad u_n|$$

- HF determinants use **occupied orbitals only**
- **Inclusion of unoccupied (virtual) orbitals** may increase accuracy and take care of correlation

Example: Virtual Orbital of He

- Double zeta orbital

$$\phi_{1s}(\vec{r}) = c_1 S_{1s}(\vec{r}; \zeta = 1.454) + c_2 S_{1s}(\vec{r}; \zeta = 2.911)$$

- Ground-state orbital (by Hartree-Fock)

$$\psi_0 = 0.844 S_{1s}(\zeta = 1.454) + 0.181 S_{1s}(\zeta = 2.911)$$

- Virtual orbital (orthogonal to ψ_0)

$$\psi_v = 1.624 S_{1s}(\zeta = 1.454) - 1.821 S_{1s}(\zeta = 2.911)$$

Last class

Example: Virtual Orbital of He

- Hartree-Fock wavefunction $\phi_{\text{HF}} = \psi_0(\vec{r}_1)\psi_0(\vec{r}_2)$
- Better wavefunction

$$\phi_{\text{CI}} = c_1\psi_0(\vec{r}_1)\psi_0(\vec{r}_2) + c_2\psi_v(\vec{r}_1)\psi_v(\vec{r}_2)$$

Using a variational method, one can obtain

$$\phi_{\text{CI}} = 0.999 \psi_0(\vec{r}_1)\psi_0(\vec{r}_2) - 0.0476 \psi_v(\vec{r}_1)\psi_v(\vec{r}_2)$$

with optimized energy $E = -2.87541 E_{\text{h}}$

(HF energy $E = -2.86167 E_{\text{h}}$)

Last class

Optimization of Wavefunction

Method		crude $\longrightarrow$ accurate	Applications
Exact solution			H only
Pure VM/PT		depending on complexity	Very few (He)
Orbital approx.	HF		All elements
	CI		All elements
	CC		All elements
	MP		All elements
DFT			All elements

Not in text

Now We Are Ready...




for **Hands-On Practice!**

Not in text


Atomic Orbitals

- **Angular part:** same with H atom (spherical harmonics)
- **Radial part:** slightly different from H atom (but not much)



(Image: <https://chem.libretexts.org/@go/page/4534>)

Orbital Energies (Experimental)



* e-e repulsion leads to *l*-splitting

Electron Configurations

- Distribution of electrons in **atomic orbitals**

e.g. $1s^2$, $1s^1 2s^1$, $1s^2 2s^2 2p^2$

- **Aufbau principle:** ground-state electron configurations

“Electrons first fill orbitals with the lowest available energy”

Electronic States of Atoms

- Ground-state electron configuration of C: $1s^2 2s^2 2p^2$
→ Same configurations can give “different” states
- Angular momentum is a key to the good description
 - For each electron i , we have $\vec{L}_i$ and $\vec{S}_i$ (and $\vec{J}_i = \vec{L}_i + \vec{S}_i$)
- Two possible ways to obtain quantum numbers
 - Russell-Saunders coupling: $\vec{L}_{\text{total}}$ and $\vec{S}_{\text{total}} \rightarrow \vec{J}_{\text{total}}$
 - j-j coupling: $\vec{J}_i$ for each electron $\rightarrow \vec{J}_{\text{total}}$

Russell-Saunders Coupling

- Total orbital angular momentum

$$\vec{L}_{\text{total}} = \sum_i \vec{L}_i$$

- Total spin angular momentum

$$\vec{S}_{\text{total}} = \sum_i \vec{S}_i$$

- Total angular momentum

$$\vec{J}_{\text{total}} = \vec{L}_{\text{total}} + \vec{S}_{\text{total}}$$

Case of Two Electrons

$$\vec{L}_{\text{total}} = \vec{L}_1 + \vec{L}_2, \quad \vec{S}_{\text{total}} = \vec{S}_1 + \vec{S}_2$$

For **nonequivalent electrons** (e.g. $1s^1 2s^1$), the possible quantum numbers are

$$L = (l_1 + l_2), (l_1 + l_2 - 1), \dots, |l_1 - l_2|$$

$$S = (s_1 + s_2), (s_1 + s_2 - 1), \dots, |s_1 - s_2|$$

$$J = (L + S), (L + S - 1), \dots, |L - S|$$

$$M_J = -J, -(J - 1), \dots, J$$

Atomic Term Symbol

$$^{2S+1}[L]_J$$

- Symbols for $[L]$

$L =$	0	1	2	3	4	...
	S	P	D	F	G	

- Closed (sub)shells do not contribute
→ only **open subshells** need to be counted

Two Nonequivalent Electrons

$$1s^1 2s^1: l_1 = 0, l_2 = 0, s_1 = 1/2, s_2 = 1/2$$

$$L = 0; S = 0, 1$$

- $L = 0, S = 0$:

$$J = 0 \rightarrow M_J = 0 (^1S_0)$$

- $L = 0, S = 1$:

$$J = 1 \rightarrow M_J = 0, \pm 1 (^3S_1)$$

Two Nonequivalent Electrons

$$2p^1 3p^1: l_1 = 1, l_2 = 1, s_1 = 1/2, s_2 = 1/2$$

$$L = 0, 1, 2; S = 0, 1$$

- $L = 0, S = 0: J = 0$ (1S_0)
- $L = 0, S = 1: J = 1$ (3S_1)
- $L = 1, S = 0: J = 1$ (1P_1)
- $L = 1, S = 1: J = 0, 1, 2$ ($^3P_0, ^3P_1, ^3P_2$)
- $L = 2, S = 0: J = 2$ (1D_2)
- $L = 2, S = 1: J = 1, 2, 3$ ($^3D_1, ^3D_2, ^3D_3$)

Equivalent Electrons

- **Pauli exclusion principle:** (a, a) impossible
- **Indistinguishability:** (a, b) = (b, a)

1. Construct a table for microstates

$$M_L = m_{l1} + m_{l2} + \cdots, \quad M_S = m_{s1} + m_{s2} + \cdots$$

2. Eliminate the states that violate the two conditions

$$\text{Total number of microstates} = G!/[N!(G - N)!]$$

3. Deduce the possible values of L and S (start from largest M_L and then largest M_S)



Two Equivalent Electrons

$$2p^2: l_1 = 1, l_2 = 1, s_1 = 1/2, s_2 = 1/2$$

$$L = 0, 1, 2; S = 0, 1$$

$$M_L = -2, -1, 0, 1, 2; M_S = -1, 0, 1$$

		M_S		
		1	0	-1
M_L	2	(1⁺, 1⁺)	(1 ⁺ , 1 ⁻)	(1⁻, 1⁻)
	1	(0 ⁺ , 1 ⁺)	(1 ⁺ , 0 ⁻); (1 ⁻ , 0 ⁺)	(0 ⁻ , 1 ⁻)
	0	(0⁺, 0⁺) ; (1 ⁺ , -1 ⁺)	(1 ⁺ , -1 ⁻); (-1 ⁺ , 1 ⁻); (0 ⁺ , 0 ⁻)	(1 ⁻ , -1 ⁻); (0⁻, 0⁻)
	-1	(0 ⁺ , -1 ⁺)	(0 ⁺ , -1 ⁻); (0 ⁻ , -1 ⁺)	(0 ⁻ , -1 ⁻)
	-2	(-1⁺, -1⁺)	(-1 ⁺ , -1 ⁻)	(-1⁻, -1⁻)

RS 9.9
M 9.10

Allowed Microstates

		M_S		
		1	0	-1
M_L	2		$(1^+, 1^-)$	
	1	$(0^+, 1^+)$	$(1^+, 0^-); (1^-, 0^+)$	$(0^-, 1^-)$
	0	$(1^+, -1^+)$	$(1^+, -1^-); (-1^+, 1^-); (0^+, 0^-)$	$(1^-, -1^-)$
	-1	$(0^+, -1^+)$	$(0^+, -1^-); (0^-, -1^+)$	$(0^-, -1^-)$
	-2		$(-1^+, -1^-)$	

Total number of microstates = $6!/(2!4!) = 15$

RS 9.9
M 9.10

Deducing L and S

		M_S		
		1	0	-1
M_L	2			
	1		 	
	0		  	
	-1		 	
	-2			


- $M_L = 0, \pm 1, \pm 2$ and $M_S = 0 \rightarrow L = 2$ and $S = 0$ (1D)
- $M_L = 0, \pm 1$ and $M_S = 0, \pm 1 \rightarrow L = 1$ and $S = 1$ (3P)
- $M_L = 0$ and $M_S = 0 \rightarrow L = 0$ and $S = 0$ (1S)

RS 9.9
M 9.10

Configurations and Term Symbols

Configuration	Term symbol
<i>Nonequivalent electrons</i>	
s^1s^1	$^1S, ^3S$
s^1p^1	$^1P, ^3P$
s^1d^1	$^1D, ^3D$
p^1p^1	$^1S, ^3S, ^1P, ^3P, ^1D, ^3D$
<i>Equivalent electrons</i>	
$s^2; p^6; d^{10}$	1S
s^1	2S
$p^1; p^5$	2P
$p^2; p^4$	$^1S, ^1D, ^3P$
p^3	$^2P, ^2D, ^4S$
$d^1; d^9$	2D
$d^2; d^8$	$^1S, ^1D, ^1G, ^3P, ^3F$
$d^3; d^7$	$^2P, ^2D (\times 2), ^2F, ^2G, ^2H, ^4P, ^4F$
$d^4; d^6$	$^1S (\times 2), ^1D (\times 2), ^1F, ^1G (\times 2), ^1I, ^3P (\times 2), ^3D, ^3F (\times 2), ^3G, ^3H, ^5D$
d^5	$^2S, ^2P, ^2D (\times 3), ^2F (\times 2), ^2G (\times 2), ^2H, ^2I, ^4P, ^4D, ^4F, ^4G, ^6S$

Term Symbols and Energies



Energy levels of C atom

Hund's Rules

Empirical rules to find the ground-state term symbol

1. The state with the **largest S** is the most stable
2. For states with the same S , the state with the **largest L** is the most stable
3. For states with the same S and L ,
 1. If a subshell is **less than half-filled**, the state with the **smallest J** is the most stable
 2. If a subshell is **more than half-filled**, the state with the **largest J** is the most stable



Examples

- Beryllium $1s^2 2s^1 3s^1$ (excited state, but ...)

For $s^1 s^1$, we have 3S_1 and 1S_0 as possible term symbols.

Rule 1 $\rightarrow$ 3S_1 is the most stable state.


- Carbon $1s^2 2s^2 2p^2$

For p^2 , we have 1S_0 , 3P_0 , 3P_1 , 3P_2 , 1D_2 .

Rule 1 $\rightarrow$ one of 3P states is the most stable state.

Rule 3 $\rightarrow$ 3P_0 is the most stable state.

Application of Hund's Rules



(Image: <https://texample.net/tikz/examples/energy-levels/>)

Selection Rules


(for multi-electron atoms)

$$\Delta L = 0, \pm 1 \quad (L = 0 \nrightarrow L = 0)$$

$$\Delta S = 0$$

$$\Delta J = 0, \pm 1 \quad (J = 0 \nrightarrow J = 0)$$

Allowed Transitions of Helium




(Image: <https://www1.udel.edu/pchem/C444/spLectures/05062008.pdf>)

Relativistic Effects

- Origin of the fine structure of H atom
- The spin-orbit coupling is also often called as relativistic effect
- Both effects become more significant as Z increases
- Larger spin-orbit coupling: j - j coupling
- Breakdown of the selection rules

Not in text

Orbital Energies of Inert Gas




(Image: Jerabek et al., *Phys Rev Lett*, 2018)

Not in text

Orbital Energies of Au and Hg

Without relativistic effect

With relativistic effect



- Au: $6s^1 4f^{14} 5d^{10}$
- Hg: $6s^2 4f^{14} 5d^{10}$



Not in text

Mercury Spectrum

$$\Delta L = 0, \pm 1 \quad (L = 0 \rightarrow L = 0)$$

$$\Delta S = 0$$


$$\Delta J = 0, \pm 1 \quad (J = 0 \rightarrow J = 0)$$



(Image: <https://zeiss-campus.magnet.fsu.edu/articles/lightsources/mercuryarc.html>
Liu et al., *Chinese Phys B*, 2013)

RS 9.9
M 9.13

Energy Splitting of Helium



(Image: McQuarrie, *Quantum Chemistry*, 2nd Ed., Fig. 9.11)