

# 화학수학

두 번째 수업

기초 미적분학 복습

# Today's Class

- Operations: powers and roots
- Numbers
  - Complex numbers
- Functions
  - Logarithms and exponentials
  - Trigonometric functions
- Calculus
  - Differentiation
  - Integration



# Numerical Mathematics

- Powers (거듭제곱)

$$x^2 = x \times x, \quad x^3 = x \times x \times x,$$

$$x^n = x \times x \times \cdots \times x$$

- Roots (제곱근)

$$(\sqrt{x})^2 = x, \quad (\sqrt[3]{x})^3 = x,$$

$$(\sqrt[n]{x})^n = x$$

## Exponents (지수)

$n$  in  $x^n$  = exponent of  $x$

- Negative exponents

$$x^{-1} = \frac{1}{x}, \quad x^{-3} = \frac{1}{x^3}$$

- Fractional exponents

$$x^{1/2} = \sqrt{x}, \quad x^{1/3} = \sqrt[3]{x}$$

# Properties of Exponents


- Property 1

$$x^a x^b = x^{a+b}$$

- Property 2

$$(x^a)^b = x^{ab}$$

# Numbers



(Image: <https://www.mathsisfun.com/sets/number-types.html>)

## Complex Numbers (복소수)

- Imaginary unit (허수 단위)

$$i = \sqrt{-1}$$

- Complex number ( $a, b$ : real)

$$c = a + ib$$

where  $a = \operatorname{Re}(c)$  is the **real part** of  $c$  and  $b = \operatorname{Im}(c)$  is the **imaginary part** of  $c$ .

# Operations with Complex Numbers

If  $c_1 = a_1 + ib_1$  and  $c_2 = a_2 + ib_2$ ,

- $c_1 + c_2 = (a_1 + a_2) + i(b_1 + b_2)$
- $c_1 c_2 = (a_1 + ib_1)(a_2 + ib_2)$   
 $= a_1 a_2 + i(a_1 b_2 + a_2 b_1) + i^2 b_1 b_2$   
 $= (a_1 a_2 - b_1 b_2) + i(a_1 b_2 + a_2 b_1)$



## Related Quantities

If a complex number is given as  $c = a + ib$ , its ...

- Complex conjugate (켈레복소수)

$$c^* = \bar{c} = (a + ib)^* = a - ib$$


- Magnitude (크기)

$$|c| = \sqrt{cc^*} = \sqrt{a^2 + b^2}$$

- Reciprocal (역수)

$$c^{-1} = \frac{a - ib}{a^2 + b^2} = \frac{c^*}{|c|^2}$$

# Function (함수)



Example:  $f(x) = x^2 + 2x$


$$f(1) = 1^2 + 2 \times 1 = 3$$

$$f(-1) = (-1)^2 + 2 \times (-1) = -1$$

$\vdots$

# Linear Functions (선형함수)

$$y = mx + b$$



(Image: Mortimer, *Mathematics for Physical Chemistry*, 4<sup>th</sup> ed., Figure 2.2)

# Polynomial Functions (다항함수)

$$y = a_0 + a_1x + a_2x^2 + a_3x^3 + \dots$$

- Quadratic functions

$$y = ax^2 + bx + c$$

- Cubic functions

$$y = ax^3 + bx^2 + cx + d$$

## Logarithms (로그함수)

$$x = a^y \Leftrightarrow y = \log_a x$$

- Common logarithms (상용로그):  $a = 10$
- Natural logarithms (자연로그):  $a = e$ ,  $\log_e x = \ln x$

$$e = \lim_{n \rightarrow \infty} \left( 1 + \frac{1}{n} \right)^n = 2.718 \dots$$

# Properties of Exponents and Logarithms

| Exponent fact                           | Logarithm fact                                                        |
|-----------------------------------------|-----------------------------------------------------------------------|
| $a^0 = 1$                               | $\log_a (1) = 0$                                                      |
| $a^{1/2} = \sqrt{a}$                    | $\log_a (\sqrt{a}) = \frac{1}{2}$                                     |
| $a^1 = a$                               | $\log_a (a) = 1$                                                      |
| $a^{x_1} a^{x_2} = a^{x_1+x_2}$         | $\log_a (y_1 y_2) = \log_a (y_1) + \log_a (y_2)$                      |
| $a^{-x} = \frac{1}{a^x}$                | $\log_a \left( \frac{1}{y} \right) = -\log_a (y)$                     |
| $\frac{a^{x_1}}{a^{x_2}} = a^{x_1-x_2}$ | $\log_a \left( \frac{y_1}{y_2} \right) = \log_a (y_1) - \log_a (y_2)$ |
| $(a^x)^z = a^{xz}$                      | $\log_a (y^z) = z \log_a (y)$                                         |
| $a^\infty = \infty$                     | $\log_a (\infty) = \infty$                                            |
| $a^{-\infty} = 0$                       | $\log_a (0) = -\infty$                                                |

(Image: Mortimer, *Mathematics for Physical Chemistry*, 4<sup>th</sup> ed., Table 2.2)

## Exponentials (지수함수)

$$y = e^{bx} = \exp(bx)$$

- Conversion of  $y = a^{bx}$



Assume that  $a = e^n$

This gives  $n = \log_e a = \ln a$

Substitution leads to


$$y = (e^n)^{bx} = e^{nbx} = e^{bx \ln a}$$

# Logarithms and Exponentials





# Trigonometric Functions (삼각함수)



$$\sin \alpha = \frac{y}{r},$$

$$\cos \alpha = \frac{x}{r},$$

$$\tan \alpha = \frac{y}{x},$$

$$\csc \alpha = \frac{1}{\sin \alpha} = \frac{r}{y}$$

$$\sec \alpha = \frac{1}{\cos \alpha} = \frac{r}{x}$$

$$\cot \alpha = \frac{1}{\tan \alpha} = \frac{x}{y}$$

# Trigonometric Identities (삼각함수 항등식)

$$\sin \alpha = -\sin(-\alpha)$$

$$\cos \alpha = \cos(-\alpha)$$

$$\tan \alpha = -\tan(-\alpha)$$

$$\sin \alpha = \sin(\alpha + 2\pi) = \sin(\alpha + 4\pi) = \dots$$

$$\cos \alpha = \cos(\alpha + 2\pi) = \cos(\alpha + 4\pi) = \dots$$

$$\sin^2 \alpha + \cos^2 \alpha = 1$$

# Trigonometric Functions




FIGURE 2.7 The sine of an angle.




FIGURE 2.8 The cosine of an angle.




FIGURE 2.9 The tangent of an angle.

(Image: Mortimer, *Mathematics for Physical Chemistry*, 4<sup>th</sup> ed., Figures 2.7, 2.8, and 2.9)

# Inverse Trigonometric Functions

## (역 삼각함수)

$$y = \sin x \Leftrightarrow x = \arcsin y = \operatorname{asin} y = \sin^{-1} y$$

$$y = \cos x \Leftrightarrow x = \arccos y = \operatorname{acos} y = \cos^{-1} y$$

$$y = \tan x \Leftrightarrow x = \arctan y = \operatorname{atan} y = \tan^{-1} y$$

# Euler's Formula

$$e^{ix} = \cos x + i \sin x$$

- Application of Euler's formula

$$e^{ix} = \cos x + i \sin x$$

$$e^{-ix} = \cos x - i \sin x$$

Addition and subtraction leads to

$$\cos x = \frac{e^{ix} + e^{-ix}}{2}, \quad \sin x = \frac{e^{ix} - e^{-ix}}{2i}$$

# Hyperbolic Trigonometric Functions

(쌍곡 삼각함수)


$$\sinh x = \frac{e^x - e^{-x}}{2}, \quad \cosh x = \frac{e^x + e^{-x}}{2}$$

$$\tanh x = \frac{\sinh x}{\cosh x}, \quad \coth x = \frac{1}{\tanh x}$$

$$\operatorname{sech} x = \frac{1}{\cosh x}, \quad \operatorname{csch} x = \frac{1}{\sinh x}$$

# Differentiation


“The slope of a function”



$$\lim_{\Delta x \rightarrow 0} \frac{\Delta y}{\Delta x} = \frac{dy}{dx}$$

(Image: [http://wwwf.imperial.ac.uk/metric/metric\\_public/differentiation/index.html](http://wwwf.imperial.ac.uk/metric/metric_public/differentiation/index.html))


# First Derivative: Slope



(Image: Atkins *et al.*, *Atkins' Physical Chemistry*, 11<sup>th</sup> ed., the chemist's toolkit 5, Sketch 1)



# Second Derivative: Sharpness



# Some Examples of Derivatives

| Function, $y = y(x)$ | Derivative, $dy/dx = y'(x)$ |
|----------------------|-----------------------------|
| $ax^n$               | $anx^{n-1}$                 |
| $ae^{bx}$            | $abe^{bx}$                  |
| $a$                  | $0$                         |
| $a \sin(bx)$         | $ab \cos(bx)$               |
| $a \cos(bx)$         | $-ab \sin(bx)$              |
| $a \ln(x)$           | $a/x$                       |
| $a \tan(x)$          |                             |

Check Appendix D!



## Derivative Identities (1)

- If  $y$  is a function of  $x$  and  $c$  is a constant,

$$\frac{d(y + c)}{dx} = \frac{dy}{dx}, \quad \frac{d(cy)}{dx} = c \frac{dy}{dx}$$

- If  $y$  and  $z$  are both functions of  $x$ ,

$$\frac{d(y + z)}{dx} = \frac{dy}{dx} + \frac{dz}{dx}$$
$$\frac{d(yz)}{dx} = z \frac{dy}{dx} + y \frac{dz}{dx}$$

## Derivative Identities (2)

- If  $y$  and  $z$  are both functions of  $x$ ,

$$\frac{d}{dx} \left( \frac{y}{z} \right) = \frac{1}{z^2} \left( z \frac{dy}{dx} - y \frac{dz}{dx} \right)$$

- If  $u$  is a function of  $x$  and  $f$  is a function of  $u$ ,

$$\frac{df}{dx} = \frac{df}{du} \frac{du}{dx}$$

(Chain rule)

## Example 6.6

Find  $dP/dT$  if  $P(T) = ke^{-Q/T}$ .

Let  $u = -Q/T$ . Then,  $P(T) = ke^u$ . From the chain rule,


$$\frac{dP}{dT} = \frac{dP}{du} \frac{du}{dT} = \frac{d(ke^u)}{du} \times \frac{d}{dT} \left( -\frac{Q}{T} \right) = ke^u \times \frac{Q}{T^2}$$

Therefore,

$$\frac{dP}{dT} = \frac{kQ}{T^2} e^{-Q/T}$$

# Integration

“The areas under curves”



$$\lim_{\delta x \rightarrow 0} \sum_i f(x_i) \delta x = \int_a^b f(x) dx$$

## Indefinite Integrals (부정적분)

Consider an integral

$$\int_a^{x'} f(x) dx = F(x') + C$$

constant of  
integration  
(적분상수)

where  $x'$  is a variable and  $a$  is an arbitrary constant.

Generally, it is written as

$$\int f(x) dx = F(x) + C$$

# Calculation of Indefinite Integrals



When evaluating the integral  $F(x) = \int f(x)dx$ ,  
find a function  $F(x)$  whose derivative is  $f(x)$ :

$$f(x) = \frac{dF(x)}{dx}$$



# Some Examples of Derivatives

Integration



| Function, $y = y(x)$ | Derivative, $dy/dx = y'(x)$ |
|----------------------|-----------------------------|
| $ax^n$               | $anx^{n-1}$                 |
| $ae^{bx}$            | $abe^{bx}$                  |
| $a$                  | $0$                         |
| $a \sin(bx)$         | $ab \cos(bx)$               |
| $a \cos(bx)$         | $-ab \sin(bx)$              |
| $a \ln(x)$           | $a/x$                       |
| $a \tan(x)$          |                             |

Check Appendix E!

# Calculation of Definite Integrals

If the indefinite integral of  $f(x)$  is  $F(x)$ ,

$$\int_a^b f(x) \, dx = F(b) - F(a)$$

No constant of integration!

# Techniques of Integration

Method of substitution (변수치환법)

Integration by parts (부분적분법)

Method of partial fractions (부분분수법)

# Method of Substitution

1. Define a new variable
2. Calculate its infinitesimal change and the limits
3. Rewrite the integral in terms of the new variable

## Example 7.9

Evaluate the integral

$$\int_0^{\infty} x e^{-x^2} dx.$$

Let  $u = x^2$ . Then,

$$du = 2x dx$$

$$x = 0 \rightarrow u = 0$$

$$x = \infty \rightarrow u = \infty$$

Hence, substitution leads to

$$\int_0^{\infty} \frac{1}{2} e^{-u} du = \left[ -\frac{1}{2} e^{-u} \right]_0^{\infty} = \frac{1}{2}$$

# Integration by Parts

If  $u$  and  $v$  are both functions of  $x$ ,

$$\frac{d(uv)}{dx} = v \frac{du}{dx} + u \frac{dv}{dx}$$

Integrating both sides,

$$uv = \int v \frac{du}{dx} dx + \int u \frac{dv}{dx} dx + C$$

which gives

$$\int u \frac{dv}{dx} dx = uv - \int v \frac{du}{dx} dx + C$$

# Integration by Parts

For indefinite integrals,

$$\int u \frac{dv}{dx} dx = uv - \int v \frac{du}{dx} dx + C$$

For definite integrals,

$$\int_a^b u \frac{dv}{dx} dx = uv \Big|_a^b - \int_a^b v \frac{du}{dx} dx$$

## Example 7.11

Find the definite integral

$$\int_0^{\pi} x \sin x \, dx.$$

Let  $u = x$  and  $dv/dx = \sin x$ . Then,  $v = -\cos x$ .

$$\int_0^{\pi} x \sin x \, dx = \int_0^{\pi} u \frac{dv}{dx} dx = uv \Big|_0^{\pi} - \int_0^{\pi} v \frac{du}{dx} dx$$

Hence, substitution leads to

$$x(-\cos x) \Big|_0^{\pi} - \int_0^{\pi} (-\cos x) \, dx = \pi + (\sin x) \Big|_0^{\pi} = \pi$$



## Method of Partial Fractions

If  $P(x)$  and  $Q(x)$  are polynomials, and  $Q(x)$  can be factorized as

$$Q(x) = (a_1x + b_1)(a_2x + b_2) \cdots (a_nx + b_n),$$

$P(x)/Q(x)$  can be decomposed as

$$\frac{P(x)}{Q(x)} = \frac{A_1}{a_1x + b_1} + \frac{A_2}{a_2x + b_2} + \cdots + \frac{A_n}{a_nx + b_n}$$

Simpler combination of fractions

## Example 7.12

Decompose the integral

$$\int \frac{6x - 30}{x^2 + 3x + 2} dx.$$

The denominator is decomposed as  $x^2 + 3x + 2 = (x + 2)(x + 1)$ . Thus,

$$\frac{6x - 30}{x^2 + 3x + 2} = \frac{A_1}{x + 2} + \frac{A_2}{x + 1}$$

Multiplying both sides by  $(x + 2)(x + 1)$ ,

$$6x - 30 = A_1(x + 1) + A_2(x + 2)$$

$$6x - 30 = (A_1 + A_2)x + (A_1 + 2A_2)$$

Since  $A_1 + A_2 = 6$  and  $A_1 + 2A_2 = -30$ ,  $A_1 = 42$  and  $A_2 = -36$ . Hence,

$$\int \frac{6x - 30}{x^2 + 3x + 2} dx = \int \frac{42}{x + 2} dx - \int \frac{36}{x + 1} dx.$$

