
Polynomial Ham Sandwich Theorem and Its Applications in Combinatorics

~ * ~

— Proof of the Szemerédi–Trotter Theorem —

Park Jeong-woo^{*}

Student ID : 20190262

Department of Mathematical Sciences,
Korea Advanced Institute of Science and Technology

^{*}My family name is *Park*, but please call me *Jeong-woo*.

email jeongwoo-math@kaist.ac.kr

site jeongwoo-math.github.io

Abstract

One of the most basic and powerful tool in combinatorics is *counting case by case*. This article includes a polynomial method for partitioning sets, which can be used to apply case-by-case-counting method. Also, I'll introduce its applications, like the Szemerédi–Trotter theorem.

Contents

1	Brief Introduction	3
2	The Ham Sandwich theorems	3
3	Cell Decomposition	4
4	Szemerédi–Trotter Theorem	4

1 Brief Introduction

The [ham sandwich theorem](#) states that, for any d bounded open subsets of a Euclidean space \mathbb{R}^d , there is a *hyperplane* bisecting each of them, in the sense of the *Lebesgue measure*. One can consider *hypersurfaces* instead of hyperplanes. In this case, we can bisect *more than d* bounded open subsets of \mathbb{R}^d . Also, there are discrete analogues of them, by replacing the Lebesgue measure to the *counting measure*, and bounded open subsets to finite subsets. We can use this to make a *nice division* of a finite set, which can be applied to prove Szemerédi–Trotter theorem.

2 The Ham Sandwich theorems

Theorem 2.1 (Borsuk–Ulam theorem).

The classical ham sandwich theorem can be stated as follows.

Definition 2.2 (bisection). Let A be a Lebesgue-measurable subset of \mathbb{R}^d . We call a hyperplane $H = \{x \in \mathbb{R}^d; x \cdot v = b\}$ *bisects* A , if and only if, $m(A \cap H^+) = m(A \cap H^-) = \frac{1}{2} \cdot m(A)$, where $H^+ := \{x \in \mathbb{R}^d; x \cdot v > b\}$ and $H^- := \{x \in \mathbb{R}^d; x \cdot v < b\}$ are two half-spaces.¹

Theorem 2.3 (classical ham sandwich theorem). Let $(O_i)_{1 \leq i \leq d}$ be a collection of bounded open subsets of \mathbb{R}^d . There is a hyperplane H bisecting each of $(O_i)_{1 \leq i \leq d}$.

Proof. Use Borsuk–Ulam theorem. Standard proofs can be found in (???) □

Theorem 2.4 (polynomial ham sandwich theorem).

Theorem 2.5 (general ham sandwich theorem, cf. [ST42]).

Theorem 2.6 (discretized classical sandwich theorem).

Theorem 2.7 (discretized polynomial sandwich theorem).

Definition 2.8 (Veronese embedding).

¹The well-definedness is not too hard to show, because H^+ and H^- can be considered as the two connected components of $\mathbb{R}^d \setminus H$. However, I prefer the former definition, because it is more fundamental and easy to generalize.

3 Cell Decomposition

Lemma 3.1 (cell decomposition).

Theorem 3.2 ([Tho15]).

Lemma 3.3 (Bézout’s theorem, cf. [Ful08]).

4 Szemerédi–Trotter Theorem

Theorem 4.1 (Szemerédi–Trotter theorem).

[KMS11] [Mun00] [Tho15] [ST42] [Ta01] [Yu15]

References

- [ST42] A. H. Stone and J. W. Tukey. “Generalized “sandwich” theorems”. In: *Duke Mathematical Journal* 9.2 (1942), pp. 356–359. DOI: [10.1215/S0012-7094-42-00925-6](https://doi.org/10.1215/S0012-7094-42-00925-6). URL: <https://doi.org/10.1215/S0012-7094-42-00925-6>.
- [Mun00] James Munkres. *Topology*. 2nd ed. Pearson, 2000. ISBN: 978-0131816299.
- [Ful08] William Fulton. *Algebraic Curves — An Introduction to Algebraic Geometry*. 2008.
- [KMS11] Haim Kaplan, Jiří Matoušek, and Micha Sharir. *Simple Proofs of Classical Theorems in Discrete Geometry via the Guth–Katz Polynomial Partitioning Technique*. 2011. arXiv: [1102.5391](https://arxiv.org/abs/1102.5391) [math.CO].
- [Tao11] Terence Tao. *The Szemerédi–Trotter theorem via the polynomial ham sandwich theorem*. 2011. URL: <https://terrytao.wordpress.com/2011/02/18/the-szemerédi-trotter-theorem-via-the-polynomial-ham-sandwich-theorem>. accessed: 05.28.2021.
- [Tho15] René Thom. “Sur L’Homologie des Variétés Algébriques Réelles”. In: *Differential and Combinatorial Topology*. Princeton University Press, 2015, pp. 255–265. DOI: [doi: 10.1515/9781400874842-016](https://doi.org/10.1515/9781400874842-016). URL: <https://doi.org/10.1515/9781400874842-016>.
- [Wig15] Yuval Wigderson. *The Ham Sandwich Theorem and Friends*. 2015. URL: <http://web.stanford.edu/~yuvalwig/math/teaching>. accessed: 05.28.2021.