

Picard's Existence and Uniqueness Theorem

Author: Jeremiah Givens

Professor: Dr. Claudio Morales

Department of Mathematics, University of Alabama in Huntsville

1 Introduction

The main objective of this paper is to examine a very practical application of Banach's fixed point theorem: Picard's Existence and Uniqueness Theorem for Ordinary Differential Equations. We will begin by proving some useful lemmas, which we will then use to prove Picard's theorem. After proving the theorem, we will then look at a few example differential equations that Picard's theorem proves the existence of solutions to, and we will use an iterative technique, known as the Picard Iteration, to approximate the solutions.

2 Preliminaries

Lemma 1. *Let $x : [a, b] \rightarrow \mathbb{R}$ be continuous. Then, x is bounded, and x achieves its maximum.*

Proof. We have

$$\begin{aligned} [a, b] \text{ is compact} &\implies f([a, b]) \text{ is compact} && \text{continuous image of compact set} \\ &\implies f([a, b]) \text{ is closed and bounded} && \text{by the Heine-Borel Theorem, since } f([a, b]) \subseteq \mathbb{R}^2 \\ &\implies f \text{ is bounded.} \end{aligned}$$

Now we need to show that x achieves its maximum. We know from our argument above, that x has an upper bound. Thus, the least upper bound property of the real numbers tells us that the supremum $M = \sup\{x(t) : t \in [a, b]\}$ exists. By the definition of a supremum, we can choose a sequence $\{t_n\}_{n \in \mathbb{N}}$ such that $M - \frac{1}{n} \leq x(t_n) \leq M$. Clearly, the sequence $\{x(t_n)\}_{n \in \mathbb{N}}$ converges to M . Since $[a, b]$ is compact, and therefore sequentially compact, we have that $\{t_n\}_{n \in \mathbb{N}}$ has a subsequence $\{t_{n_i}\}_{i \in \mathbb{N}}$ that converges in $[a, b]$. By definition of continuity, $\{x(t_{n_i})\}_{i \in \mathbb{N}}$ converges. Since a subsequence of any convergent sequence converges to the same limit, we have that $\{x(t_{n_i})\}_{i \in \mathbb{N}}$ converges to M . Therefore, x achieves its maximum at $t = \lim_{i \rightarrow \infty} t_{n_i}$. \square

Lemma 2. *Let C equal the set of all continuous real-valued functions on the closed interval $[a, b]$. Then, C forms a metric space with the metric d defined by*

$$d(x, y) = \max_{t \in [a, b]} |x(t) - y(t)|.$$

Proof. We must first start with showing that d is well defined. Let $x, y : [a, b] \rightarrow \mathbb{R}$ be continuous functions. We know that the difference of two continuous functions is continuous, thus the function $x(t) - y(t)$ is continuous. Likewise, we have that the composition of continuous functions is continuous. Thus, since the absolute value function is continuous, we have that the function $|x(t) - y(t)|$ is continuous. Thus, by Lemma 1, $\max_{t \in [a, b]} |x(t) - y(t)|$ exists, and d is well defined.

Now we must show that d is a metric. Let $x, y \in C$. By definition absolute value, we clearly have that $d(x, y) \geq 0$. Now we have

$$\begin{aligned} d(x, y) = 0 &\iff \max_{t \in [a, b]} |x(t) - y(t)| = 0 \\ &\iff (\forall t \in [a, b])(0 \leq |x(t) - y(t)| \leq 0) && \text{definition of absolute value and max} \\ &\iff (\forall t \in [a, b])(|x(t) - y(t)| = 0) \\ &\iff (\forall t \in [a, b])(x(t) = y(t)) \\ &\iff x = y. \end{aligned}$$

Thus, d fits the first criteria of a metric.

Following easily from the definition of absolute value, we have

$$\begin{aligned} d(x, y) &= \max_{t \in [a, b]} |x(t) - y(t)| \\ &= \max_{t \in [a, b]} |y(t) - x(t)| \\ &= d(y, x), \end{aligned}$$

and d fits the second criteria of a metric.

Let $x, y, z \in C$. We have

$$\begin{aligned} d(x, z) &= \max_{t \in [a, b]} |x(t) - z(t)| \\ &= \max_{t \in [a, b]} (|x(t) - y(t)| + |y(t) - z(t)|) && \text{Since the absolute value forms a metric on } \mathbb{R}. \end{aligned}$$

We want to show that

$$\max_{t \in [a, b]} (|x(t) - y(t)| + |y(t) - z(t)|) \leq \max_{t \in [a, b]} |x(t) - y(t)| + \max_{t \in [a, b]} |y(t) - z(t)|.$$

By Lemma 1, all of these maxes are achieved. Therefore, we can define $t_1, t_2, t_3 \in [a, b]$ to be such that

$$\begin{aligned} |x(t_1) - y(t_1)| + |y(t_1) - z(t_1)| &= \max_{t \in [a, b]} (|x(t) - y(t)| + |y(t) - z(t)|) \\ |x(t_2) - y(t_2)| &= \max_{t \in [a, b]} |x(t) - y(t)| \\ |y(t_3) - z(t_3)| &= \max_{t \in [a, b]} |y(t) - z(t)|. \end{aligned}$$

In terms of t_1, t_2 , and t_3 , we want to show that

$$|x(t_1) - y(t_1)| + |y(t_1) - z(t_1)| \leq |x(t_2) - y(t_2)| + |y(t_3) - z(t_3)|.$$

We have

$$\begin{aligned} |x(t_1) - y(t_1)| + |y(t_1) - z(t_1)| &\leq |x(t_2) - y(t_2)| + |y(t_1) - z(t_1)|, \text{ since } |x(t_1) - y(t_1)| \leq |x(t_2) - y(t_2)| \text{ by def. of } t_2 \\ &\leq |x(t_2) - y(t_2)| + |y(t_3) - z(t_3)|, \text{ since } |y(t_1) - z(t_1)| \leq |y(t_3) - z(t_3)| \text{ by def. of } t_3, \end{aligned}$$

and we have shown that d satisfies the triangle inequality. With this, we have finally shown that d is a metric on C . \square

Lemma 3. *The metric space (C, d) from the previous lemma is complete.*

Proof. Let $\{x_n\}_{n \in \mathbb{N}}$ be a Cauchy sequence in C . Define $f = \lim_{n \rightarrow \infty} x_n$. We wish to show that $f \in C$. Let $\epsilon > 0$, and let $p_0 \in [a, b]$. Since $\{x_n\}_{n \in \mathbb{N}}$ converges to f , there exists an $n \in \mathbb{N}$ such that $d(x_n, f) < \epsilon/3$. By definition of our metric, this implies $|x_n(p) - f(p)| < \epsilon/3$ for all $p \in [a, b]$. Now, since x_n is continuous,

$$(\exists \delta > 0)(\forall p \in [a, b])(|p - p_0| < \delta \implies |x_n(p) - x_n(p_0)| < \epsilon/3).$$

Then, for all $p \in [a, b]$ with $|p - p_0| < \delta$, we have

$$\begin{aligned} |f(p) - f(p_0)| &\leq |f(p) - x_n(p)| + |f(p_0) - x_n(p_0)| && \text{by triangle inequality} \\ &\leq |f(p) - x_n(p)| + |f(p_0) - x_n(p_0)| + |x_n(p) - x_n(p_0)| && \text{again, by triangle inequality} \\ &\leq \epsilon/3 + \epsilon/3 + \epsilon/3 \\ &= \epsilon. \end{aligned}$$

Thus, f is continuous at p_0 . Since p_0 was arbitrary, f is continuous, and we can conclude that $f \in C$. Thus, C is complete. \square

Lemma 4. *Let $x_0 \in \mathbb{R}$, and let $c \in (0, \infty)$. Let \tilde{C} be a subspace of the metric space from Lemma 1, consisting of all functions $x \in C$ such that*

$$d(x, x_0) \leq c.$$

Then, \tilde{C} is closed.

Proof. Let $f \in \tilde{C}$. We have

$$\begin{aligned} f \in \tilde{C} &\iff (\forall \epsilon > 0)(B(f; \epsilon) \cap \tilde{C} \neq \emptyset) \\ &\iff (\forall \epsilon > 0)(\exists x \in \tilde{C})(d(f, x) \leq \epsilon) \\ &\iff (\forall n \in \mathbb{N})(\exists x_n \in \tilde{C})(d(f, x_n) \leq \frac{1}{n}). \end{aligned}$$

With this, we have

$$\begin{aligned} d(f, x_0) &\leq d(f, x_n) + d(x_n, x_0) && \text{triangle inequality} \\ &\leq \frac{1}{n} + d(x_n, x_0) \\ &\leq \frac{1}{n} + c && \text{since } x_n \in \tilde{C} \end{aligned}$$

Since this is true for all $n \in \mathbb{N}$, we have $d(f, x_0) \leq c$ which implies $f \in \tilde{C}$. \square

Lemma 5. *Let C be a complete metric space. Let $A \subseteq C$ be a closed. Then, A is complete.*

Proof. Let A be closed, and let $\{x_n\}_{n \in \mathbb{N}}$ be a Cauchy sequence in A . Since X is complete, we know that the limit converges to some point x in X . We have

$$\begin{aligned}
 \lim_{n \rightarrow \infty} x_n &\implies (\forall \epsilon > 0)(\exists N \in \mathbb{N})(n \geq N \implies d(x_n, x) \leq \epsilon) \\
 &\implies (\forall \epsilon > 0)(\exists x_n \in A)(x_n \in B(x; \epsilon)) \\
 &\implies (\forall \epsilon > 0)(B(x; \epsilon) \cap A \neq \emptyset) \\
 &\implies x \in \bar{A} \\
 &\implies x \in A.
 \end{aligned}$$

since A is closed

Since this is true for any Cauchy sequence in A , A is complete. \square

3 Main Topic

The focus of this paper is on solutions to explicit ordinary differential equations of the form

$$x' = f(t, x) \tag{1}$$

where $x : \mathbb{R} \rightarrow \mathbb{R}$, $f : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$, and the prime denotes differentiation of x with respect to t . Specifically, we are interested in initial value problems, where

$$x(t_0) = x_0.$$

Theorem 1. *Let f be continuous on a rectangle*

$$R = \{(t, x) \in \mathbb{R}^2 : |t - t_0| \leq a \wedge |x - x_0| \leq b\},$$

and thus bounded on R , say

$$|f(t, x)| \leq c, \text{ for all } (t, x) \in R. \tag{2}$$

Suppose f satisfies a Lipschitz condition on R with respect to its second argument, that is, there is a constant k (Lipschitz constant) such that for $(t, x), (t, v) \in R$

$$|f(t, x) - f(t, v)| \leq k|x - v|. \tag{3}$$

Then the initial value problem (1) has a unique solution. This solution exists on an interval $[t_0 - \beta, t_0 + \beta]$ where

$$\beta < \min \left\{ a, \frac{b}{c}, \frac{1}{k} \right\} \tag{4}$$

Proof. Let $C(J)$ be the metric space of all real-valued continuous functions on the interval $J = [t_0 - \beta, t_0 + \beta]$ with metric d defined by

$$d(x, y) = \max_{t \in J} |x(t) - y(t)|.$$

By Lemma 2 we know that $C(J)$ is a metric space, and by Lemma 3 we know that $C(J)$ is complete. Let \tilde{C} be a subspace of $C(J)$ consisting of all functions $x \in C(J)$ such that

$$|x(t) - x_0| \leq c\beta. \tag{5}$$

By Lemma 4, we have that \tilde{C} is closed, so by Lemma 5 \tilde{C} is complete.

From the fundamental theorem of calculus, (1) can be written in the form $x = Tx$, where $T : \tilde{C} \rightarrow \tilde{C}$ is defined by

$$Tx(t) = x_0 + \int_{t_0}^t f(\tau, x(\tau)) d\tau \tag{6}$$

We have $\tau \in J \implies |\tau - t_0| \leq a$ (by 4), and

$$\begin{aligned} x \in \tilde{C} &\implies |x(\tau) - x_0| \leq c\beta && \text{by (5)} \\ &\implies |x(\tau) - x_0| \leq b && \text{by (4).} \end{aligned}$$

Thus, $(\tau, x(\tau)) \in R$ and the integral in (6) exists since f is continuous on R . To see that T maps \tilde{C} into itself,

$$\begin{aligned} |Tx(t) - x_0| &= \left| \int_{t_0}^t f(\tau, x(\tau)) d\tau \right| \\ &\leq c|t - t_0| && \text{by (2)} \\ &\leq c\beta && \text{since } t \in [t_0 - \beta, t_0 + \beta]. \end{aligned}$$

Thus, by (5) we have that $Tx \in \tilde{C}$.

We will now show that T is a contraction on \tilde{C} . We have

$$\begin{aligned} |Tx(t) - Tv(t)| &= \left| \int_{t_0}^t f(\tau, x(\tau)) - f(\tau, v(\tau)) d\tau \right| \\ &\leq |t - t_0| \max_{\tau \in J} k|x(\tau) - v(\tau)| && \text{by the Lipschitz condition (3)} \\ &\leq k\beta d(x, v) && \text{by (4) and by definition of a the metric } d. \end{aligned}$$

Since the right hand side does not depend on t , we can take the max on both sides to yield

$$d(Tx, Tv) \leq k\beta d(x, v).$$

By (4), we have that $k\beta < 1$. Thus, we have shown that T is a contraction mapping on \tilde{C} .

By the Banach Fixed Point Theorem, T has a unique fixed point $x \in \tilde{C}$, such that $Tx = x$. That is, there exists a unique $x \in \tilde{C}$ such that

$$x(t) = x_0 + \int_{t_0}^t f(\tau, x(\tau)) d\tau. \quad (7)$$

Since $(\tau, x(\tau)) \in R$ where f is continuous, we have that both sides of (7) are differentiable, and x satisfies (1). Since every solution x to (1) must satisfy (7), we have that x is unique, and our proof is complete. \square

Examples

As we saw in the proof of the Banach Fixed Point Theorem, fixed points of a contraction mapping can be written as the limit of the sequence $\{T^n x_0\}_{n=0}^\infty$. This provides us with an iterative technique for approximating the solution to differential equations that are shown to have a unique solution by Picard's Theorem. In this section we will go over some examples that illustrate this technique.

Example 1

We start first, with the well known equation

$$x' = x, \text{ and } x(0) = 1.$$

In this case, $f(t, x) = x$, which is clearly continuous on \mathbb{R}^2 . Also, f satisfies our Lipschitz Condition, as

$$|f(t, x) - f(t, v)| = |x - v|,$$

and we can see that our Lipschitz constant is any $k \geq 1$. With all of this, we can see that Picard's theorem gives us a solution for t in any closed interval centered at 0.

Lets go ahead and compute the first few terms:

$$\begin{aligned} x_0 &= 1 \\ Tx_0 &= x_0 + \int_{t_0}^t x_0 d\tau = 1 + \int_0^t 1 d\tau = 1 + t \\ T^2 x_0 &= 1 + \int_0^t (1 + \tau) d\tau = 1 + t + \frac{1}{2}t^2. \end{aligned}$$

Continuing in this manner, it becomes quite obvious that Picard's iteration is computing the Taylor series of the exponential function

$$x(t) = \sum_{n=0}^{\infty} \frac{t^n}{n!} = e^t,$$

which we know to be the solution of this differential equation.

Example 2

We will now consider a more interesting equation

$$x' = t \cos(x), \text{ and } x(0) = 0.$$

Once again, we have that f is continuous \mathbb{R}^2 . We then need to show that f fits the Lipschitz condition. We have

$$\begin{aligned} |f(t, x) - f(t, v)| &= |t \cos(x) - t \cos(v)| \\ &\leq |t| |\cos(x) - \cos(v)| \\ &= |t| |\cos(x) - \cos(v)|. \end{aligned}$$

Now we know the derivative

$$\frac{d}{dt}(\cos(\theta)) = -\sin(\theta).$$

Assuming, without loss of generality, that $x \leq y$, the mean value theorem from elementary calculus tells us that $\exists c \in (x, y)$ such that

$$\begin{aligned} -\sin(c) &= \frac{\cos(v) - \cos(x)}{v - x} \implies \sin(c)(x - v) = \cos(v) - \cos(x) \\ &\implies |\cos(x) - \cos(v)| = |\sin(c)||x - v| \\ &\implies |\cos(x) - \cos(v)| \leq |x - v| \quad \text{Since } \sin(c) \in [-1, 1]. \end{aligned}$$

Therefore,

$$\begin{aligned} |f(t, x) - f(t, v)| &\leq |t| |\cos(x) - \cos(v)| \\ &\leq |t| |x - v| \\ &\leq k |x - v| \quad \text{if we restrict } t \in [-k, k] \text{ for some } k \in (0, \infty), \end{aligned}$$

and we have determined f fits the Lipschitz condition, and Picard's theorem tells us there exists a unique solution to this differential equation.

Jumping right in, let's calculate the first few terms of the Picard Iteration.

$$\begin{aligned} x_0 &= 0 \\ Tx_0 &= x_0 + \int_0^t \tau \cos(x_0) d\tau = \int_0^t \tau d\tau = \frac{1}{2}t^2 \\ T^2x_0 &= \int_0^t \tau \cos\left(\frac{1}{2}\tau^2\right) d\tau = \sin\left(\frac{1}{2}t^2\right) \\ T^3x_0 &= \int_0^t \tau \cos\left(\sin\left(\frac{1}{2}\tau^2\right)\right) d\tau = \frac{1}{2}t^2 \cos\left(\sin\left(\frac{t^2}{2}\right)\right) \\ T^4x_0 &= \int_0^t \tau \cos\left(\frac{1}{2}\tau^2 \cos\left(\sin\left(\frac{\tau^2}{2}\right)\right)\right) d\tau. \end{aligned}$$

We did not compute a closed form of that last term, as it exceeded the time of our computer algebra software. This highlights an unpleasant feature of this technique: it can quickly become difficult to compute higher terms in the sequence.

It's interesting to note that, in contrast to the last example, the Picard iteration is not computing the Taylor series of our solution. However, we have all the information we need to compute the Taylor series of x . For the

purpose of comparison, let's define a new sequence, $\{x_n\}_{n=0}^{\infty}$, where the n^{th} term is the first $n + 1$ terms of the Taylor series. Then, we have

$$\begin{aligned}x_0 &= 0 \\x_1 &= 0 \\x_2 &= \frac{t^2}{2} \\x_3 &= \frac{t^2}{2} - \frac{t^6}{48} \\x_4 &= \frac{t^2}{2} - \frac{t^6}{48} + \frac{t^{10}}{768} \\x_5 &= \frac{t^2}{2} - \frac{t^6}{48} + \frac{t^{10}}{768} - \frac{61t^{14}}{645120}\end{aligned}$$