

## Problem 1

Use Holder's inequality to show

$$\int_0^1 \sqrt{x}(1-x)^{-1/3} dx \leq \frac{2}{5^{1/3}}.$$

### Solution

Using Holder's inequality, let  $p = 3$ , and let  $p' = 3/2$ . Then, we have

$$\begin{aligned} \int_0^1 \sqrt{x}(1-x)^{-1/3} dx &\leq \left( \int_0^1 \sqrt{x}^3 dx \right)^{1/3} \left( \int_0^1 ((1-x)^{-1/3})^{3/2} dx \right)^{2/3} \\ &= \left( \int_0^1 x^{3/2} dx \right)^{1/3} \left( \int_0^1 u^{-1/2} du \right)^{2/3} \\ &= \left( \frac{2}{5} \right)^{1/3} 2^{2/3} \\ &= \frac{2}{5^{1/3}}, \end{aligned}$$

as desired.

## Problem 2

Let  $E \subseteq \mathbb{R}^n$  be measurable with  $|E| = 1$ . Let  $h \geq 0$  be measurable on  $E$ . Let  $A = \int_E h dx$ . Show that

$$\sqrt{1 + A^2} \leq \int_E \sqrt{1 + h^2} dx \leq 1 + A.$$

### Solution

(I've not yet figured out the proof of the first inequality, which would go here.)

For the second inequality, we have

$$\begin{aligned} \int_E \sqrt{1 + h^2} dx &\leq \int_E \sqrt{1 + h^2 + 2h} dx \\ &= \int_E \sqrt{(1 + h)^2} dx \\ &= \int_E 1 + h dx \\ &= 1 + A. \end{aligned}$$

## Problem 3

Find all nonnegative functions  $g \in L^3(0, 1)$  such that

$$\left( \int_0^1 xg(x) dx \right)^3 = \frac{4}{25} \int_0^1 g^3(x) dx$$

**Solution**

Using Holders inequality, we can see that if  $p = \frac{3}{2}$ , and  $p' = 3$ , then

$$\begin{aligned} \int_0^1 xg(x)dx &\leq \left( \int_0^1 x^{3/2}dx \right)^{2/3} \left( \int_0^1 g^3(x)dx \right)^{1/3} \\ &= \left( \frac{2}{5} \right)^{2/3} \left( \int_0^1 g^3(x)dx \right)^{1/3} \\ &= \left( \frac{4}{25} \int_0^1 g^3(x)dx \right)^{1/3} \\ \left( \int_0^1 xg(x)dx \right)^3 &= \frac{4}{25} \int_0^1 g^3(x)dx. \end{aligned}$$

Now, as we proved in class, the equality holds if and only if  $\alpha x^{3/2} = g^3(x)$  almost everywhere for some real  $\alpha$ . Thus, we have

$$g(x) = \alpha x^{\frac{1}{2}}$$

for some nonnegative real number  $\alpha$  and for almost every  $x$ .

**Problem 4**

Let  $f \in L^\infty(0,1)$  and  $\|f\|_\infty \leq 1$ . Show that

$$\int_0^1 \sqrt{1-f^2(x)}dx \leq \sqrt{1 - \left( \int_0^1 f(x)dx \right)^2},$$

and describe the class of functions  $f$  for which equality takes place.

**Solution**

We have

$$\begin{aligned} \int_0^1 \sqrt{1-f^2(x)}dx &= \int_0^1 \sqrt{(1-f)(1+f)}dx \\ &= \int_0^1 \sqrt{1-f} \sqrt{1+f} dx \\ &\leq \sqrt{\int_0^1 (1-f)dx} \sqrt{\int_0^1 (1+f)dx} && \text{Cauchy-Schwarz's inequality} \\ &= \sqrt{1 - \int_0^1 f dx} \sqrt{1 + \int_0^1 f dx} \\ &= \sqrt{1 - \left( \int_0^1 f(x)dx \right)^2}. \end{aligned}$$

From the proof of Holder's Inequality, we know that the equality holds iff there exists some  $\alpha \in \mathbb{R}$  such that  $1-f = \alpha(1+f)$ . Thus,

$$\begin{aligned} 1-f &= \alpha(1+f) \implies 1-f = \alpha + \alpha f \\ &\implies 1-\alpha = (1+\alpha)f \\ &\implies f = \frac{1-\alpha}{1+\alpha}. \end{aligned}$$

## Problem 5

**Theorem 1.** *Prove that*

$$\int_0^\infty e^{-x} \sqrt{x^4 + 3x^2 + 2} dx \leq \sqrt{12},$$

*and that the equality does not hold.*

### Solution

*Proof.* We have

$$\begin{aligned} \int_0^\infty e^{-x} \sqrt{x^4 + 3x^2 + 2} dx &= \int_0^\infty \sqrt{e^{-2x}(x^4 + 3x^2 + 2)} dx \\ &= \int_0^\infty \sqrt{e^{-x}(x^2 + 2)e^{-x}(x^2 + 1)} dx \\ &= \int_0^\infty \sqrt{e^{-x}(x^2 + 2)} \sqrt{e^{-x}(x^2 + 1)} dx \\ &\leq \sqrt{\int_0^\infty e^{-x}(x^2 + 2) dx} \sqrt{\int_0^\infty e^{-x}(x^2 + 1) dx} \quad \text{Cauchy-Schwarz's inequality} \\ &= \sqrt{2 + 2\sqrt{2 + 1}} \quad \text{Apply integration by parts twice} \\ &= \sqrt{12}. \end{aligned}$$

Assume that the equality holds. Then, there exists some  $\alpha \in \mathbb{R}$  such that for almost every  $x \in [0, \infty)$ , we have

$$e^{-x}(x^2 + 2) = \alpha e^{-x}(x^2 + 1).$$

With this, we see

$$\begin{aligned} x^2 + 2 &= \alpha x^2 + \alpha \implies x^2(1 - \alpha) = 2 + \alpha \\ &\implies x^2 = \frac{1 - \alpha}{2 + \alpha}. \end{aligned}$$

Thus, for any given  $\alpha$ , this can only be true for at most two points in  $[0, \infty)$ , which contradicts our assumption that this is true almost everywhere. Therefore, we can conclude that the equality does not hold, as desired.  $\square$

## Problem 6

Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  be continuous and bounded. Show that

$$\|f\|_\infty = \sup\{|f(x)| : x \in \mathbb{R}^n\}.$$

### Solution

Let  $M = \sup\{|f(x)| : x \in \mathbb{R}^n\}$ , and let  $\alpha < M$ . Then, by definition of supremum, we have

$$\begin{aligned} \{x \in \mathbb{R}^n : |f(x)| > M\} &= \emptyset \implies |\{x \in \mathbb{R}^n : |f(x)| > M\}| = 0 \\ &\implies M \geq \|f\|_\infty. \end{aligned}$$

Thus, we just need to show that  $M \leq \|f\|_\infty$ , and our proof will be complete. To do this, let's suppose for sake of contradiction that  $M > \|f\|_\infty$ . Then, there exists some nonnegative  $\alpha < M$  such that

$$|\{x \in \mathbb{R}^n : |f(x)| > \alpha\}| = 0.$$

We have

$$\{x \in \mathbb{R}^n : |f(x)| > \alpha\} = |f|^{-1}((\alpha, \infty)).$$

Since  $f$  is continuous,  $|f|$  is continuous. Thus, since  $(\alpha, \infty)$  is open, we can conclude that  $|f|^{-1}((\alpha, \infty))$  is open. Now, by definition of supremum, we have that there exists some  $x \in \mathbb{R}^n$  such that  $\alpha < |f(x)| \leq M$ . Thus,  $|f|^{-1}((\alpha, \infty))$  is nonempty. Since nonempty open subsets of  $\mathbb{R}^n$  have positive measure, we can conclude that

$$|\{x \in \mathbb{R}^n : |f(x)| > \alpha\}| > 0$$

and we have reached a contradiction. Thus,  $M \leq \|f\|_\infty$ , and our proof is complete.

## Problem 7

**Theorem 2.** Let  $\{f_k\} \subset L^p(E)$  for some  $1 \leq p < \infty$ . Assume that  $|E| < \infty$ ,  $\|f_k\|_p \leq A$  for each natural  $k$ , and that

$$\lim_{k \rightarrow \infty} f_k(x) = f(x)$$

for almost every  $x \in E$ . Then  $f_k \rightarrow f$  in  $L^1$ .

### Solution

To prove this, we will start by proving an intuitive lemma.

**Lemma 1.** Let  $E \subseteq \mathbb{R}^n$ , and let  $f \in L(E)$ . Then for any  $\epsilon > 0$ , there exists a  $\delta > 0$  such that if  $A \subseteq E$  with  $|A| < \delta$ , then

$$\int_A |f| < \epsilon.$$

Informally, this is known as the fact that integrals over small sets are small.

*Proof.* Let  $\epsilon > 0$ , and let  $A \subseteq E$ . Let  $g$  be a simple function with  $0 \leq g \leq |f|$ . Then

$$\int_A (|f| - g) \leq \int_E (|f| - g) \implies \int_A |f| \leq \int_A g + \int_E |f| - \int_E g.$$

By theorem 4.13 in our textbook, and the monotone convergence theorem, we can choose  $g$  such that

$$\int_E |f| - \int_E g < \frac{\epsilon}{2}.$$

If we define  $M = \max g$ , then

$$\int_A g \leq |A|M.$$

Thus, if we let  $\delta = \frac{\epsilon}{2M}$ , and  $A$  is such that  $|A| < \delta$ , then we have

$$\begin{aligned} \int_A |f| &\leq \int_A g + \int_E |f| - \int_E g \\ &< \frac{\epsilon}{2} + \frac{\epsilon}{2} \\ &= \epsilon, \end{aligned}$$

and our proof is complete. □

Now we are ready for the proof of the main theorem.

*Proof.* Let  $q$  be the conjugate exponent of  $p$ , and let  $F \subseteq E$  be measurable. If  $p > 1$ , we have

$$\begin{aligned} \int_F |f_k| &\leq \left( \int_F 1 \right)^{1/q} \left( \int_F |f|^p \right)^{1/p} && \text{By Holder's inequality} \\ &\leq \left( \int_F 1 \right)^{1/q} \left( \int_E |f|^p \right)^{1/p} \\ &= |F|^{1/q} \|f_k\|_p \\ &\leq |F|^{1/q} A. \end{aligned}$$

Let  $\epsilon > 0$ . By Ergorov's theorem, there exists a closed set  $F \subseteq E$  such that  $f_k$  converges uniformly to  $f$  on  $F$ , and  $|E \setminus F| < \epsilon$ . We have

$$\begin{aligned}
 \lim_{k \rightarrow \infty} \int_E |f_k - f| &= \lim_{k \rightarrow \infty} \int_F |f_k - f| + \int_{E \setminus F} |f_k - f| \\
 &= 0 + \lim_{k \rightarrow \infty} \int_{E \setminus F} |f_k - f| && \text{By the uniform convergence theorem} \\
 &\leq \int_{E \setminus F} |f_k| + \int_{E \setminus F} |f| \\
 &\leq |E \setminus F|^{1/q} A + \int_{E \setminus F} \liminf |f_k| \\
 &\leq |E \setminus F|^{1/q} A + \liminf \int_{E \setminus F} |f_k| && \text{By Fatou's Lemma} \\
 &\leq 2|E \setminus F|^{1/q} A \\
 &\leq 2\epsilon^{1/q} A.
 \end{aligned}$$

Thus, taking the limit as  $\epsilon \rightarrow 0$ , we have proven that  $f_k$  converges to  $f$  in  $L^1$  if  $p > 1$ .

Now consider the case of  $p = 1$ . Then, we have

$$\begin{aligned}
 \lim_{k \rightarrow \infty} \int_E |f_k - f| &= \lim_{k \rightarrow \infty} \int_F |f_k - f| + \int_{E \setminus F} |f_k - f| \\
 &= 0 + \lim_{k \rightarrow \infty} \int_{E \setminus F} |f_k - f| && \text{By the uniform convergence theorem} \\
 &\leq \int_{E \setminus F} |f_k| + \int_{E \setminus F} |f| \\
 &\leq \int_{E \setminus F} |f_k| + \int_{E \setminus F} \liminf |f_k| \\
 &\leq \int_{E \setminus F} |f_k| + \liminf \int_{E \setminus F} |f_k| && \text{By Fatou's Lemma.}
 \end{aligned}$$

Thus, taking the limit as  $\epsilon \rightarrow 0$ , the above lemma allows us to conclude that  $f_k$  converges to  $f$  in  $L^1$ , and our proof is complete.  $\square$

## Problem 8

**Theorem 3.** Let  $1 \leq p < \infty$  and assume that  $f \in L^p(\mathbb{R})$ . Then

$$\lim_{t \rightarrow \infty} \int_t^{t+1} f(x) dx = 0.$$

### Solution

*Proof.* We have

$$\begin{aligned}
 \lim_{t \rightarrow \infty} \int_t^{t+1} |f(x)| dx = 0 &\implies \lim_{t \rightarrow \infty} \left| \int_t^{t+1} f(x) dx \right| = 0 \\
 &\implies \lim_{t \rightarrow \infty} \int_t^{t+1} f(x) dx = 0,
 \end{aligned}$$

so it will suffice to show that  $\lim_{t \rightarrow \infty} \int_t^{t+1} f(x) dx = 0$ .

Suppose, for sake of contradiction, that  $\lim_{t \rightarrow \infty} \int_t^{t+1} f(x) dx \neq 0$ . Then, there exists some  $\epsilon > 0$  such that for any natural  $N$ , there exists an  $t \geq N$  such that

$$\int_t^{t+1} f(x) dx > \epsilon.$$

Let  $t_0$  be any natural number such that

$$\int_{t_0}^{t_0+1} f(x)dx > \epsilon,$$

and define a sequence  $\{t_n\}_{n=1}^{\infty}$  such that  $t_n \geq t_{n-1} + 1$ , and

$$\int_{t_n}^{t_n+1} f(x)dx > \epsilon.$$

Then, the intervals  $[t_n, t_n + 1]$  are non overlapping, and we have

$$\begin{aligned} \int_{\mathbb{R}} |f(x)|dx &\geq \sum_{n=0}^{\infty} \int_{t_n}^{t_n+1} f(x)dx \\ &> \sum_{n=0}^{\infty} \epsilon \\ &= \infty. \end{aligned}$$

Thus,  $f \notin L^1$  and by theorem 8.2  $f \notin L^p$ , which is a contradiction. Thus, we have proven the theorem.  $\square$

## Problem 9

**Theorem 4.** Let  $p, q, r \in [1, \infty)$  such that

$$\frac{1}{r} = \frac{1}{p} + \frac{1}{q}.$$

If  $f \in L^p(\mathbb{R}^n)$  and  $g \in L^q(\mathbb{R}^n)$ , then

$$\|fg\|_r \leq \|f\|_p \cdot \|g\|_q.$$

### Solution

*Proof.* Define  $p' = \frac{p}{r}$  and  $q' = \frac{q}{r}$ . Then  $p'$  and  $q'$  are conjugate exponents. We have

$$\begin{aligned} \|fg\|_r &= \left( \int_{\mathbb{R}^n} |fg|^r \right)^{1/r} \\ &= \left( \int_{\mathbb{R}^n} |f^r g^r| \right)^{1/r} \\ &\leq \left( \left( \int_{\mathbb{R}^n} |f^r|^{p'} \right)^{1/p'} \left( \int_{\mathbb{R}^n} |g^r|^{q'} \right)^{1/q'} \right)^{1/r} && \text{By Holder's inequality} \\ &= \left( \int_{\mathbb{R}^n} |f|^{rp'} \right)^{1/rp'} \left( \int_{\mathbb{R}^n} |g|^{rq'} \right)^{1/rq'} \\ &= \left( \int_{\mathbb{R}^n} |f|^p \right)^{1/p} \left( \int_{\mathbb{R}^n} |g|^q \right)^{1/q} \\ &= \|f\|_p \cdot \|g\|_q, \end{aligned}$$

and our proof is complete.  $\square$

## Problem 10

**Theorem 5.** Let  $1 \leq p < r < q < +\infty$  and define  $\theta \in (0, 1)$  by

$$\frac{1}{r} = \frac{\theta}{p} + \frac{1-\theta}{q}.$$

Let  $f \in L^p \cap L^q$ . Prove that

$$\|f\|_r \leq \|f\|_p^\theta \cdot \|f\|_q^{1-\theta}.$$

**Solution**

*Proof.* Define  $p' = \frac{p}{\theta}$  and  $q' = \frac{q}{1-\theta}$ . Then, we have

$$\frac{1}{r} = \frac{1}{p'} + \frac{1}{q'}.$$

Utilizing the results of the previous problem, we have

$$\begin{aligned} \|f\|_r &= \|f^\theta f^{1-\theta}\|_r \\ &\leq \|f^\theta\|_{p'} \cdot \|f^{1-\theta}\|_{q'} \\ &= \left( \int_{\mathbb{R}^n} |f^\theta|^{p'} \right)^{1/p'} \left( \int_{\mathbb{R}^n} |f^{1-\theta}|^{q'} \right)^{1/q'} \\ &= \left( \int_{\mathbb{R}^n} |f|^{\theta p'} \right)^{1/p'} \left( \int_{\mathbb{R}^n} |f|^{(1-\theta)q'} \right)^{1/q'} \\ &= \left( \int_{\mathbb{R}^n} |f|^p \right)^{\theta/p} \left( \int_{\mathbb{R}^n} |f|^q \right)^{(1-\theta)/q} \\ &= \|f\|_p^\theta \cdot \|f\|_q^{1-\theta}, \end{aligned}$$

and our proof is complete. □

**Problem 11**

**Theorem 6.** Let  $0 < r < \infty$  and  $f \in L^r(\mathbb{R}^n) \cap L^\infty(\mathbb{R}^n)$ . Then the following statements are true:

Part (i):

For all  $p \in (r, \infty)$ ,

$$\|f\|_p \leq \|f\|_r^{r/p} \cdot \|f\|_\infty^{1-r/p}.$$

Part (ii):

$$\lim_{p \rightarrow \infty} \|f\|_p = \|f\|_\infty.$$

**Solution**

*Proof. Part (i):*

We have

$$\begin{aligned} \|f\|_p &= \|f^{r/p} f^{1-r/p}\|_p \\ &= \left( \int_{\mathbb{R}^n} |f^{r/p} f^{1-r/p}|^p dx \right)^{1/p} \\ &= \left( \int_{\mathbb{R}^n} |f^r f^{p-r}| dx \right)^{1/p} \\ &\leq \left( \|f^{p-r}\|_\infty \cdot \int_{\mathbb{R}^n} |f^r| dx \right)^{1/p} && \text{Holder's Inequality} \\ &= \left( \|f\|_\infty^{p-r} \cdot \int_{\mathbb{R}^n} |f|^r dx \right)^{1/p} \\ &= \left( \|f\|_\infty^{p-r} \cdot \|f\|_r^r \right)^{1/p} \\ &= \|f\|_r^{r/p} \cdot \|f\|_\infty^{1-r/p}, \end{aligned}$$

as desired.

*Part (ii):*

Using the results from the previous part, we have

$$\begin{aligned}\lim_{p \rightarrow \infty} \|f\|_p &\leq \lim_{p \rightarrow \infty} \|f\|_r^{r/p} \cdot \|f\|_\infty^{1-r/p} \\ &= \|f\|_r^0 \cdot \|f\|_\infty^{1-0} \\ &= \|f\|_\infty.\end{aligned}$$

Thus, all we must do to complete the proof is show that  $\lim_{p \rightarrow \infty} \|f\|_p \geq \|f\|_\infty$  (but I've not figured this part out yet).  $\square$