

Mathematical notes

Ancillary material for “The Last Human Gate”

Jeremy Canale

September 2026

These notes derive the algebraic statements used in the paper. Results are referred to by name, as in the manuscript. Nothing here is estimated from data.

1 Workload ratio and break-even thresholds

With common visits λ and useful hours K , the labor account gives

$$\rho = \frac{N_A}{N_H} = \frac{\lambda \bar{h}_A + B_A}{\lambda h_0 + B_H} = \frac{\bar{h}_A/h_0 + b_A}{1 + b_H}, \quad b_s = \frac{B_s}{\lambda h_0}.$$

Write $h_E = \kappa \eta h_0$, $h_R = \mu h_S^H = \mu \sigma h_0 = m h_0$ and $h_W = r h_0$. Then $\bar{h}_A/h_0 = q\kappa\eta + (1-q)\sigma\mu + r$. The baseline partition states $q\kappa + (1-q)\sigma = 1$, so $(1-q)\sigma = 1 - \phi$ with $\phi = q\kappa$. Substituting gives both forms of the workload ratio:

$$\rho = \frac{q\kappa\eta + (1-q)\sigma\mu + r + b_A}{1 + b_H} = \frac{\phi\eta + (1-\phi)\mu + r + b_A}{1 + b_H}.$$

Exception-share threshold. In overall-mean units, $\rho < 1$ is $qk + (1-q)m + r + b_A < 1 + b_H$, that is $q(k - m) < 1 + b_H - m - r - b_A$. For $k > m$ this is the exception-share threshold $q < q^*$; for $k < m$ the inequality reverses; for $k = m$ the case share drops out. Solving the same equality for k at a given $q > 0$ gives $k^* = m + (1 + b_H - m - r - b_A)/q$, the line drawn in the frontier figure, once per route.

Exception-effort margin. In group-relative units, $\rho < 1$ is $\phi\eta < 1 + b_H - (1 - \phi)\mu - r - b_A$. For $\phi > 0$,

$$\eta^* = \frac{1 + b_H - (1 - \phi)\mu - r - b_A}{\phi} = \mu + \frac{1 + b_H - \mu - r - b_A}{\phi}, \quad \frac{\partial \eta^*}{\partial \phi} = -\frac{1 + b_H - \mu - r - b_A}{\phi^2}.$$

The margin therefore falls with ϕ exactly when $\mu + r + b_A < 1 + b_H$, that is, when a process made only of standard cases would save labor. In the synthetic example m is held common, so $(1 - \phi)\mu = (1 - q)m$ and the code evaluates $\eta^* = [1 + b_H - (1 - q)m - r - b_A]/\phi$.

2 Selection identity

Let $T \geq 0$ be baseline handling time with $\mathbb{E}[T] = h_0 > 0$, $J \in \{0, 1\}$ the residual indicator, $e(T) = \Pr(J = 1 \mid T)$ and $q = \mathbb{E}[e(T)] > 0$. Iterated expectations give $\mathbb{E}[TJ] = \mathbb{E}[Te(T)] = \text{Cov}(T, e(T)) + h_0 q$. Dividing by q yields $\mathbb{E}[T \mid J = 1] - h_0 = \text{Cov}(T, e(T))/q$; dividing by h_0 yields $\phi = q + \text{Cov}(T, e(T))/h_0$. Since $\kappa = \mathbb{E}[T \mid J = 1]/h_0$, also $\phi = q\kappa$. For independent copies T, T' , $2 \text{Cov}(T, e(T)) = \mathbb{E}[(T - T')(e(T) - e(T'))] \geq 0$ when e is nondecreasing, because the two factors then share a sign. The identities restrict first moments and one cross-moment only: truncating a distribution to a narrow band of difficult cases raises the conditional mean and can lower the conditional variance.

3 Strategic gate model

Setup. Let ε have distribution function F and density f . With $s = \xi + u + g$, approval probability is $p(s) = \Pr(s + \varepsilon \geq \tau) = 1 - F(\tau - s)$, so $p'(s) = f(\tau - s) \geq 0$ and $p''(s) = -f'(\tau - s)$. The project maximizes $\Pi(u, g) = Vp(s) - c_u(u) - c_g(g) - D(g, a)$ over $u, g \geq 0$.

First- and second-order conditions. At an interior optimum,

$$Vp'(s) - c'_u(u) = 0, \quad Vp'(s) - c'_g(g) - D_g(g, a) = 0.$$

Put $t = Vp''(s)$, $\omega_u = c''_u(u) > 0$ and $\omega_g = c''_g(g) + D_{gg}(g, a) > 0$. The Hessian is

$$H = \begin{pmatrix} t - \omega_u & t \\ t & t - \omega_g \end{pmatrix}, \quad \det H = \omega_u \omega_g - t(\omega_u + \omega_g) = \Delta.$$

A strict local maximum requires $t - \omega_u < 0$ and $\Delta > 0$. Both hold automatically when $t \leq 0$.

Assurance intensity. Differentiating the two conditions with respect to a ,

$$H \begin{pmatrix} du \\ dg \end{pmatrix} = \begin{pmatrix} 0 \\ D_{ga} da \end{pmatrix} \implies \frac{\partial u^*}{\partial a} = -\frac{t D_{ga}}{\Delta}, \quad \frac{\partial g^*}{\partial a} = \frac{(t - \omega_u) D_{ga}}{\Delta}, \quad \frac{\partial s^*}{\partial a} = -\frac{\omega_u D_{ga}}{\Delta}.$$

With $D_{ga} > 0$: cosmetic effort falls whenever the second-order conditions hold; the observable score falls as well; genuine remediation rises weakly if and only if $t \leq 0$. If $t > 0$ (and the second-order conditions hold), both efforts fall.

Lemma 1. *If f is differentiable and unimodal with mode zero, then $p''(s) \leq 0$ if and only if $s \geq \tau$.*

Proof. Unimodality with mode zero means $f'(x) \geq 0$ for $x \leq 0$ and $f'(x) \leq 0$ for $x \geq 0$. Since $p''(s) = -f'(\tau - s)$, the sign condition is $\tau - s \leq 0$. $\square$

The redirection result therefore applies to projects whose expected score already meets the threshold; below it, stronger assurance can discourage both kinds of effort.

Corner solutions. If $g^* = 0$ because $Vp'(s) \leq c'_g(0) + D_g(0, a)$, a marginal increase in a leaves (u^*, g^*) unchanged. If $u^* = 0$ is binding and $g^* > 0$, the single remaining condition gives $\partial g^* / \partial a = D_{ga} / (t - \omega_g) < 0$ under its second-order condition $t - \omega_g < 0$.

Clearer criteria. Let a clarity parameter θ lower both marginal costs: $c'_{u\theta} < 0$ and $c'_{g\theta} < 0$. Then

$$\frac{\partial u^*}{\partial \theta} = \frac{c'_{u\theta}(t - \omega_g) - t c'_{g\theta}}{\Delta}, \quad \frac{\partial g^*}{\partial \theta} = \frac{(t - \omega_u) c'_{g\theta} - t c'_{u\theta}}{\Delta}.$$

For $t \leq 0$ the first term of each numerator is positive and the second is nonpositive, so neither sign is determined: clearer criteria can raise genuine work, cosmetic work, or both.

Numerical check. `test_reproduce.py` solves the model with quadratic costs, logistic noise and $D(g, a) = ag^2/2$ at two assurance levels and confirms the predicted signs in the region $s > \tau$.

4 Reliability bounds and the readiness example

For events $\mathcal{C}_1, \dots, \mathcal{C}_n$ with $\Pr(\mathcal{C}_i) = p_i$, the Fréchet inequalities give $\max(0, 1 - \sum_i (1 - p_i)) \leq \Pr(\bigcap_i \mathcal{C}_i) \leq \min_i p_i$. With $n = 20$ and $p_i = 0.99$ the bounds are 0.80 and 0.99; independence gives $0.99^{20} \approx 0.8179$.

The geometric readiness index of $(0.01, 1, 1, 1, 1)$ is $0.01^{1/5} = 10^{-0.4} \approx 0.398$, which shows that the index is only partially sensitive to a near-zero dimension and motivates reporting a minimum index and hard blockers alongside it.