

Basis and Linear Independence

Jeremy Teitelbaum

Basis

A set of vectors in \mathbf{R}^n (or in any vector space V) is called a **basis** if

- ▶ it spans V
- ▶ it is linearly independent.

Examples: if A is an invertible $n \times n$ matrix, its columns are linearly independent and span \mathbf{R}^n and therefore are a basis for \mathbf{R}^n .

The vectors $1, x, x^2, \dots, x^n$ span the polynomials of degree at most n and are linearly independent.

The “standard vectors” e_i for $i = 1, \dots, n$ are a basis for \mathbf{R}^n .

Subspace basis

The vectors $(1, 3, 2)$ and $(-1, -1, 0)$ are linearly independent and span a subspace H of \mathbf{R}^3 .

Therefore they are a basis for H .

Every spanning set contains a basis

If a set S of vectors v_1, \dots, v_n spans a subspace H , then a subset of S is a basis.

Proof: If the vectors are linearly independent, they are already a basis.

If they are dependent, then one is a linear combination of the others. Remove that one from S . The result still spans.

Continue removing dependent vectors until the remaining vectors are independent, and you've found your basis.

A basis is a minimal spanning set

If H is a subspace of V , suppose you have a bunch of vectors in H .

Too many vectors makes them dependent. Too few means they can't span. If they are a basis, there are enough to span, but not to become dependent.

Basis for $\text{Nul}(A)$.

The null space of A is spanned by the vectors with weights given by the free variables in the row reduced form of A .

Those vectors are independent and therefore form a basis.

Basis for $\text{Col}(A)$.

Given vectors v_1, \dots, v_k , make an $m \times k$ matrix with the v_i as columns.

To find a linear relation among the columns of A , we need to solve $Ax = 0$.

But $Ax = 0$ if and only if $EAx = 0$ where E is an elementary matrix.

Put another way, row reduction doesn't change the x such that $Ax = 0$.

So we can assume A is in row reduced echelon form.

More on basis for $\text{Col}(A)$.

Once A is in row reduced form, we see that:

- ▶ the columns corresponding to free variables are linear combinations of the pivot columns
- ▶ the pivot columns are linearly independent.

Basis for $\text{Col}(A)$.

The **columns of** A corresponding to the pivot columns in the row reduced version of A are a basis for the column space. (note that these are *not* the columns of the reduced matrix).

So: a basis for the null space is made up of k vectors where k is the number of free variables, and a basis for the column space is made up of r vectors where r is the number of pivot columns.

Notice that $k + r = n$ where n is the total number of columns of A .

Example

Suppose that

$$A = \begin{bmatrix} 2 & 4 & 5 & 1 \\ 1 & -3 & -2 & 0 \\ 0 & -2 & -3 & 1 \end{bmatrix}$$

The row reduced form of A is

$$\begin{bmatrix} 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & -1 \end{bmatrix}$$

Since the first three columns are pivot columns, the first three columns of A span the column space of A , and the last column satisfies $c_4 = c_1 + c_2 - c_3$.

Example continued

The nullspace of A is the solution to the homogeneous system, and it is given by the equations

$$x_1 = -x_4$$

$$x_2 = -x_4$$

$$x_3 = x_4$$

so the null space is spanned by

$$\begin{bmatrix} -1 \\ -1 \\ 1 \\ 1 \end{bmatrix}$$

Null Space and Col Space

Contrast Between Nul A and Col A for an $m \times n$ Matrix A

Nul A	Col A
1. Nul A is a subspace of \mathbb{R}^n .	1. Col A is a subspace of \mathbb{R}^m .
2. Nul A is implicitly defined; that is, you are given only a condition ($A\mathbf{x} = \mathbf{0}$) that vectors in Nul A must satisfy.	2. Col A is explicitly defined; that is, you are told how to build vectors in Col A .
3. It takes time to find vectors in Nul A . Row operations on $[A \ \mathbf{0}]$ are required.	3. It is easy to find vectors in Col A . The columns of A are displayed; others are formed from them.
4. There is no obvious relation between Nul A and the entries in A .	4. There is an obvious relation between Col A and the entries in A , since each column of A is in Col A .
5. A typical vector \mathbf{v} in Nul A has the property that $A\mathbf{v} = \mathbf{0}$.	5. A typical vector \mathbf{v} in Col A has the property that the equation $A\mathbf{x} = \mathbf{v}$ is consistent.
6. Given a specific vector \mathbf{v} , it is easy to tell if \mathbf{v} is in Nul A . Just compute $A\mathbf{v}$.	6. Given a specific vector \mathbf{v} , it may take time to tell if \mathbf{v} is in Col A . Row operations on $[A \ \mathbf{v}]$ are required.
7. Nul $A = \{\mathbf{0}\}$ if and only if the equation $A\mathbf{x} = \mathbf{0}$ has only the trivial solution.	7. Col $A = \mathbb{R}^m$ if and only if the equation $A\mathbf{x} = \mathbf{b}$ has a solution for every \mathbf{b} in \mathbb{R}^m .
8. Nul $A = \{\mathbf{0}\}$ if and only if the linear transformation $\mathbf{x} \mapsto A\mathbf{x}$ is one-to-one.	8. Col $A = \mathbb{R}^m$ if and only if the linear transformation $\mathbf{x} \mapsto A\mathbf{x}$ maps \mathbb{R}^n onto \mathbb{R}^m .

Figure 1: Null Space vs Col Space

Row space

The *row space* of a matrix is the span of its rows.

Row operations do not change the row space, so one can find a basis for the row space of A by putting A in reduced form.

The rows with a pivot (that is, the nonzero rows) form a basis for the row space.

This is because they are clearly linearly independent (and they span by definition).