

Math 2710

Sep 16-20

Characterization of the gcd

Proposition: (2.29, page 34) Suppose $b \neq 0$. An integer d is the greatest common divisor of a and b if and only if

- $d \geq 0$
- d is a common divisor of a and b
- If r is a common divisor of a and b , then $r|d$.

Proof: First suppose that these three conditions are true. Then d is a common divisor, and by Proposition 2.1 (iv), if r is any other common divisor of a and b , then $r|d$ so $|r| \leq d$. So d is the greatest common divisor.

Now suppose d is the greatest common divisor of a and b . Then $d \geq 0$ and d is a common divisor, so we just need to check the third condition. By the extended euclidean algorithm there are x and y so that $ax + by = d$. By Proposition 2.1 (ii), any common divisor of a and b divides $ax + by = d$, as we wanted to show.

Least common multiple

Definition: A common multiple of two integers a and b , with $b \neq 0$, is any integer m such that $a|m$ and $b|m$. The **least common multiple** of a and b is the smallest positive integer which is a common multiple of a and b .

Theorem: The lcm of a and b is $|ab/g|$ where g is the gcd of a and b .

Proof: We can assume a and b are non-negative as this does not affect the lcm. Because g divides both a and b , we have $ab/g = a(b/g) = b(a/g)$ so ab/g is an integer and it is a common multiple of a and b . Now let t be any common multiple of a and b . Find x and y so that $ax + by = g$. Then $tax + tby = tg$. Since t is a common multiple of a and b , we have tax and tby are both multiples of ab . So $tax + tby = abs$ for some integer s . We conclude that $t = (ab/g)s$, so that t is a multiple of ab/g . This means $t \geq (ab/g)$ so ab/g must be the least common multiple.

Linear Diophantine Equations

A *diophantine equation* is an equation where the variables are restricted to integer values.

A linear diophantine equation in one variable is of the form

$$ax = b$$

where a and b are integers and we want x to be an integer. Clearly this has a solution exactly when $a|b$.

Linear Diophantine Equations in 2 variables

A linear diophantine equation in two variables is an equation of the form

$$ax + by = c$$

where a , b , and c are integers.

Solving such an equation means finding *integers* x and y that satisfy the condition.

Theorem on Linear Diophantine Equations

Theorem:

- The linear diophantine equation $ax + by = c$ has a solution if and only if $\gcd(a, b)|c$.
- If x_0, y_0 is one solution to the equation, and x and y is any other solution, then there exists an integer n so that

$$x = x_0 + n\frac{b}{d} \quad \text{and} \quad y = y_0 - n\frac{a}{d}$$