

Higher Theory and Applications of Differential Equations
MATH2221 UNSW

Jeremy Le

2024T2

Contents

1	Linear ODEs	3
1.1	Introduction	3
1.2	Linear Differential Operators	4
1.3	Differential Operators with Constant Coefficients	6
1.4	Wronskians and Linear Independence	7
1.5	Methods for Inhomogeneous Equations	8
1.5.1	Judicious Guessing Method	8
1.5.2	Annihilator Method	9
1.5.3	Judicious Guessing Method Continued	10
1.5.4	Variation of Parameters	10
1.6	Solution via Power Series	11
1.7	Singular ODEs	11
1.8	Bessel and Legendre Equations	13
1.8.1	Bessel Equations and Functions	13
1.8.2	Legendre Equation	14
2	Dynamical Systems	15
2.1	Terminology	15
2.2	Existence and Uniqueness	16
2.3	Linear Dynamical Systems	18

Chapter 1

Linear ODEs

1.1 Introduction

Recall that a first-order ordinary differential equation (ODE) has, in its most general realisation, the form

$$y'(t) = f(t, y(t)).$$

A special case is the equation

$$a(t)y'(t) + b(t)y(t) = f(t),$$

with $a(t) \neq 0$ on some interval $I \in \mathbb{R}$. This special first-order ODE is called a **linear first-order ODE**. Another special case is

$$y'(t) = f(t)g(y),$$

which is known as a **separable first-order ODE**.

For a separable equation the solution is found (at least, implicitly by) writing:

$$\int \frac{1}{g(y)} dy = \int f(t) dt.$$

Solving Seperable ODEs Consider $y' = t^2y, y(0) = 3$. This is seperable with $f(t) = t^2$ and $g(y) = y$. Then

$$\int \frac{1}{y} dy = \int t^2 dt$$

so that

$$\ln |y(t)| = \frac{1}{3}t^3 + C.$$

Now apply e^t to both sides to obtain

$$|y(t)| = e^{\frac{1}{3}t^3 + C} = e^C e^{\frac{1}{3}t^3}.$$

Thus, a general solution of the equation is

$$y(t) = Ae^{\frac{1}{3}t^3}.$$

Since $y(0) = 3$, we see that the unique solution is $y(t) = 3e^{\frac{1}{3}t^3}$.

In the case of a linear first-order equation, i.e. $y' + a(t)y = f(t)$, a useful solution method is the integrating factor technique. The idea is to find a function μ so that when we multiply both sides of the equation with μ we find that

$$[\mu y](t)' = \mu(t)(y' + a(t)y) = \mu(t)f(t),$$

for if this happens, then the general solution of the ODE should be

$$y(t) = \frac{1}{\mu(t)} \int \mu(t)f(t) dt + \frac{C}{\mu(t)}.$$

Solving Linear First-Order ODE Solve $y' - 2ty = 3t$. We pick

$$\mu(t) = e^{\int -2t dt} = e^{-t^2}.$$

Then

$$\begin{aligned}(e^{-t^2}y)' &= 3te^{-t^2} \\ e^{-t^2}y &= \int 3te^{-t^2} dt = -\frac{3}{2}e^{-t^2} + C \\ y(t) &= -\frac{3}{2} + Ce^{t^2}.\end{aligned}$$

1.2 Linear Differential Operators

In linear algebra, you have seen the compact notation $A\mathbf{x} = \mathbf{b}$ for system of linear equations. A similar notation when dealing with a linear ordinary differential equations is

$$Lu = f.$$

Here, L is an operator (or transformation) that acts on a function u to create a new function Lu .

Given coefficients $a_0(x), a_1(x), \dots, a_m(x)$ we define the **linear differential operator** L of **order** m ,

$$\begin{aligned}Lu(x) &= \sum_{j=0}^m a_j(x) D^j u(x) \\ &= a_m D^m u + a_{m-1} D^{m-1} u + \dots + a_0 u,\end{aligned}$$

where $D^j u = d^j u / dx^j$ (with $D^0 u = u$).

We refer to a_m as the **leading coefficient** of L and assume that each $a_j(x)$ is a smooth function of x .

The ODE $Lu = f$ is said to be **singular** with respect to an interval $[a, b]$ if the leading coefficient $a_m(x)$ vanishes for any $x \in [a, b]$.

Example $Lu = (x - 3)u''' - (1 + \cos x)u' + 6u$ is a linear differential of order 3, with leading coefficient $x - 3$. Thus, L is singular on $[1, 4]$, but not singular on $[0, 2]$.

Example $N(u) = u'' + u^2 u' - u$ is a nonlinear differential operator of order 2.

Linearity For any constants c_1 and c_2 and any m -times differentiable functions u_1 and u_2 ,

$$L(c_1 u_1 + c_2 u_2) = c_1 L u_1 + c_2 L u_2.$$

Ordinary differential equations of the form $Lu = 0$ are known as **homogenous**. Those of the form $Lu = f$ are known as **inhomogeneous**.

When the solution to a differential equation is prescribed at a particular point $x = x_0$, that is

$$u(x_0) = v_0, \quad u'(x_0) = v_1, \quad \dots, \quad u^{(m-1)}(x_0) = v_{m-1},$$

we call it an **initial value problem**. Where a differential equation is order 2 or greater, solutions at 2 or more locations can be prescribed. Such problems are called **boundary value problems**.

Unique Solution to Linear Initial Problem For an ODE $Lu = f$ which is not singular with respect to a, b , with f continuous on $[a, b]$, the IVP for an m th-order linear differential operator with m initial values has a unique solution.

Solution to m th Order Problem has Dimension m Assume that the linear, m th-order differential operator L is not singular on $[a, b]$. Then the set of all solutions to the homogenous equation $Lu = 0$ on $[a, b]$ is a vector space of dimension m .

If $\{u_1, u_2, \dots, u_m\}$ is **any** basis for the solution space of $Lu = 0$, then every solution can be written in a unique way as

$$u(x) = c_1 u_1(x) + c_2 u_2(x) + \dots + c_m u_m(x) \quad \text{for } a \leq x \leq b.$$

We refer to this as the **general solution** of the homogenous equation $Lu = 0$ on $[a, b]$.

Linear superposition refers to this technique of constructing a new solution out of a linear combination of old ones.

Example The general solution to $u'' - u' - 2u = 0$ is $u(x) = c_1 e^{-x} + c_2 e^{2x}$.

Consider the inhomogeneous equation $Lu = f$ on $[a, b]$, and fix a particular solution u_P . For *any* solution u , the difference $u - u_P$ is a solution of the homogenous equation because

$$L(u - u_P) = Lu - Lu_P = f - f = 0 \text{ on } [a, b].$$

Hence, $u(x) - u_P(x) = c_1 u_1(x) + \dots + c_m u_m(x)$ for some constants c_1, \dots, c_m and so

$$u(x) = u_P(x) + \underbrace{c_1 u_1(x) + \dots + c_m u_m(x)}_{u_H(x)}, \quad a \leq x \leq b,$$

is the **general solution** of the inhomogeneous equation $Lu = f$.

Example The inhomogeneous ODE $u'' - u' - 2u = -2e^x$ has a particular solution $u_P(x) = e^x$. The general solution for its homogenous counterpart is $u_H(x) = c_1 e^{-x} + c_2 e^{2x}$. So the general solution of the inhomogeneous ODE is

$$u(x) = u_P(x) + u_H(x) = e^x + c_1 e^{-x} + c_2 e^{2x}.$$

Reduction of Order For $u = u_1(x) \neq 0$, a solution to the ODE

$$u'' + p(x)u' + q(x)u = 0,$$

on some interval I , then a second solution is

$$u = u_1(x) \int \frac{1}{u_1^2 \exp(\int p dx)} dx.$$

Example For the ODE $u'' - 6u' + 9u = 0$, take $u_1 = e^{3x}$ and find v . **Answer** xe^{3x} .

1.3 Differential Operators with Constant Coefficients

If L has constant coefficients, then the problem of solving $Lu = 0$ reduces to that of factorising the polynomial having the same coefficients.

Suppose that a_j is constant for $0 \leq j \leq m$, with $a_m \neq 0$. We define the associated polynomial of degree m ,

$$p(z) = \sum_{j=0}^m a_j z^j = a_m z^m + a_{m-1} z^{m-1} + \cdots + a_1 z + a_0,$$

so that if

$$Lu = a_m u^{(m)} + a_{m-1} u^{(m-1)} + \cdots + a_1 u' + a_0 u,$$

then formally, $L = p(D)$.

By the fundamental theorem of algebra,

$$p(z) = a_m (z - \lambda_1)^{k_1} (z - \lambda_2)^{k_2} \cdots (z - \lambda_r)^{k_r}$$

where $\lambda_1, \lambda_2, \dots, \lambda_r$ satisfying

$$k_1 + k_2 + \cdots + k_r = m.$$

Lemma $(D - \lambda)x^j e^{\lambda x} = jx^{j-1}e^{\lambda x}$ for $j \geq 0$.

Lemma $(D - \lambda)^k x^j e^{\lambda x} = 0$ for $j = 0, 1, \dots, k - 1$.

Basic Solutions If $(z - \lambda)^k$ is a factor of $p(z)$ then the function $u(x) = x^j e^{\lambda x}$ is a solution of $Lu = 0$ for $0 \leq j \leq k - 1$.

General Solution For the constant-coefficient case, the general solution of the homogenous equation $Lu = 0$ is

$$u(x) = \sum_{q=1}^r \sum_{l=0}^{k_q-1} c_{ql} x^l e^{\lambda_q x},$$

where the c_{ql} are arbitrary constants.

Repeated Real Root From the factorisation

$$D^4 + 6D^3 + 9D^2 - 4D - 12 = (D - 1)(D + 2)^2(D + 3)$$

we see that the general solution of

$$u'''' + 6u''' + 9u'' - 4u' - 12u = 0$$

is

$$u = c_1 e^x + c_2 e^{-2x} + c_3 x e^{-2x} + c_4 e^{-3x}.$$

Complex Root From the factorisation

$$\begin{aligned} D^3 - 7D^2 + 17D - 15 &= (D^2 - 4D + 5)(D - 3) \\ &= (D - 2 - i)(D - 2 + i)(D - 3) \end{aligned}$$

we see that the general solution of

$$u''' - 7u'' + 17u' - 15u = 0$$

is

$$\begin{aligned} u(x) &= c_1 e^{(2+i)x} + c_2 e^{(2-i)x} + c_3 e^{3x} \\ &= c_4 e^{2x} \cos x + c_5 e^{2x} \sin x + c_3 e^{3x}. \end{aligned}$$

Second-order ODEs arise naturally in classical mechanics for example a harmonic simple oscillator.

1.4 Wronskians and Linear Independence

We introduce a function, called the Wronskain that provides us with a way of testing whether a family of solutions to $Lu = 0$ is linearly independent.

Let $u_1(x), u_2(x), \dots, u_m(x)$ be functions defined on an interval $I \in \mathbb{R}$. The functions u_1, \dots, u_m are called **linearly dependent** if there exist constant a_1, a_2, \dots, a_m **not all zero** such that

$$a_1 u_1(x) + a_2 u_2(x) + \dots + a_m u_m(x) = 0 \quad \forall x \in I.$$

If the above equation only holds for

$$a_i = 0, \quad i = 1, 2, \dots, m$$

then the functions are **linearly independent**.

Example $u_1 = \sin 2x$ and $u_2 = \sin x \cos x$ are linearly dependent.
 $u_1 = \sin x$ and $u_2 = \cos x$ are linearly indepdent.

The **Wronskian** of the functions u_1, u_2, \dots, u_m is the $m \times m$ determinant

$$W(x) = W(x; u_1, u_2, \dots, u_m) = \det[D^{i-1}u_j].$$

Example The Wronskian of the functions $u_1 = e^{2x}$, $u_2 = xe^{2x}$ and $u_3 = e^{-x}$ is

$$W = \begin{vmatrix} e^{2x} & xe^{2x} & e^{-x} \\ 2e^{2x} & e^{2x} + 2xe^{2x} & -e^{-x} \\ 4e^{2x} & 4e^{2x} + 4xe^{2x} & e^{-x} \end{vmatrix} = 9e^{3x}.$$

Lemma If u_1, \dots, u_m are linearly dependent over an interval $[a, b]$ then $W(x; u_1, \dots, u_m) = 0$ for $a \leq x \leq b$.

Lemma If u_1, u_2, \dots, u_m are solutions of $Lu = 0$ on the interval $[a, b]$ then their Wronskian satisfies

$$a_m(x)W'(x) + a_{m-1}(x)W(x) = 0, \quad a \leq x \leq b.$$

Linear Independence of Solutions Let u_1, u_2, \dots, u_m be solutions of a non-singular, linear, homogenous, m -th order ODE $Lu = 0$ on the interval $[a, b]$.

Either

$W(x) = 0$ for $a \leq x \leq b$ and the m solutions are linearly **dependent**,
or else

$W(x) \neq 0$ for $a \leq x \leq b$ and the m solutions are linearly **independent**.

1.5 Methods for Inhomogeneous Equations

1.5.1 Judicious Guessing Method

You would have learned the method of undetermined coefficients for constructing a particular solution u_p to an inhomogeneous second-order linear ODE $Lu = f$ in some simple cases. We will study this method systematically for higher-order linear ODEs with constant coefficients.

Superposition of Solutions Suppose that u_1 solves $Lu = e^{3x}$, and u_2 solves $Lu = \sin x$, where L is a linear differential operator. Then the solution of

$$Lu = e^{3x} + \sin x$$

is

$$u(x) = u_1(x) + u_2(x).$$

And a solution of

$$Lu = \frac{1}{2}e^{3x} - 5\sin x$$

is

$$u(x) = \frac{1}{2}u_1(x) - 5u_2(x).$$

Now we want to investigate some methods for finding particular solutions - i.e., finding a solution of $Lu = f$. One such method is the method of judicious guessing. For example:

1. If f is a polynomial, then guess that u_p is a polynomial.
2. If f is a exponential, then guess that u_p is exponential.

3. If f is a sine or cosine, then guess that u_p is a combination of such functions.

One problem with this method: it will only work for the types of functions identified above.

Example Suppose that $u'' - u' = t^2 + 2t$. Note as before that,

$$u_h(t) = c_1 + c_2 e^t.$$

So guess,

$$u_p(t) = At^3 + Bt^2 + Ct + D.$$

Then

$$t^2 + 2t = u_p'' - u_p' = -3At^2 + (6A - 2B)t + (2B - C).$$

So, equating coefficients of like power terms, we see that

$$A = -\frac{1}{3}, B = -2, C = -4, \text{ and } D \text{ is unrestricted.}$$

Therefore, reabsorbing D into c_1 , we see that

$$u(t) = u_h(t) + u_p(t) = c_1 + c_2 e^t - \frac{1}{3}t^3 - 2t^2 - 4t.$$

Now we look at this idea of judicious guessing in a more systematic way. Let $L = p(D)$ be a linear differential operator of order m with constant coefficients.

Polynomial Solutions Assume that $a_0 = p(0) \neq 0$. For any integer $r \geq 0$, there exists a unique polynomial u_P of degree r such that $Lu_P = x^r$.

Exponential Solutions Let $L = p(D)$, $M \in \mathbb{R}$ and $\mu \in \mathbb{C}$. If $p(\mu) \neq 0$, then the function

$$u_P(x) = \frac{Me^{\mu x}}{p(\mu)}$$

satisfies $Lu_P = Me^{\mu x}$.

Example A particular solution of $u'' + 4u' - 3i = 3e^{2x}$ is $u_P = e^{2x}/3$.

Product of Polynomial and Exponential Let $L = p(D)$ and assume that $p(\mu) \neq 0$. For any integer $r \geq 0$, there exists a unique polynomial v of degree r such that $u_P = v(x)e^{\mu x}$ satisfies $Lu_P = x^r e^{\mu x}$.

1.5.2 Annihilator Method

In the previous cases we proposed a solution $u = u_P$ and showed that it satisfied $Lu = f$. The following is a method to derive a particular solution given $Lu = f$. If $f(x)$ is differentiable at least n times and

$$[a_n D^n + a_{n-1} D^{n-1} + \cdots + a_1 D^1 + a_0]f(x) = 0$$

then $[a_n D^n + a_{n-1} D^{n-1} + \cdots + a_1 D^1 + a_0]$ **annihilates** f .

Example D^n annihilates x^{m-1} for $m \leq n$.
 $(D - \alpha)^n$ annihilates $x^{m-1}e^{\alpha x}$ for $m \leq n$.

Annihilator Method: Simple Example Given $Lu = f$ we can apply the appropriate annihilator to both sides and solving the resulting homogenous DE.

Let $Lu = u'' - u'$ and suppose we want a solution such that $Lu = x^2$. Annihilating both sides we have

$$D^3(u'' - u') = u^{(5)} - u^{(4)} = 0.$$

Setting $w = u^{(4)}$, clearly $w = Ce^x$ is the general solution. Integrating four times yields

$$u = Ce^x + Ex^3 + Fx^2 + Gx + H.$$

Clearly $u_h = Ae^x + H$ and the form of the particular solution is $u_P = x(Ex^2 + Fx + G)$. Substituting find $E = -1/3, F = -1$ and $G = -2$.

1.5.3 Judicious Guessing Method Continued

Polynomial Solutions: The Remaining Case Let $L = p(D)$ and assume $p(0) = p'(0) = \dots = p^{(k-1)}(0) = 0$ but $p^{(k)}(0) \neq 0$ where $1 \leq k \leq m - 1$. For any integer $r \geq 0$, there exists a unique polynomial v of degree r such that $u_P(x) = x^K v(x)$ satisfies $Lu_P = x^r$.

Exponential Times Polynomial: Remaining Case Let $L = p(D)$ and assume $p(\mu) = p'(\mu) = \dots = p^{(k-1)}(\mu) = 0$. But $p^{(k)}(\mu) \neq 0$, where $1 \leq k \leq m - 1$. For any integer $r \geq 0$, there exists a unique polynomial v of degree r such that $u_P(x) = x^k v(x)e^{\mu x}$ satisfies $Lu_P = x^r e^{\mu x}$.

1.5.4 Variation of Parameters

Example Find the general solution to $u'' - 4u' + 4u = (x + 1)\exp 2x$.

Note first that the general solution, u_h , to $u'' - 4u' + 4u = 0$ is

$$u(x) = c_1 e^{2x} + c_2 x e^{2x}$$

since the characteristic equation is $0 = r^2 - 4r + 4 = (r - 2)^2$. Then

$$W(x) = \begin{vmatrix} e^{2x} & x e^{2x} \\ 2e^{2x} & e^{2x} + 2x e^{2x} \end{vmatrix} = e^{4x} + 2x e^{4x} - 2x e^{4x} = e^{4x}.$$

So by the method of variation of parameters:

$$v_1'(x) = e^{-4x} \cdot -x e^{2x} (x + 1) e^{2x} \text{ and } v_2'(x) = e^{-4x} \cdot e^{2x} (x + 1) e^{2x}.$$

In other words,

$$v_1'(x) = -x^2 - x \text{ and } v_2'(x) = x + 1.$$

Therefore $u(x) = c_1 e^{2x} + c_2 x e^{2x} - (\frac{1}{3}x^3 + \frac{1}{2}x^2)e^{2x} + (\frac{1}{2}x^2 + x)x e^{2x}$.

1.6 Solution via Power Series

General Case Consider a general second-order, linear, homogenous ODE

$$Lu = a_2(x)u'' + a_1(x)u' + a_0(x)u = 0.$$

Equivalently,

$$u'' + p(x)u' + q(x)u = 0,$$

where

$$p(x) = \frac{a_1(x)}{a_2(x)} \text{ and } q(x) = \frac{a_0(x)}{a_2(x)}.$$

Assume that a_j is **analytic** at 0 for $0 \leq j \leq 2$. Then p and q are analytic at 0, that is, they admit power series expansions

$$p(z) = \sum_{k=0}^{\infty} p_k z^k \text{ and } q(z) = \sum_{k=0}^{\infty} q_k z^k \text{ for } |z| < \rho,$$

for some $\rho > 0$.

Convergence Theorem If the coefficients $p(z)$ and $q(z)$ are analytic for $|z| < \rho$, then the formal power series for the solution $u(z)$, constructed above, is also analytic for $|z| < \rho$.

Power Series at Zero Consider

$$Lu = (1 - x^2)u'' - 5xu' - 4u = 0, \quad u(0) = 1, \quad u'(0) = 2.$$

In this case,

$$p(z) = \frac{-5z}{1 - z^2} = -5 \sum_{k=0}^{\infty} z^{2k+1} \text{ and } q(z) = \frac{-4}{1 - z^2} = -4 \sum_{k=0}^{\infty} z^{2k}$$

are analytic for $|z| < 1$, so the theorem guarantees that $u(z)$, given by the formal power series, is also analytic for $|z| < 1$.

Expansion about a Point other than Zero Suppose we want a power series expansion about a point $c \neq 0$, for instance because the initial conditions are given at $x = c$.

A simple change of the independent variable allows us to write

$$u = \sum_{k=0}^{\infty} A_k (z - c)^k = \sum_{k=0}^{\infty} A_k Z^k \text{ where } Z = z - c.$$

Since $du/dx = du/dZ$ and $d^2u/dz^2 = d^2u/dZ^2$, we obtain the translated equation

$$\frac{d^2u}{dZ^2} + p(Z + c) \frac{du}{dZ} + q(Z + c)u = 0.$$

Now compute that A_k using the series expansions of $p(Z + c)$ and $q(Z + c)$ in powers of Z .

1.7 Singular ODEs

In general, we do not want L to be singular on an interval for which we wish to solve $Lu = f$. However, some important applications lead to singular ODEs so we now address this case.

A second-order **Euler-Cauchy ODE** has the form

$$Lu = ax^2u'' + bxu' + cu = f(x),$$

where a, b and c are constants with $a \neq 0$. This ODE is singular at $x = 0$.
Noticing that

$$Lx^r = [ar(r-1) + br + c]x^r,$$

we see that $u = x^r$ is a solution of the homogenous equation ($f = 0$) iff

$$ar(r-1) + br + c = 0.$$

Factorisation Suppose $ar(r-1) + br + c = a(r-r_1)(r-r_2)$. If $r_1 \neq r_2$ then the general solution of the homogenous equation $Lu = 0$ is

$$u(x) = C_1x^{r_1} + C_2x^{r_2}, \quad x > 0.$$

Lemma If $r_1 = r_2$ then the general solution of the homogenous Euler-Cauchy equation $Lu = 0$ is

$$u(x) = C_1x^{r_1} + C_2x^{r_1} \ln x, \quad x > 0.$$

Euler-Cauchy Equations with Nonreal Indicial Roots Suppose that $r_{1,2} = \alpha \pm \beta i$ are the roots of the indicial equation

$$ar(r-1) + br + c = 0$$

associated to the Euler-Cauchy equation

$$at^2u'' + btu' + cu = 0.$$

Then the real-valued solutions can be derived as follows. First note that

$$t^{\alpha+\beta i} = t^\alpha t^{\beta i}$$

is a solution. Then notice that

$$t^{\beta i} = e^{\ln t^{\beta i}} = e^{i \ln t^\beta} = \cos(\ln(t^\beta)) + i \sin(\ln(t^\beta)).$$

So,

$$t^\alpha t^{\beta i} = t^\alpha e^{\ln t^{\beta i}} = t^\alpha e^{i \ln t^\beta} = t^\alpha (\cos(\ln(t^\beta)) + i \sin(\ln(t^\beta)))$$

is a solution. Finally, since each of the real part and the imaginary part is separately a (linear independent) solution, we see that the general solution in this case is (for $t > 0$)

$$u(t) = t^\alpha (c_1 \cos(\ln(t^\beta)) + i \sin(\ln(t^\beta))).$$

Example Consider $t^2u'' - tu' + 5u = 0$. Then the indicial equation is

$$r(r-1) - r + 5 = 0 \implies r = 1 \pm 2i.$$

So the general solution is,

$$u(t) = t(c_1 \cos \ln t^2 + c_2 \sin \ln t^2).$$

A number of important applications lead to ODEs that can be written in the **Frobenius normal form**

$$z^2u'' + zP(z)u' + Q(z)u = 0,$$

where $P(z)$ and $Q(z)$ are analytic at $z = 0$:

$$P(z) = \sum_{k=0}^{\infty} P_k z^k \text{ and } Q(z) = \sum_{k=0}^{\infty} Q_k z^k, \quad |z| < \rho.$$

Now consider $z^2 u'' + zP(z)u' + Q(z)u = 0$. Formal manipulations show that $Lu(z)$ equals

$$I(r)A_0 z^r + \sum_{k=1}^{\infty} \left(I(k+r)A_k + \sum_{j=0}^{k-1} [(j+r)P_{k-j} + Q_{k-j}] A_j \right) z^{k+r},$$

where $I(r)$ is the indicial polynomial $I(r) := r(r-1)P_0r + Q_0$, so we define $A_0(r) = 1$ and

$$A_k(r) = \frac{-1}{I(k+r)} \sum_{j=0}^{k-1} [(j+r)P_{k-j} + Q_{k-j}] A_j(r), \quad k \geq 1,$$

provided $I(k+r) \neq 0$ for all $k \geq 1$.

1.8 Bessel and Legendre Equations

1.8.1 Bessel Equations and Functions

The **Bessel equation with parameter ν** is

$$z^2 u'' + zu' + (z^2 - \nu^2)u = 0.$$

This ODE is in Frobenius normal form, with indicial polynomial $I(r) = (r+\nu)(r-\nu)$, and we seek a series solution

$$u(z) = \sum_{k=0}^{\infty} A_k z^{k+r}.$$

We assume $\operatorname{Re} \nu \geq 0$, so $r_1 = \nu$ and $r_2 = -\nu$.

With the normalisation

$$A_0 = \frac{1}{2^\nu \Gamma(1+\nu)}$$

the series solution is called the **Bessel function of order ν** and is denoted

$$J_\nu(z) = \frac{(z/2)^\nu}{\Gamma(1+\nu)} \left[1 - \frac{(z/2)^2}{1+\nu} + \frac{(z/2)^4}{2!(1+\nu)(2+\nu)} - \cdots \right].$$

From the functional equation $\Gamma(1+z) = z\Gamma(z)$ we see that

$$J_\nu(z) = \frac{(z/2)^\nu}{\Gamma(1+\nu)} - \frac{(z/2)^{\nu+2}}{\Gamma(2+\nu)} + \frac{(z/2)^{\nu+4}}{2!\Gamma(3+\nu)} - \frac{(z/2)^{\nu+6}}{3!\Gamma(4+\nu)} + \cdots$$

and so

$$J_\nu(z) = \sum_{k=0}^{\infty} \frac{(-1)^k (z/2)^{2k+\nu}}{k! \Gamma(k+1+\nu)}.$$

1.8.2 Legendre Equation

The **Legendre equation** with parameter ν is

$$(1 - z^2)u'' - 2zu' + \nu(\nu + 1)u = 0.$$

This ODE is not singular at $z = 0$ so the solution has an ordinary Taylor series expansion

$$u = \sum_{k=0}^{\infty} A_k z^k.$$

The A_k must satisfy

$$(k + 1)(k + 2)A_{k+2} - [k(k + 1) - \nu(\nu + 1)A_k] = 0$$

for $k \geq 0$, and since

$$k(k + 1) - \nu(\nu + 1) = (k - \nu)(k + \nu + 1),$$

the recurrence relation is

$$A_{k+1} = \frac{(k - \nu)(k + \nu + 1)}{(k + 1)(k + 2)} A_k \text{ for } k \geq 0.$$

We have

$$u(z) = A_0 u_0(z) + A_1 u_1(z)$$

where

$$u_0(z) = 1 - \frac{\nu(\nu + 1)}{2!} z^2 + \frac{(\nu - 2)\nu(\nu + 1)(\nu + 3)}{4!} z^4 - \dots$$

and

$$u_1(z) = z - \frac{(\nu - 1)(\nu + 2)}{3!} z^3 + \frac{(\nu - 3)(\nu - 1)(\nu + 2)(\nu + 4)}{5!} z^5 - \dots$$

Suppose now that $\nu = n$ is a non-negative integer. If n is even the series for $u_0(z)$ terminates, whereas if n is odd then the series for $u_1(z)$ terminates.

The terminating solution is called the **Legendre polynomial** of degree n and is denoted by $P_n(z)$ with the normalization

$$P_n(1) = 1.$$

Legendre Polynomials The first few Legendre polynomials are

$$\begin{aligned} P_0(z) &= 1, & P_3(z) &= \frac{1}{2}(5z^3 - 3z), \\ P_1(z) &= z, & P_4(z) &= \frac{1}{8}(35z^4 - 30z^2 + 3), \\ P_2(z) &= \frac{1}{2}(3z^2 - 1), & P_5(z) &= \frac{1}{8}(63z^5 - 70z^3 + 15z). \end{aligned}$$

Notice that P_n is an even or odd function according to whether n is even or odd.

Chapter 2

Dynamical Systems

2.1 Terminology

We begin with some examples of how systems of differential equations arise in applications, and see how all such problems can be formulated as a **first-order** system

$$\frac{d\mathbf{x}}{dt} = \mathbf{F}(\mathbf{x}).$$

Such a formulation leads to a natural geometric interpretation of a solution.

Lotka-Volterra Equations Simplified ecology with two species:

$$\begin{aligned} F(t) &= \text{number of foxes at time } t, \\ R(t) &= \text{number of rabbits at time } t. \end{aligned}$$

Assume populations large enough that F and R can be treated as smoothly varying in time. In the 1920s, Alfred Lotka and Vito Volterra independently proposed the predator-prey model

$$\begin{aligned} \frac{dF}{dt} &= -aF + \alpha FR, & F(0) &= F_0, \\ \frac{dR}{dt} &= bR - \beta FR, & R(0) &= R_0. \end{aligned}$$

Here a, α, b and β are non-negative constants.

Any first-order system for N ODEs in the form

$$\begin{aligned} \frac{dx}{dt} &= F_1(x, y, \dots, x_N), & x(0) &= x_{10}, \\ \frac{dy}{dt} &= F_2(x, y, \dots, x_N), & y(0) &= x_{20}, \\ &\vdots & &\vdots \\ \frac{dx_N}{dt} &= F_N(x, y, \dots, x_N), & x_N(0) &= x_{N0}, \end{aligned}$$

can be written in vector notation as

$$\frac{d\mathbf{x}}{dt} = \mathbf{F}(\mathbf{x}) \quad \mathbf{x}(0) = \mathbf{x}_0.$$

The system of ODEs is determined by the **vector field** $\mathbf{F} : \mathbb{R}^N \rightarrow \mathbb{R}^N$.

A system of ODEs of the form

$$\frac{d\mathbf{x}}{dt} = \mathbf{F}(\mathbf{x})$$

is said to be **autonomous**.

In a **non-autonomous** system, \mathbf{F} will depend explicitly on t :

$$\frac{d\mathbf{x}}{dt} = \mathbf{F}(\mathbf{x}, t).$$

It can be shown that it is sufficient (in principle) to develop theory for the autonomous case as a non-autonomous system can be converted into an autonomous system.

Second-order ODE Consider an initial-value problem for a general (possibly non-autonomous) second-order ODE

$$\frac{d^2x}{dt^2} = f\left(x, \frac{dx}{dt}, t\right), \text{ with } x = x_0 \text{ and } \frac{dx}{dt} = y_0 \text{ at } t = 0.$$

If $x = x(t)$ is a solution, and if we let $y = dx/dt$, then

$$\frac{dy}{dt} = \frac{d^2x}{dt^2} = f\left(x, \frac{dx}{dt}, t\right) = f(x, y, t),$$

that is, (x, y) is a solution of the first-order system

$$\begin{aligned} \frac{dx}{dt} &= y, & x(0) &= x_0, \\ \frac{dy}{dt} &= f(x, y, t) & y(0) &= y_0. \end{aligned}$$

2.2 Existence and Uniqueness

The most fundamental question about a dynamical system

$$\frac{d\mathbf{x}}{dt} = \mathbf{F}(\mathbf{x}, t)$$

is

For a given initial value \mathbf{x}_0 , does a solution $\mathbf{x}(t)$ satisfying $\mathbf{x}(0) = \mathbf{x}_0$ exist, and if so is this solution unique?

Answer is **yes**, whenever the vector field \mathbf{F} is **Lipschitz**.

The number L is a **Lipschitz constant** for a function $f : [a, b] \rightarrow \mathbb{R}$ if

$$|f(x) - f(y)| \leq L|x - y| \quad \text{for all } x, y \in [a, b].$$

Example Consider $f(x) = 2x^2 - x + 1$ for $0 \leq x \leq 1$. Since

$$\begin{aligned} f(x) - f(y) &= 2(x^2 - y^2) - (x - y) = 2(x + y)(x - y) - (x - y) \\ &= (2x + 2y - 1)(x - y) \end{aligned}$$

we have $|f(x) - f(y)| = |2x + 2y - 1||x - y|$ so a Lipschitz constant is

$$L = \max_{x,y \in [0,1]} |2x + 2y - 1| = 3.$$

We say that the function $f : [a, b] \rightarrow \mathbb{R}$ is Lipschitz if a Lipschitz constant for f exists.

Lipschitz Continuity If f is Lipschitz then f is (uniformly) continuous.

Continuous does not imply Lipschitz Consider the (uniformly) continuous function

$$f(x) = 3 + \sqrt{x} \text{ for } 0 \leq x \leq 4.$$

In this case, if $x, y \in (0, 4]$ then

$$f(x) - f(y) = \sqrt{x} - \sqrt{y} = \left(\sqrt{x} - \sqrt{y} \times \frac{\sqrt{x} + \sqrt{y}}{\sqrt{x} + \sqrt{y}} \right) = \frac{x - y}{\sqrt{x} + \sqrt{y}}$$

so if a Lipschitz constant L exists then

$$L \geq \frac{|f(x) - f(y)|}{|x - y|} = \frac{1}{\sqrt{x} + \sqrt{y}}$$

for arbitrarily small x and y , a contradiction.

A function $f : I \rightarrow \mathbb{R}$ is C^k if $f, f', f'', \dots, f^{(k)}$ all exist and are continuous on the interval I .

Theorem For any closed and bounded interval $I = [a, b]$, if f is C^1 on I then $L = \max_{x \in I} |f'(x)|$ is a Lipschitz constant for f on I .

A vector field $\mathbf{F} : S \subseteq \mathbb{R}^N$ is Lipschitz on $S \subseteq \mathbb{R}^N$ if

$$\|\mathbf{F}(\mathbf{x}) - \mathbf{F}(\mathbf{y})\| \leq L\|\mathbf{x} - \mathbf{y}\| \quad \text{for all } \mathbf{x}, \mathbf{y} \in S$$

Here,

$$\|\mathbf{x}\| = \left(\sum_{j=1}^N x_j^2 \right)^{\frac{1}{2}}$$

denotes the **Euclidean norm** of the vector $\mathbf{x} \in \mathbb{R}^N$.

We say that $\mathbf{F}(\mathbf{x}, t)$ is **Lipschitz in \mathbf{x}** if

$$\|\mathbf{F}(\mathbf{x}, t) - \mathbf{F}(\mathbf{y}, t)\| \leq L\|\mathbf{x} - \mathbf{y}\|.$$

Local Existence and Uniqueness Let $\mathbf{x}_0 \in \mathbb{R}^N$, fix $r > 0$ and $\tau > 0$, and put

$$S = \{(\mathbf{x}, t) \in \mathbb{R}^N \times \mathbb{R} : \|\mathbf{x} - \mathbf{x}_0\| \leq r \text{ and } |t| \leq \tau\}.$$

If $\mathbf{F}(\mathbf{x}, t)$ is Lipschitz in \mathbf{x} for $\mathbf{x}, t \in S$, and if

$$\|\mathbf{F}(\mathbf{x}, t)\| \leq M \quad \text{for } (\mathbf{x}, t) \in S,$$

then there exists a unique C^1 function $\mathbf{x}(t)$ satisfying

$$\frac{d\mathbf{x}}{dt} = \mathbf{F}(\mathbf{x}, t) \quad \text{for } |t| \leq \min\{r/M, \tau\}, \quad \text{with } \mathbf{x}(0) = \mathbf{x}_0.$$

2.3 Linear Dynamical Systems

Linear differential equations are generally much easier to solve than nonlinear ones. Fortunately, linear DEs suffice for describing many important applications.

We say that the $N \times N$, first order system of ODEs

$$\frac{d\mathbf{x}}{dt} = \mathbf{F}(\mathbf{x}, t)$$

is **linear** if the RHS has the form

$$\mathbf{F}(\mathbf{x}, t) = A(t)\mathbf{x} + \mathbf{b}(t)$$

for some $N \times N$ matrix-valued function $A(t) = [a_{ij}(t)]$ and a vector-valued function $\mathbf{b} = [b_i(t)]$.

The system is autonomous precisely when A and \mathbf{b} are constant.

Global Existence and Uniqueness If the elements of $A(t)$ and components of \mathbf{b} are continuous for $0 \leq t \leq T$, then the linear initial-value problem

$$\frac{d\mathbf{x}}{dt} = A(t)\mathbf{x} + \mathbf{b}(t) \quad \text{for } 0 \leq t \leq T, \quad \text{with } \mathbf{x}(0) = \mathbf{x}_0,$$

has a unique solution $\mathbf{x}(t)$ for $0 \leq t \leq T$.

We now investigate the special case when A is constant and $\mathbf{b}(t) = \mathbf{0}$:

$$\frac{d\mathbf{x}}{dt} = A\mathbf{x}$$

General Solution via Eigensystem If \mathbf{v} is a constant vector and $A\mathbf{v} = \lambda\mathbf{v}$, we define $\mathbf{x}(t) = e^{\lambda t}\mathbf{v}$. Then

$$\frac{d\mathbf{x}}{dt} = \lambda e^{\lambda t}\mathbf{v} = e^{\lambda t}(\lambda\mathbf{v}) = e^{\lambda t}(A\mathbf{v}) = A(e^{\lambda t}\mathbf{v}) = A\mathbf{x}$$

that is, \mathbf{x} is a solution of $d\mathbf{x}/dt = A\mathbf{x}$. If $A\mathbf{v}_j = \lambda_j\mathbf{v}_j$ for $1 \leq j \leq N$, then the linear combination

$$\mathbf{x}(t) = \sum_{j=1}^N c_j e^{\lambda_j t} \mathbf{v}_j$$

is also a solution because the ODE is linear and homogenous. Provided the \mathbf{v}_j are linearly independent, then the above equation is a **general solution** because given any $\mathbf{x}_0 \in \mathbb{R}^N$ there exist unique c_j such that

$$\mathbf{x}(0) = \sum_{j=1}^N c_j \mathbf{v}_j = \mathbf{x}_0.$$

Example Consider

$$\begin{aligned}\frac{dx}{dt} &= -5x + 2y, & x(0) &= 5, \\ \frac{dy}{dt} &= -6x + 3y & y(0) &= 7.\end{aligned}$$

Note that the initial value problem can be written in the vector form

$$\mathbf{x}'(t) = \mathbf{A}\mathbf{x}(t), \mathbf{x}(0) = \mathbf{x}_0,$$

$$\mathbf{A} = \begin{bmatrix} -5 & 2 \\ -6 & 3 \end{bmatrix} \quad \text{and } \mathbf{x}_0 := \begin{bmatrix} 5 \\ 7 \end{bmatrix}$$

Solving the system, using the eigenpair approach, we would need to find the eigenvectors and eigenvalues.

Characteristic equation is

$$0 = |\mathbf{A} - \lambda \mathbf{I}| = (-5 - \lambda)(3 - \lambda) + 12 \implies \lambda_1 := -3 \text{ and } \lambda_2 = 1.$$

Next we find the associated eigenvectors.

$$\lambda_1 = -3 : (\mathbf{A} + 3\mathbf{I})\mathbf{v} = \mathbf{0} \implies \mathbf{v}_1 := \begin{bmatrix} 1 \\ 1 \end{bmatrix} \quad \lambda_2 = 1 : (\mathbf{A} - \mathbf{I})\mathbf{v} = \mathbf{0} \implies \mathbf{v}_2 := \begin{bmatrix} 1 \\ 3 \end{bmatrix}$$

This means that a general solution of the system $\mathbf{x}' = \mathbf{A}\mathbf{x}$ is

$$\mathbf{x}(t) = c_1 e^{-3t} \begin{bmatrix} 1 \\ 1 \end{bmatrix} + c_2 e^t \begin{bmatrix} 1 \\ 3 \end{bmatrix}$$

Applying the initial value we can see that the unique solution is $x(t) = 4e^{-3t} + e^t$ and $y(t) = 4e^{-3t} + 3e^t$.

A square matrix $A \in \mathbb{C}^{N \times N}$ is **diagonalisable** if there exists a non-singular matrix $Q \in \mathbb{C}^{N \times N}$ such that $Q^{-1}AQ$ is diagonal.

Theorem A square matrix $A \in \mathbb{C}^{N \times N}$ is diagonalisable if and only if there exists a basis $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_N\}$ for \mathbb{C}^N consisting of eigenvectors of A . Indeed if,

$$A\mathbf{v}_j = \lambda_j \mathbf{v}_j \text{ for } j = 1, 2, \dots, N,$$

and we put $Q = [\mathbf{v}_1 \ \mathbf{v}_2 \ \dots \ \mathbf{v}_N]$ then $Q^{-1}AQ = \Lambda$ where

$$A = \begin{bmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_N \end{bmatrix}$$

Consider a diagonalisable matrix A . Since $Q^{-1}AQ = \Lambda$, it follows that A has an eigenvalue decomposition

$$A = Q\Lambda Q^{-1}.$$

In general, we see by induction on k that

$$A^k = Q\Lambda^k Q^{-1} \text{ for } k = 0, 1, 2, \dots$$

Example

$$A = \begin{bmatrix} -5 & 2 \\ -6 & 3 \end{bmatrix}$$

then

$$\Lambda = \begin{bmatrix} -3 & 0 \\ 0 & 1 \end{bmatrix}, \quad Q = \begin{bmatrix} 1 & 1 \\ 1 & 3 \end{bmatrix}, \quad Q^{-1} = \frac{1}{2} \begin{bmatrix} 3 & -1 \\ -1 & 1 \end{bmatrix}$$

so

$$A^k = Q\Lambda^k Q^{-1} = \frac{1}{2} \begin{bmatrix} (-1)^k \times 3^{k+1} - 1 & (-1)^{k+1} \times 3^k + 1 \\ (-1)^k \times 3^{k+1} - 3 & (-1)^{k+1} \times 3^k + 3 \end{bmatrix}.$$

For any polynomial

$$p(z) = c_0 + c_1 z + c_2 z^2 + \cdots + c_m z^m$$

and any square matrix A , we define

$$p(A) = c_0 I + c_1 A + c_2 A^2 + \cdots + c_m A^m.$$

When A is diagonalisable, $A^k = Q\Lambda^k Q^{-1}$ so

$$\begin{aligned} p(A) &= c_0 Q I Q^{-1} + c_1 Q \Lambda Q^{-1} + \cdots + c_m Q \Lambda^m Q^{-1} \\ &\vdots \\ &= Q p(\Lambda) Q^{-1} \end{aligned}$$

Lemma For any polynomial p and any diagonal matrix Λ ,

$$p(\Lambda) = \begin{bmatrix} p(\lambda_1) & & \\ & \ddots & \\ & & p(\lambda_N) \end{bmatrix}$$

Theorem If two polynomials p and q are equal on the spectrum of a diagonalisable matrix A , that is, if

$$p(\lambda_j) = q(\lambda_j) \text{ for } j = 1, 2, \dots, N,$$

then $p(A) = q(A)$.

Example Recall that

$$A = \begin{bmatrix} -5 & 2 \\ -6 & 3 \end{bmatrix}$$

has eigenvalues $\lambda_1 = -3$ and $\lambda_2 = 1$. Let

$$p(z) = z^2 - 4 \quad \text{and} \quad q(z) = -2z - 1,$$

and observe

$$p(-3) = 5 = q(-3) \text{ and } p(1) = -3 = q(1).$$

We find

$$p(A) = A^2 - 4I = \begin{bmatrix} 9 & -4 \\ 12 & -7 \end{bmatrix} = -2A - I = q(A).$$

Exponential of a Diagonalisable Matrix If $A = Q\Lambda Q^{-1}$ is diagonalisable, then

$$e^A = Qe^\Lambda Q^{-1} \text{ and } e^\Lambda = \begin{bmatrix} e^{\lambda_1} & & \\ & e^{\lambda_2} & \\ & & \ddots \\ & & & e^{\lambda_N} \end{bmatrix}$$

Nilpotent Matrix A matrix is nilpotent if there exists a positive integer k such that $\mathbf{A}^k = \mathbf{O}$, where \mathbf{O} denotes the zero matrix.

Example

$$\mathbf{A}^2 = \mathbf{A}\mathbf{A} = \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} = \mathbf{O}.$$

Therefore \mathbf{A} is nilpotent and, in particular,

$$e^t \mathbf{A} = \mathbf{I} + t\mathbf{A} = \begin{bmatrix} 1 & 0 \\ t & 1 \end{bmatrix}$$