

Abstract Algebra and Fundamental Analysis

Jeremy Le — UNSW MATH2701 24T3

Contents

1	Algebra (Geometry)	3
1.1	Transformations and Groups	3
1.2	Subgroups and the Group of Isometries	4
1.3	Reflections and Isometries	5
1.4	Translations and Rotations on \mathbb{R}^2	6
1.5	Classification of Plane Isometries	7
1.6	Similarities	8
1.7	Dilatations	9
1.8	Classification of Plane Similarities	9
1.9	Normal Subgroups	10
1.10	Collineations	10
1.11	Darboux's Theorem	10
1.12	Affine Transformations	11
1.13	The Real Projective Line $\mathbb{R}P^1$, Plane $\mathbb{R}P^2$ and Space $\mathbb{R}P^n$	12
1.14	The Principle of Duality in $\mathbb{R}P^2$	12
1.15	Desargues' Theorem and Pappus' Theorem	13
1.16	Projective Transformations in $\mathbb{R}P^n$	13
1.17	Projective Plane Transformations	14
2	Analysis	15
2.1	Asymptotics	15
2.2	Inequalities	16
2.2.1	Basic Inequalities	16
2.2.2	Role of Convexity	16
2.2.3	The Cauchy-Schwarz and Hölder Inequalities	16
2.3	Norms and Convex Bodies	17
2.3.1	p -norms	17
2.3.2	Convex Bodies	17
2.4	Duality	18
2.4.1	Dual Norms	18
2.4.2	Polar Bodies	18
2.4.3	Seperating Hyperplanes	18
2.4.4	Mahler Volume	18
2.5	Prime Numbers	19
2.5.1	Infinitude of Primes	19
2.5.2	Elementary Estimates for the Growth of $\pi(x)$	19
2.5.3	Statement of the Prime Number Theorem	19

2.6	The Real Numbers	19
2.7	Absolute Values and p -adic Numbers	20

1 Algebra (Geometry)

1.1 Transformations and Groups

Definition 1.1. A *transformation* on \mathbb{R}^n is a **bijection** from \mathbb{R}^n to \mathbb{R}^n . We will denote $\mathcal{B}(\mathbb{R}^n)$ the set of all transformations on \mathbb{R}^n .

In particular, a transformation on the Euclidean plane \mathbb{R}^2 is called a **plane transformation**.

Definition 1.2 (Group). A group is a set G equipped with a map

$$* : G \times G \rightarrow G, (g, h) \mapsto g * h = gh,$$

that satisfies the following axioms:

- (G1) **Associativity**, i.e. $g, h, k \in G$, then $(gh)k = g(hk)$.
- (G2) **Existence of identity**, i.e. there is an element denoted by e in G called the *identity* of G such that $eg = g = ge$ for any $g \in G$. (Such e is unique; notation: 1_G .)
- (G3) **Existence of inverse**, i.e. for any $g \in G$, there is an element denoted by $h \in G$ called the inverse of g such that $gh = hg = e$. (h is also unique; notation: g^{-1} .)

A group G is called commutative or abelian if $gh = hg$ for all $g, h \in G$.

Proposition 1.3. *Examples of Transformation Groups*

- (1) *The set $\mathcal{B}(\mathbb{R}^n)$ of all transformations on \mathbb{R}^n together with the operation of composition forms a group.*
- (2) *The set $\mathcal{T}(\mathbb{R}^n)$ of all translations on \mathbb{R}^n together with the operation of composition forms a group.*
- (3) *The set $\mathcal{C}(\mathbb{R}^n)$ of collineations of \mathbb{R}^n together with the operation of composition forms a group.*

Definition 1.4 (Subgroup). Let $(G, *)$ be a group. A nonempty subset $H \subseteq G$ is said to be a subgroup of G , denoted by $H \leq G$, if $(H, *)$ is a group.

Lemma 1.5 (Subgroup Lemma). *A nonempty subset H of a group G is a subgroup if and only if the following two closure conditions are satisfied:*

- (SG1) *Closure under multiplication, i.e. if $h, k \in H$, then $hk \in H$;*
- (SG2) *Closure under inverse, i.e. if $h \in H$, then $h^{-1} \in H$.*

In particular, $1_H = 1_G \in H$.

Definition 1.6 (Group Isomorphisms). For groups G, H , a map $f : G \rightarrow H$ is called a group homomorphism if $f(xy) = f(x)f(y)$ for all $x, y \in G$. A bijective group homomorphism is called an isomorphism. In this case, we say that G is isomorphic to H . Notation $G \cong H$.

1.2 Subgroups and the Group of Isometries

Lemma 2.1. *If S is a subset of a group $(G, *)$, then $\langle S \rangle = \bigcap_{S \subseteq H \leq G} H$. In other words, $\langle S \rangle$ is the **smallest** subgroup of G that contains all the elements of S .*

Definition 2.2. We call $\langle S \rangle$ the **subgroup of G generated by S** . A group generated by one element is called a **cyclic group**.

Notation:

- space: \mathbb{R}^n ;
- points: A, B, C, P, Q, R, \dots with position vectors $\mathbf{a}, \mathbf{b}, \mathbf{c}, \mathbf{p}, \mathbf{q}, \mathbf{r}, \dots$;
- transformations: $\tau, \pi, \sigma, \delta, \dots$;
- lines: l, m, n, \dots ; line equations in \mathbb{R}^n : $\mathbf{x} = \mathbf{a} + \lambda \mathbf{v}$ for all $\lambda \in \mathbb{R}$;
- planes in \mathbb{R}^n : $\mathbf{x} = \mathbf{a} + \lambda \mathbf{u} + \mu \mathbf{v}$ for all $\lambda, \mu \in \mathbb{R}$;
- **Hyperplanes** through $\mathbf{a} \in \mathbb{R}^n$ with normal $\mathbf{n} \in \mathbb{R}^n = \mathbf{0}$:

$$\mathbb{H}_{\mathbf{n}, \mathbf{a}} = \{\mathbf{x} \in \mathbb{R}^n \mid (\mathbf{x} - \mathbf{a}) \cdot \mathbf{n} = 0\} = \langle \mathbf{n} \rangle^\perp + \mathbf{a}.$$

- For points P, Q in \mathbb{R}^n , we may also define the **perpendicular bisector** of the line segment \overline{PQ} to be the hyperplane \mathbb{H} that passes through the midpoint of \overline{PQ} and perpendicular to \overline{PQ} . So \mathbb{H} has the equation $(\mathbf{x} - \mathbf{m}) \cdot (\mathbf{p} - \mathbf{q}) = 0$ where $\mathbf{m} = \frac{1}{2}(\mathbf{p} + \mathbf{q})$.
- It is clear that, for all $X \in \mathbb{H}$,

$$d(X, P) = \sqrt{\|\mathbf{x} - \mathbf{m}\|^2 + \|\mathbf{p} - \mathbf{m}\|^2} = \sqrt{\|\mathbf{x} - \mathbf{m}\|^2 + \|\mathbf{q} - \mathbf{m}\|^2} = d(X, Q).$$

The Euclidean space \mathbb{R}^n

- Length of a vector: $\|\mathbf{a}\| = \sqrt{\mathbf{a} \cdot \mathbf{a}}$;
- Distance between two points P, Q : $d(P, Q) := \|\mathbf{p} - \mathbf{q}\|$;
- Projection of \mathbf{a} on \mathbf{b} : $\text{proj}_{\mathbf{b}}(\mathbf{a}) = \frac{\mathbf{a} \cdot \mathbf{b}}{\mathbf{b} \cdot \mathbf{b}} \mathbf{b}$;
- Angle between \mathbf{a} and \mathbf{b} : $\cos(\theta) = \frac{\mathbf{a} \cdot \mathbf{b}}{\|\mathbf{a}\| \|\mathbf{b}\|}$;
- Orthogonality: $\mathbf{a} \perp \mathbf{b} \iff \mathbf{a} \cdot \mathbf{b} = 0$;

Definition 2.3. An *isometry* on \mathbb{R}^n is a map $\tau : \mathbb{R}^n \rightarrow \mathbb{R}^n$ which preserves distance between points: $d(P, Q) = d(\tau(P), \tau(Q))$, $\forall P, Q \in \mathbb{R}^n$.

Lemma 2.4. *The set of isometries which fix the zero vector is equal to the set of (linear) maps that represent multiplication by an orthogonal matrix.*

Theorem 2.5. *An isometry can be decomposed into a translation multiplied by a linear transformation, which can be represented by an orthogonal matrix. In other words, for every $\tau \in \mathcal{I}(\mathbb{R}^n)$, there exist an orthogonal $n \times n$ matrix Q and a vector $\mathbf{b} \in \mathbb{R}^n$ such that $\tau = T_{Q, \mathbf{b}} = T_{I, \mathbf{b}} \circ T_{Q, \mathbf{0}}$. In particular, an isometry is a **transformation**.*

Theorem 2.6. *The group of Isometries*

- (1) *The set $\mathcal{I}(\mathbb{R}^n)$ of all isometries forms a subgroup of the group $\mathcal{B}(\mathbb{R}^n)$ of all transformations.*
- (2) *The group $\mathcal{I} = \mathcal{I}(\mathbb{R}^n)$ contains two subgroups: the group \mathcal{T} of translations and the group \mathcal{O} of all orthogonal linear transformations. Moreover, we have $\mathcal{I} = \mathcal{T}\mathcal{O} := \{\tau\sigma \mid \tau \in \mathcal{T}, \sigma \in \mathcal{O}\}$.*

1.3 Reflections and Isometries

Definition 3.1. Let \mathbb{H} be a hyperplane. The reflection $\sigma_{\mathbb{H}}$ in \mathbb{H} is the mapping defined by:

$$\sigma_{\mathbb{H}}(P) = \begin{cases} P & \text{if } P \in \mathbb{H}; \\ P' & \text{if } P \text{ is off } \mathbb{H} \text{ and } \mathbb{H} \text{ is the perpendicular bisector of } P\bar{P}'. \end{cases}$$

(in the sense that $d(P, X) = d(P', X)$ for all $X \in \mathbb{H}$.)

Proposition 3.2. Let \mathbb{H} be a hyperplane.

- (1) A reflection $\sigma_{\mathbb{H}}$ is an isometry satisfying $\sigma_{\mathbb{H}}^2 = 1$.
- (2) $\sigma_{\mathbb{H}}$ fixes a line $m \not\subseteq \mathbb{H}$ if and only if $m \perp \mathbb{H}$.
- (3) $\sigma_{\mathbb{H}}$ fixes a line **pointwise** if and only if $m \subseteq \mathbb{H}$.

Theorem 3.3. If $\mathbb{H} = \mathbb{H}_{\mathbf{n}, \mathbf{a}}$, then there exist $Q = I - \frac{2}{\mathbf{n} \cdot \mathbf{n}} \mathbf{n} \mathbf{n}^T \in O_n(\mathbb{R})$ and $\mathbf{b} = 2 \frac{\mathbf{a} \cdot \mathbf{n}}{\mathbf{n} \cdot \mathbf{n}} \mathbf{n}$ such that

$$\sigma_{\mathbb{H}}(\mathbf{x}) = Q\mathbf{x} + \mathbf{b}.$$

Corollary 3.4. In \mathbb{R}^2 , if line ℓ has equation $aX + bY + c = 0$, then the reflection σ_{ℓ} in ℓ has equation:

$$\begin{aligned} \sigma_{\ell}(\mathbf{x}) &= \frac{1}{a^2 + b^2} \begin{bmatrix} b^2 - a^2 & -2ab \\ -2ab & a^2 - b^2 \end{bmatrix} \mathbf{x} + \frac{1}{a^2 + b^2} \begin{bmatrix} -2ac \\ -2bc \end{bmatrix} \\ &= \begin{pmatrix} x \\ y \end{pmatrix} - 2 \frac{(ax + by + c)}{a^2 + b^2} \begin{pmatrix} a \\ b \end{pmatrix}. \end{aligned}$$

Definition 3.5 (Points in Generic Position). We say that m points $P_1(\mathbf{p}_1), P_2(\mathbf{p}_2), \dots, P_m(\mathbf{p}_m)$ in \mathbb{R}^n are in **generic position** if the vectors $\mathbf{p}_i - \mathbf{p}_1$, for $i = 2, 3, \dots, m$, are linearly independent. In particular, $n + 1$ points in \mathbb{R}^n are in generic position if every hyperplane contains at most n of the $n + 1$ points.

Theorem 3.6. (1) An isometry on \mathbb{R}^n that fixes $n + 1$ points in generic position is the identity map.

(2) An isometry on \mathbb{R}^n that fixes n points in generic position is a reflection **or** the identity.

(3) An isometry that fixes $n - 1$ but not n points in generic position is a product of two **reflections**.

(4) Every isometry (in \mathbb{R}^n) is a product of **at most** $n + 1$ reflections.

Corollary 3.7. The group $\mathcal{I}(\mathbb{R}^n)$ is generated by reflections $\mathbb{H}_{\mathbf{n}, \mathbf{a}}$ for all $\mathbf{0} \neq \mathbf{n}, \mathbf{a} \in \mathbb{R}^n$.

Corollary 3.8. (1) A plane isometry that fixes three vertices of a triangle is the identity map.

(2) Every plane isometry $\tau \in \mathcal{I}(\mathbb{R}^2)$ is a product of at most three reflections in three lines.

1.4 Translations and Rotations on \mathbb{R}^2

Theorem 4.1. *An isometry τ in \mathbb{R}^n is a **translation** if and only if τ is the product of two reflections in parallel hyperplanes.*

Corollary 4.2. *A plane isometry is a translation if and only if it is a product of two reflections in parallel lines.*

Definition 4.3. A **rotation** on \mathbb{R}^2 about a point C , through angle θ , is the transformation that fixes C and otherwise sends a point P to a point P' , where $d(C, P) = d(C, P')$, and the angle from \vec{CP} to $\vec{CP'}$ is θ (in anti-clockwise direction) if $\theta > 0$, and clockwise if $\theta < 0$). We denote this transformation by $\rho_{C,\theta}$.

Theorem 4.4. *A plane isometry is a **rotation** if and only if it is the product of two reflections in intersecting lines. Further we have*

- (1) *if lines l, m intersect at C , and the directed angle from l to m is $\frac{\theta}{2} \in (-\frac{\pi}{2}, \frac{\pi}{2}]$, then $\sigma_m \sigma_l = \rho_{C,\theta}$;*
- (2) *if lines p, q, r are concurrent, then there exists a line l such that $\sigma_r \sigma_q \sigma_p = \sigma_l$.*

Corollary 4.5. (1) *A non-identity rotation (on \mathbb{R}^2) fixes exactly one point.*

(2) *A rotation with centre C fixes every circle with centre C .*

(3) *The set of all rotations about a particular point (i.e., with centre at a particular point) is a subgroup of the group $\mathcal{I}(\mathbb{R}^2)$ of isometries; further still, it is a **commutative** subgroup. In other words,*

$$\mathcal{R}_C := \{\rho_{C,\theta} : \theta \in \mathbb{R}\} \leq \mathcal{I}(\mathbb{R}^2) \text{ and } \rho \rho' = \rho' \rho, \forall \rho, \rho' \in \mathcal{R}_C.$$

Theorem 4.6 (Equation of a rotation). (1) *The rotation $\rho_{\mathbf{0},\theta} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ about the origin $\mathbf{0}$ and through angle θ is the linear isomorphism $T_{Q,\mathbf{0}}(\mathbf{x}) = Q\mathbf{x}$, where Q is the following matrix:*

$$Q = \begin{bmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{bmatrix}.$$

(2) *If \mathbf{c} is the position vector of C , then $\rho_{C,\theta} = T_{\mathbf{c}}(\rho_{\mathbf{0},\theta})T_{-\mathbf{c}}$. Hence, $\rho_{C,\theta}$ has the equation $\rho_{C,\theta}(\mathbf{x}) = Q\mathbf{x} + \mathbf{b}$, where Q defines $\rho_{\mathbf{0},\theta}$ as in (1) and $\mathbf{b} = (I - Q)\mathbf{c}$. At the group level, we have $\mathcal{R}_C = T_{\mathbf{c}}\mathcal{R}_0T_{-\mathbf{c}}$. Call the group \mathcal{R}_C is **conjugate** to the group \mathcal{R}_0 .*

Half-turn A rotation of the form $\rho_C := \rho_{C,\pi}$ is called a half-turn. A half-turn has the equation

$$\mathbf{x}' = -\mathbf{x} + 2\mathbf{c},$$

where \mathbf{c} is the position vector of C .

Definition 4.7. A figure $F_1 \subseteq \mathbb{R}^n$ is **congruent** to a figure $F_2 \subseteq \mathbb{R}^n$ if one can be mapped onto the other by an isometry; i.e. if there exists an isometry τ such that $\tau(F_1) = F_2$. **Notation:** $F_1 \cong F_2$ means F_1 is congruent to F_2 .

Theorem 4.8. *If $\triangle ABC \cong \triangle A'B'C'$ in \mathbb{R}^2 (same side lengths), then there exists a **unique** plane isometry τ such that*

$$\tau(A) = A', \tau(B) = B', \tau(C) = C'.$$

1.5 Classification of Plane Isometries

Definition 5.1. A plane isometry τ is called a **glide reflection** with axis c (a line) if there exist distinct lines a, b which are perpendicular to c such that $\tau = \sigma_c \sigma_b \sigma_a (= \sigma_b \sigma_a \sigma_c)$.

Proposition 5.2. (1) *A glide reflection is a composition of a reflection in line a and a halfturn centred at a point off a .*

(2) *A glide reflection is a translation followed by a reflection.*

(3) *A glide reflection fixes no points.*

(4) *A glide reflection fixes exactly one line, the axis, c .*

(5) *The midpoint of any point and its image under a glide reflection lies on its axis (c).*

Theorem 5.3. *Distinct lines p, q, r are neither concurrent, nor parallel, if and only if $\sigma_r \sigma_q \sigma_p$ is a glide reflection.*

Definition 5.4. An isometry that is a product of an even (resp., odd) number of reflections is said to be even (resp., odd) isometry.

Theorem 5.5. 1. *The set \mathcal{E} of even isometries in \mathbb{R}^n forms a subgroup of \mathcal{I} .*

2. *If \mathcal{E}' denotes the set of odd isometries, then $\mathcal{E} \cap \mathcal{E}' = \emptyset$.*

3. *If $\sigma = \sigma_H$ is a reflection, then $\mathcal{E}' = \sigma \mathcal{E} := \{\sigma \pi \mid \pi \in \mathcal{E}\}$.*

4. *We also have $\sigma \mathcal{E} = \mathcal{E} \sigma$ and $\mathcal{I} = \mathcal{E} \sqcup \sigma \mathcal{E}$.*

Corollary 5.6. *For any non-identity plane isometries, it is either even or odd. All even isometries are either translations or rotations. All odd isometries are reflections or glide reflections.*

Theorem 5.7. *A product of 4 reflections in \mathbb{R}^2 is a product of 2 reflections.*

Definition 5.8. Let $\Omega \subseteq \mathbb{R}^n$ be a geometric figure (or a subset). A **symmetry** of Ω is an isometry τ such that $\tau(\Omega) = \Omega$.

All the symmetries of Ω form a group $\text{sym}(\Omega)$, the **symmetry group** of Ω .

1.6 Similarities

Definition 6.1. A transformation $\alpha : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is called a **similarity of ratio** $r > 0$ if

$$d(\alpha(P), \alpha(Q)) = rd(P, Q), \text{ for all } P, Q \in \mathbb{R}^n.$$

Proposition 6.2. (1) *An isometry is a similarity of ratio 1.*

(2) *A similarity fixing two points is an isometry.*

(3) *A similarity fixing $n + 1$ points in generic position is the identity.*

(4) *The set of all similarities in \mathbb{R}^n forms a group, denote this set by \mathcal{S} or $\mathcal{S}(\mathbb{R}^n)$.*

Definition 6.3. A **stretch of ratio** $r > 0$ about point C is a transformation $\delta_{C,r}$ that fixes C and otherwise sends a point P to a point P' , where P' is the unique point on the **ray** from C through P such that $d(C, P') = r \cdot d(C, P)$.

Theorem 6.4. *Decomposition of a similarity If α is a similarity of ratio $r > 0$, and P is any **fixed** point, then $\alpha = \tau \delta_{P,r} = \delta_{P,r} \tau'$, for some isometries τ, τ' . Moreover, we have*

$$\mathcal{S} = \bigsqcup_{r>0} \mathcal{S} S_{P,r} = \bigsqcup_{r>0} S_{P,r} \mathcal{S} \text{ (disjoint unions),}$$

where $\mathcal{S} S_{P,r} = \{\tau S_{P,r} \mid \tau \in \mathcal{S}\}$ and $S_{P,r} \mathcal{S} = \{S_{P,r} \tau \mid \tau \in \mathcal{S}\}$.

Corollary 6.5. *A similarity is a **collineation** that preserves betweenness, midpoints, angles, perpendicularity, etc.*

Definition 6.6. (1) A **point reflection** about $C(\mathbf{c})$ is the isometry $\rho_C : \mathbb{R}^n \rightarrow \mathbb{R}^n$ defined by

$$\rho_C(\mathbf{x}) = -(\mathbf{x} - \mathbf{c}) + \mathbf{c} = -\mathbf{x} + 2\mathbf{c}.$$

(2) A **dilation** about the point C is a stretch transformation $\delta_{C,r}$ ($r > 0$) about C , or it is a stretch transformation followed by a point reflection both about C (i.e., $\rho_C \delta_{C,r}$).

Lemma 6.7. (1) *A point reflection is an isometry.*

(2) *The product of two point reflections is a translation.*

(3) *The product of a translation and a point reflection is a point reflection.*

Proposition 6.8. *All point reflections generate a subgroup \mathcal{H} (of \mathcal{S}). Moreover, \mathcal{H} is a (disjoint) union of the set \mathcal{T} of all translations and the set of all point reflections: for a fixed C ,*

$$\mathcal{H} = \mathcal{T} \sqcup \rho_C \mathcal{T} = \mathcal{T} \sqcup \mathcal{T} \rho_C = \mathcal{T} \sqcup \{\rho_P \mid P \in \mathbb{R}^n\}.$$

Proposition 6.9. *The dilation $\tau = \rho_C \delta_{C,r}$ ($r > 0$) has the following equation:*

$$\tau(\mathbf{x}) = (-r)\mathbf{x} + (1+r)\mathbf{c}.$$

Lemma 6.10. *Let $\mathbb{R}^\times = \{r \in \mathbb{R} \mid r \neq 0\}$. For any $r, s \in \mathbb{R}^\times$, and any point $P(\mathbf{p})$, we have*

$$(1) \delta_{P,-r} = \rho_O \delta_{P,r};$$

$$(2) \delta_{P,1} = 1, \delta_{P,-1} = \rho_P;$$

$$(3) \delta_{P,r} \delta_{P,s} = \delta_{P,rs};$$

$$(4) \delta_{P,r}^{-1} = \delta_{P,r^{-1}}.$$

Proposition 6.11. *The set $\{\delta_{C,r} \mid r \in \mathbb{R}^\times (:= \mathbb{R} - 0)\}$ forms a group that is isomorphic to the group $(\mathbb{R}^\times, \cdot)$.*

1.7 Dilatations

Definition 7.1. A collineation δ on \mathbb{R}^n is called a **dilatation** if, for every line ℓ in \mathbb{R}^n , $\ell \parallel \delta(\ell)$.

Proposition 7.2. The set $\mathcal{D}(\mathbb{R}^n)$ of all dilatations in \mathbb{R}^n forms a subgroup of $\mathcal{C}(\mathbb{R}^n)$.

Lemma 7.3. A dilatation that fixes two points is the identity map. Hence, for dilatations δ_1, δ_2 and distinct point A, B , if $\delta_1(A) = \delta_2(A)$ and $\delta_1(B) = \delta_2(B)$, then $\delta_1 = \delta_2$.

Lemma 7.4. (1) If A, B, C are collinear, distinct, with $\frac{CB}{CA} = r \neq 0$, then $\delta_{C,r}(A) = B$.

(2) For collinear points A, B, P, P' , if $\frac{AP}{PB} = \frac{AP'}{P'B}$, then $P = P'$.

(3) Let τ be a dilatation and let $\tau(P) = P'$ for every point P . If there exist points A, B such that \overrightarrow{AB} and $\overrightarrow{A'B'}$ have the same (resp., opposite) direction, then, for any points C, D , \overrightarrow{CD} and $\overrightarrow{C'D'}$ have the same (resp., opposite) direction.

Corollary 7.5. If points A, B, C are sent to A', B', C' under a dilatation, then

$$\frac{AB}{A'B'} = \frac{BC}{B'C'} = \frac{CA}{C'A'}.$$

Theorem 7.6. A dilatation is either a translation or a dilation. Hence, every dilatation is a similarity.

1.8 Classification of Plane Similarities

Definition 8.1. We say that figure $f_1 \subseteq \mathbb{R}^n$ and figure $f_2 \subseteq \mathbb{R}^n$ are **similar** if there is a similarity α such that $\alpha(f_1) = f_2$.

Theorem 8.2. If $\triangle ABC \sim \triangle A'B'C'$ in \mathbb{R}^2 , then there exists a **unique** plane similarity α such that

$$\alpha(A) = A', \alpha(B) = B', \alpha(C) = C'.$$

Theorem 8.3 (Equations of Similarities). If α is a similarity in \mathbb{R}^n , then there exist $Q \in O_n(\mathbb{R})$, $\mathbf{b} \in \mathbb{R}^n$ and $r \in \mathbb{R}_{>0}$ such that

$$\alpha(\mathbf{x}) = rQ\mathbf{x} + \mathbf{b}, \quad \text{for all } \mathbf{x} \in \mathbb{R}^n.$$

Lemma 8.4. A similarity without a fixed point is an isometry.

Definition 8.5. (1) A **stretch reflection** in \mathbb{R}^2 is a non-identity stretch about some point C followed by a reflection about a line through C .

(2) A **stretch rotation** in \mathbb{R}^2 is a non-identity stretch about some point C followed by a non-identity rotation about C .

Theorem 8.6. A non-identity plane similarity is exactly one of the following:

Isometry, Stretch of ratio $r \neq 1$, Stretch reflections, Stretch rotation.

Theorem 8.7. In the equation of similarities, the algebraic classification is as follows:

1. α is an isometry if $r = 1$;
2. α is a stretch (of ratio $r \neq 1$) if $r \neq 1$ and $Q = I$;
3. α is a stretch reflection if $r \neq 1, Q \neq I$ and $\det(Q) = -1$;
4. α is a stretch rotation if $r \neq 1, Q \neq I$ and $\det(Q) = 1$;

1.9 Normal Subgroups

Definition 9.1. A subgroup K of a group G is called a **normal subgroup** if $g^{-1}Kg \leq K$ (equivalently, $g^{-1}Kg = K$, or $gK = Kg$) for all $g \in G$. **Notation:** $K \trianglelefteq G$.

Theorem 9.2. Suppose $\alpha \in \mathcal{S}$ is a similarity, and $G \in \{\mathcal{I}, \mathcal{E}, \mathcal{D}, \mathcal{H}, \mathcal{T}\}$. Then $\alpha\tau\alpha^{-1} \in G$, for all $\tau \in G$. In other words, each of the groups $\mathcal{I}, \mathcal{E}, \mathcal{D}, \mathcal{H}, \mathcal{T}$ is a normal subgroup of \mathcal{S} .

Corollary 9.3. For $\alpha \in \mathcal{S}$, a point C and a hyperplane \mathbb{H} in \mathbb{R}^n , we have

$$\alpha\sigma_{\mathbb{H}}\alpha^{-1} = \sigma_{\alpha(\mathbb{H})}, \quad \alpha\rho_C\alpha^{-1} = \rho_{\alpha(C)}, \quad \alpha\delta_{C,r}\alpha^{-1} = \delta_{\alpha(C),r}.$$

In particular, in \mathbb{R}^2 , $\alpha\rho_{C,\theta}\alpha^{-1} = \rho_{\alpha(C),\pm\theta}$.

Proposition 9.4. 1. If $H \leq G$, then G is a disjoint union of **cosets** $gH, g \in G$.

2. If $K \trianglelefteq G$, then $G/K := \{gk \mid g \in G\}$ is a group with the subset multiplication. (G/K is called the **quotient group** of G by K).

1.10 Collineations

Theorem 10.1. A transformation is a collineation in \mathbb{R}^n if and only if the images of collinear points are themselves collinear.

Lemma 10.2. If α is a collineation in \mathbb{R}^n , and l, m are parallel lines, then $\alpha(l)$ and $\alpha(m)$ are parallel.

Theorem 10.3. A collineation takes the midpoint of points A, B to the midpoints of points $\alpha(A), \alpha(B)$.

Corollary 10.4. For a collineation α , if $n+1$ points P_0, P_1, \dots, P_n divide the segment $\overline{P_0P_n}$ into n congruent segments $\overline{P_{i-1}P_i}$, and $P'_i = \alpha(P_i)$, then the $n+1$ points P'_0, \dots, P'_n divide the segment $\overline{P'_0P'_n}$ into n congruent segments $\overline{P'_{i-1}P'_i}$.

In particular, if a point P is between A and B , and $\frac{AP}{PB} = r$ is **rational**, then $P' = \alpha(P)$ is between $\alpha(A)$ and $\alpha(B)$ and $\frac{A'P'}{P'B'} = r$.

1.11 Darboux's Theorem

Lemma 11.1. Let $t > 0, t \neq 1$, and P, Q be points on $\ell(A, B)$ such that $\frac{AP}{PB} = t, \frac{AQ}{QB} = -t$. Then C is the midpoint of P, Q if and only if $\frac{AC}{CB} = -t^2$.

Theorem 11.2. If α is a collineation, and point P is between points A, B then $\alpha(P)$ is between $\alpha(A), \alpha(B)$.

Corollary 11.3. A collineation on \mathbb{R}^n fixing two points on a line fixes the line pointwise.

1.12 Affine Transformations

Theorem 12.1. *A collineation in \mathbb{R}^n fixing $n + 1$ points in generic position is the identity.*

Definition 12.2. An **affine transformation** $\alpha : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is one that has an equation of the form $\alpha(\mathbf{x}) = A\mathbf{x} + \mathbf{b}$ for all $\mathbf{x} \in \mathbb{R}^n$, where $A \in GL_n(\mathbb{R})$, $\mathbf{b} \in \mathbb{R}^n$. (In other words, $\alpha = T_{A,\mathbf{b}}$.)

Lemma 12.3. *The set \mathcal{A} of all affine transformations in \mathbb{R}^n forms a group. Moreover, it contains the similarity group \mathcal{S} as a subgroup of \mathcal{A} .*

Theorem 12.4. *Let τ be a transformation. Then the following are equivalent:*

1. τ is an affine transformation;
2. τ is a collineation.

Proposition 12.5. *A (non-degenerate) conic section*

$$aX^2 + bXY + cY^2 + dX + eY + f = 0$$

*is affine equivalent to one of the following **affine standard form**:*

$$Y = X^2, \quad X^2 + Y^2 = 1, \quad XY = 1.$$

Definition 12.6. An affine transformation α with equation $\alpha(\mathbf{x}) = A\mathbf{x} + \mathbf{b}$ is called an **equi-affine transformation** if $\det(\alpha) := \det(A) = \pm 1$. An equi-affine transformation in \mathbb{R}^2 is called an **equiareal transformation**.

Proposition 12.7 (The group of equi-affine transformations). *The set \mathcal{Q} of all equi-affine transformations forms a subgroup of \mathcal{A} that has $\mathcal{Q}^+ = \{\alpha \in \mathcal{Q} \mid \det(\alpha) = 1\}$ as a normal subgroup.*

Theorem 12.8. (1) *Let α be an affine transformation in \mathbb{R}^2 with equation $\alpha(\mathbf{x}) = A\mathbf{x} + \mathbf{b}$ and let $\alpha(P) = P'$, etc., then $\text{area}(\triangle P'Q'R') = |\det A| \text{area}(\triangle PQR)$.*

(2) *If Ω is the parallelepiped spanned by the vectors $\mathbf{a}, \mathbf{b}, \mathbf{c}$ in \mathbb{R}^3 and α is an affine transformation in \mathbb{R}^3 , then $\text{vol}(\alpha\Omega) = |\det(A)| \text{vol}(\Omega)$.*

1.13 The Real Projective Line $\mathbb{R}P^1$, Plane $\mathbb{R}P^2$ and Space $\mathbb{R}P^n$

Definition 13.1. (1) The real projective plane $\mathbb{R}P^2$ is defined as the extended Euclidean plane

$$\mathbb{R}P^2 := \mathbb{R}^2 \sqcup \mathbb{R}P^1$$

The points in $b\mathbb{R}^2$ (resp., $\mathbb{R}P^1$) are called **ordinary** (resp., **ideal**) points and $\ell_\infty := \mathbb{R}P^1$ is called the **ideal (projective) line**.

(2) In general, for $n \geq 3$, define the n -dimensional real projective space

$$\mathbb{R}P^n = \underbrace{\mathbb{R}^n}_{\text{ordinary points}} \sqcup \underbrace{\mathbb{R}P^{n-1}}_{\text{ideal points}}$$

as a disjoint union of the **ordinary** part \mathbb{R}^n and the **ideal** part $\mathbb{R}P^{n-1}$

Proposition 13.2. *Two distinct **projective lines** have **exactly one** point of intersection.*

1.14 The Principle of Duality in $\mathbb{R}P^2$

Definition 14.1. • A projective point in $\mathbb{R}P^n$ is a 1-dimensional subspace of \mathbb{R}^{n+1} . For $P[x_0, x_1, \dots] \in \mathbb{R}P^n$, we also write $P = \langle \mathbf{x} \rangle$, the dimensional subspace spanned by \mathbf{x} which is the column vector $(x_0, x_1, \dots, x_n)^T$.

- A projective **line** in $b\mathbb{R}P^n$ is a 2-dimensional subspace of \mathbb{R}^{n+1} . If $P = \langle \mathbf{p} \rangle, Q = \langle \mathbf{q} \rangle$ are distinct projective points then $p\ell(P, Q) = \langle \mathbf{p}, \mathbf{q} \rangle$, the subspace spanned by \mathbf{p}, \mathbf{q} .
- A projective **plane** in $\mathbb{R}P^n$ is a 3-dimensional subspace of \mathbb{R}^{n+1} .
- A projective **hyperplane** in $b\mathbb{R}P^n$ is a n -dimensional subspace of $b\mathbb{R}^{n+1}$.
- A projective point $P = \langle \mathbf{x} \rangle$ lies on a projective line $h = \langle \mathbf{p}, \mathbf{q} \rangle$ if the one dimensional subspace $\langle \mathbf{x} \rangle$ is a **subspace** of the two dimensional subspace $\langle \mathbf{p}, \mathbf{q} \rangle$.
- The **Real Projective Plane** $\mathbb{R}P^2$ is the set of all projective points $\langle \mathbf{x} \rangle, \mathbf{x} \in \mathbb{R}^3 - \{\mathbf{0}\}$, and lines $\langle \mathbf{p}, \mathbf{q} \rangle$ with $\langle \mathbf{p} \rangle \neq \langle \mathbf{q} \rangle$, together with the above incidence structure.

Proposition 14.2. *In $b\mathbb{R}P^2$, any two projective points lie on exactly one projective line, and any two projective lines intersect in exactly one projective point.*

Lemma 14.3. *For subspaces U, V of \mathbb{R}^n , we have*

$$(U + V)^\perp = U^\perp \cap V^\perp \quad \text{and} \quad (U \cap V)^\perp = U^\perp + V^\perp.$$

Principle of Duality In $\mathbb{R}P^2$, any true statement involving points and straight lines remains true if the words “points” and “lines” are interchanged (i.e., $\langle \mathbf{x} \rangle \leftrightarrow \langle \mathbf{x} \rangle^\perp$). E.g.,

- Any two projective points **lie on** exactly one projective line.
- Any two projective lines **intersect in** exactly one projective point.

Lemma 14.4. *Projective points $\langle \mathbf{p} \rangle, \langle \mathbf{q} \rangle, \langle \mathbf{r} \rangle$ in $\mathbb{R}P^2$ are **collinear** if and only if projective lines $\langle \mathbf{p} \rangle^\perp, \langle \mathbf{q} \rangle^\perp, \langle \mathbf{r} \rangle^\perp$ are **concurrent**.*

1.15 Desargues' Theorem and Pappus' Theorem

Proposition 15.1. (1) Three distinct projective points $P = \langle \mathbf{p} \rangle, Q = \langle \mathbf{q} \rangle$, and $R = \langle \mathbf{r} \rangle$ in $\mathbb{R}P^n$, are collinear if and only if the vectors $\mathbf{p}, \mathbf{q}, \mathbf{r}$ are linearly dependent. Moreover, we may choose $\mathbf{p}, \mathbf{q}, \mathbf{r}$ to satisfy $\mathbf{p} = \mathbf{q} + \mathbf{r}$.

(2) Four distinct projective points $P = \langle \mathbf{p} \rangle, Q = \langle \mathbf{q} \rangle, R = \langle \mathbf{r} \rangle$ and $S = \langle \mathbf{s} \rangle$ in $\mathbb{R}P^n$, no three of which are collinear, are coplanar if and only if the vectors $\mathbf{p}, \mathbf{q}, \mathbf{r}, \mathbf{s}$ are linearly dependent. Moreover, we may choose $\mathbf{p}, \mathbf{q}, \mathbf{r}, \mathbf{s}$ to satisfy $\mathbf{p} = \mathbf{q} + \mathbf{r} + \mathbf{s}$.

Theorem 15.2 (Desargues' Theorem). Let A, B, C, A', B', C' be distinct points in $\mathbb{R}P^2$, such that the projective lines $pl(A, A'), pl(B, B'), pl(C, C')$ are **distinct** and **concurrent**. Then the projective points of intersections $C'' = pl(A, B) \cap pl(A', B'), A'' = pl(B, C) \cap pl(B', C'), B'' = pl(A, C) \cap pl(A', C')$ are **collinear**.

Dual Desargues' Theorem Let l, m, n, l', m', n' be distinct lines in $\mathbb{R}P^2$ such that their intersections $l \cap l', m \cap m', n \cap n'$ are distinct projective points, and collinear. Then the projective lines joining $l \cap m, l' \cap m'$, and $m \cap n, m' \cap n'$, and $n \cap l, n' \cap l'$ are concurrent.

Theorem 15.3 (Pappus' Theorem). Let A, B, C and A', B', C' be two pairs of collinear triples of distinct points in a projective plane. Then the three points $A'' = pl(B, C') \cap pl(B', C), B'' = pl(C, A') \cap pl(C', A)$ and $C'' = pl(A, B') \cap pl(A', B)$ are collinear.

Dual Pappus' Theorem Let l, m, n, l', m', n' be two pairs of concurrent projective lines in $\mathbb{R}P^2$. Then the projective lines $pl(m \cap n', m' \cap n), pl(n' \cap l, n \cap l'), pl(l \cap m', l' \cap m)$ are concurrent.

1.16 Projective Transformations in $\mathbb{R}P^n$

Definition 16.1. A map $\pi : \mathbb{R}P^n \rightarrow \mathbb{R}P^n$ is called a *projective transformation* if there exists an **invertible** matrix $A \in GL_{n+1}(\mathbb{R})$ such that $\pi \langle \mathbf{x} \rangle = \langle A\mathbf{x} \rangle$.

Proposition 16.2. (1) A **linear isomorphism** $T_A : \mathbb{R}^{n+1} \rightarrow \mathbb{R}$ induces a projective transformation $\pi_A : \mathbb{R}P^n \rightarrow \mathbb{R}P^n, \langle x \rangle \mapsto \langle A\mathbf{x} \rangle$.

(2) For $A, A' \in GL_{n+1}(\mathbb{R}), \pi_A = \pi_{A'} \iff A = \lambda A', \text{ for some } A \in \mathbb{R}^\times$.

Theorem 16.3. Let $\mathcal{P} = \mathcal{P}(\mathbb{R}P^n)$ be the set of all projective transformations on $\mathbb{R}P^n$. Then \mathcal{P} is a group, called the **group of projective transformations**.

Theorem 16.4. (1) The set $PGL_{n+1}(\mathbb{R})$ with coset multiplication forms a group, the **projective linear group**. This is the **quotient group** $GL_{n+1}(\mathbb{R})/\mathcal{K}_{n+1}$ with $\mathcal{K}_{n+1} := \mathbb{R}^\times I_{n+1} = \{\lambda I_{n+1} \mid \lambda \in \mathbb{R}^\times\}$.

(2) The map $\phi : PGL_{n+1}(\mathbb{R}) \rightarrow \mathcal{P}, [A] \mapsto \pi_A$ is a **bijection**, satisfying: $\phi([A][B]) = \phi([A])\phi([B])$. That is, the map ϕ is a group isomorphism.

Theorem 16.5. Every affine transformation $T_{A, \mathbf{b}}$ on \mathbb{R}^n can be **uniquely** extended to a projective transformation $\pi \left(\begin{smallmatrix} 1 & 0 \\ \mathbf{b} & A \end{smallmatrix} \right)$ on $\mathbb{R}P^n$ which stabilises the ideal part and the ordinary part and preserves multiplication and inverses. In group theory, terminology, \mathcal{A} is (isomorphic to) a subgroup of \mathcal{P} . That is, $\mathcal{A} \equiv \mathcal{A}' \leq \mathcal{P}$.

1.17 Projective Plane Transformations

Theorem 17.1. *Let $f = \{P, Q, R, S\}$ and P', Q', R', S' be two sets of four points, no three of which are collinear in $\mathbb{R}P^2$. Then there is a unique $\pi \in \mathcal{P}$ such that $\pi(P) = P', \pi(Q) = Q', \pi(S) = S'$.*

Proposition 17.2. *For two figures $f, g \subseteq \mathbb{R}^n$, if they are affine equivalent, then the images f, g in $\mathbb{R}P^n$ are projective equivalent.*

Theorem 17.3. *All non-degenerate conic sections are projective equivalent.*

Definition 17.4. A bijective map $\tau : \mathbb{R}P^2 \rightarrow \mathbb{R}P^2$ is called a **(projective) collineation** if τ takes collinear points to collinear points. (Equivalently, τ sends any projective line to a projective line.)

Lemma 17.5. *If τ is a collineation of $\mathbb{R}P^2$ and τ fixes points*

$$P_1 = [1, 0, 0], P_2 = [0, 1, 0], P_3 = [0, 0, 1], Q = [1, 1, 1],$$

*then τ is the **identity** map.*

Theorem 17.6. *A bijective map τ on $\mathbb{R}P^2$ is a projective collineation if and only if τ is a projective transformation.*

2 Analysis

2.1 Asymptotics

Definition 1.1 (Big-Oh). Let f, g be functions defined on an interval of the form (a, ∞) . We shall say that

$$f(x) = O(g(x)), \quad (\text{as } x \rightarrow \infty)$$

if there exists $M > 0$ and $x_0 > a$ such that for all $x > x_0$,

$$|f(x)| \leq M|g(x)|.$$

Definition 1.2 (Big-Oh at a). Let f, g be functions defined on an open interval containing a . We shall say that

$$f(x) = O(g(x)), \quad (\text{as } x \rightarrow a)$$

if there exists $M > 0$ and $\delta > 0$ such that if $|x - a| < \delta$,

$$|f(x)| \leq M|g(x)|.$$

Definition 1.3. We shall say that $f(x) = o(g(x))$ as $x \rightarrow \infty$ if for all $\epsilon > 0$, there exists $x_0 = x_0(\epsilon)$ such that if $x > x_0$, then $|f(x)| < \epsilon|g(x)|$. We say $f(x)$ is *little-oh* of $g(x)$.

Definition 1.4. We shall write that $f(x) = \theta(g(x))$ (as $x \rightarrow \infty$) if $f(x) = O(g(x))$ and $g(x) = O(f(x))$ (as $x \rightarrow \infty$). That is, there are non-zero constants M_1, M_2 and x_0 such that for all $x > x_0$

$$M_1|g(x)| \leq |f(x)| \leq M_2|g(x)|.$$

Definition 1.5. We shall say that $f(x) \sim g(x)$ as $x \rightarrow \infty$ if $\frac{f(x)}{g(x)} \rightarrow 1$ as $x \rightarrow \infty$.

2.2 Inequalities

2.2.1 Basic Inequalities

1. Triangle inequality: $|x + y| \leq |x| + |y|$.
2. The sum inequality: $\left| \sum_{k=1}^n x_k y_k \right| \leq \max\{|x_k|\} \sum_{k=1}^n |y_k|$.
3. The integral inequality: $\left| \int_a^b f(x)g(x) \, dx \right| \leq \max_{a \leq x \leq b} \{|f(x)|\} \int_a^b |g(x)| \, dx$.
4. The AM-GM inequality: if $x, y > 0$ then $(xy)^{1/2} \leq \frac{x+y}{2}$.
5. The Cauchy-Schwarz inequality: $\left| \sum_{k=1}^n x_k y_k \right| \leq \left(\sum_{k=1}^n x_k^2 \right)^{1/2} \left(\sum_{k=1}^n y_k^2 \right)^{1/2}$.

Theorem 2.1 (Generalized AM-GM inequality). *Suppose that x_1, x_2, \dots, x_n are positive. Then*

$$(x_1 x_2 \dots x_n)^{1/n} \leq \frac{1}{n} \sum_{k=1}^n x_k$$

(with equality if and only if all the x_k are equal).

2.2.2 Role of Convexity

Definition 2.2. A function $f : \mathbb{R} \rightarrow \mathbb{R}$ is **convex** if for any $x_1, x_2 \in \mathbb{R}$ and any $t \in [0, 1]$

$$f(tx_1 + (1-t)x_2) \leq tf(x_1) + (1-t)f(x_2).$$

Theorem 2.3 (Jensen's Inequality). *Suppose that $f : \mathbb{R} \rightarrow \mathbb{R}$ is convex, that $x_1, \dots, x_n \in \mathbb{R}$ and $a_1, \dots, a_n > 0$. Then*

$$f\left(\frac{\sum a_i x_i}{\sum a_j}\right) \leq \frac{\sum a_i f(x_i)}{\sum a_j}.$$

2.2.3 The Cauchy-Schwarz and Hölder Inequalities

Theorem 2.4 (Hölder Inequality). *Suppose that $1 < p, q < \infty$ with $\frac{1}{p} + \frac{1}{q} = 1$. Then for any numbers $x_1, \dots, x_n, y_1, \dots, y_n$,*

$$\left| \sum_{k=1}^n x_k y_k \right| \leq \left(\sum_{k=1}^n |x_k|^p \right)^{1/p} \left(\sum_{k=1}^n |y_k|^q \right)^{1/q}.$$

Theorem 2.5 (Hölder's inequality for integrals). *Suppose that $f, g : [a, b] \rightarrow \mathbb{R}$ (or to \mathbb{C}) are continuous, and that $1 < p, q < \infty$ with $\frac{1}{p} + \frac{1}{q} = 1$. Then*

$$\left| \int_a^b f(t)g(t) \, dt \right| \leq \int_a^b |f(t)g(t)| \, dt \leq \left(\int_a^b |f(t)|^p \, dt \right)^{1/p} \left(\int_a^b |g(t)|^q \, dt \right)^{1/q}.$$

2.3 Norms and Convex Bodies

2.3.1 p -norms

Definition 3.1. Let V be a vector space. A norm on V is a function $\mathbf{x} \mapsto \|\mathbf{x}\|$ from V to \mathbb{R} which satisfies

1. $\|\mathbf{x}\| \geq 0$ for all $\mathbf{x} \in V$ and $\|\mathbf{x}\| = 0$ if and only if $\mathbf{x} = 0$.
2. $\|\lambda\mathbf{x}\| = |\lambda|\|\mathbf{x}\|$ for all $\mathbf{x} \in V$ and scalar λ .
3. (Triangle Inequality): $\|\mathbf{x} + \mathbf{y}\| \leq \|\mathbf{x}\| + \|\mathbf{y}\|$ for all $\mathbf{x}, \mathbf{y} \in V$.

Definition 3.2. Let $1 \leq p < \infty$. For $\mathbf{x} = (x_1, \dots, x_n) \in \mathbb{R}^n$ define

$$\|\mathbf{x}\|_p = \left(\sum_{k=1}^n |x_k|^p \right)^{1/p}.$$

Theorem 3.3 (Minkowski's Inequality). *If $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$, then*

$$\|\mathbf{x} + \mathbf{y}\|_p \leq \|\mathbf{x}\|_p + \|\mathbf{y}\|_p.$$

2.3.2 Convex Bodies

Definition 3.4. A (nonempty) subset K of a vector space V is **convex** if for all $\mathbf{x}, \mathbf{y} \in K$ and all $\lambda \in [0, 1]$,

$$\lambda\mathbf{x} + (1 - \lambda)\mathbf{y} \in K.$$

That is, K contains all the line segments joining points in K .

Proposition 3.5. *Let $(V, \|\cdot\|)$ be a normed vector space. Then the set*

$$K = \{\mathbf{x} \in V : \|\mathbf{x}\| \leq 1\}$$

is convex.

Definition 3.6. Suppose that $\emptyset \neq K \subseteq \mathbb{R}^n$. Then

1. K is said to be **centrally symmetric** with respect to the origin if $\mathbf{x} \in K \iff -\mathbf{x} \in K$.
2. K is **closed** if its complement is open (\iff it contains its boundary \iff it contains all its limit points).
3. K is **bounded above and below** if there exist $0 < c \leq C < \infty$ such that $B_c \subseteq K \subseteq B_C$ where B_c and B_C are the closed Euclidean ($\|\cdot\|_2$) balls of radius c and C .

Definition 3.7. Suppose that $\emptyset \neq K \subseteq \mathbb{R}^n$. We say that K is a **convex body** if it is convex, centrally symmetric, closed and bounded above and below.

Theorem 3.8. *There is a one-to-one correspondence between convex bodies in \mathbb{R}^n and norms on \mathbb{R}^n .*

Lemma 3.9. *Let $\|\cdot\|$ be a norm on \mathbb{R}^n . Define $f : \mathbb{R}^n \rightarrow \mathbb{R}$ by $f(\mathbf{x}) = \|\mathbf{x}\|$. Then f is a continuous function from $(\mathbb{R}^n, \|\cdot\|_2)$ to \mathbb{R} . That is, if $\|\mathbf{x}_n - \mathbf{x}\|_2 \rightarrow 0$, then $f(\mathbf{x}_n) \rightarrow f(\mathbf{x})$ (ie $|f(\mathbf{x}_n) - f(\mathbf{x})| \rightarrow 0$).*

2.4 Duality

2.4.1 Dual Norms

Definition 4.1. Suppose that $\|\cdot\|$ is a norm on \mathbb{R}^n . Its **dual norm** is defined by

$$\|\mathbf{x}\|^* = \sup_{\mathbf{y} \in \mathbb{R}^n} \frac{|\mathbf{x} \cdot \mathbf{y}|}{\|\mathbf{y}\|} = \sup_{\|\mathbf{y}\|=1} |\mathbf{x} \cdot \mathbf{y}|.$$

Theorem 4.2. *The dual norm is a norm.*

2.4.2 Polar Bodies

Definition 4.3. Let K be a convex body in \mathbb{R}^n . The polar of K is the convex body associated to the dual norm to $\|\cdot\|_K$.

Theorem 4.4 (Polar Duality Theorem). *Let K be a convex body. Then $K^{\circ\circ} = K$.*

2.4.3 Separating Hyperplanes

Definition 4.5. A hyperplane $H_{\mathbf{u}} = \{\mathbf{y} \in \mathbb{R}^n : \mathbf{y} \cdot \mathbf{u} = 1\}$ is a separating hyperplane for K and \mathbf{p} if

1. $\mathbf{x} \cdot \mathbf{u} \leq 1$ for all $\mathbf{x} \in K$ (ie all of K is on the ‘low side’ of $H_{\mathbf{u}}$), and
2. $\mathbf{p} \cdot \mathbf{u} \geq 1$. (ie \mathbf{p} is on the high side of $H_{\mathbf{u}}$)

We say that $H_{\mathbf{u}}$ is a strongly separating if $\mathbf{p} \cdot \mathbf{u} > 1$.

Theorem 4.6 (Separating Hyperplane Theorem). *If K is a convex body and \mathbf{p} is a point not in K , then there exists a hyperplane that strongly separates them.*

2.4.4 Mahler Volume

Definition 4.7. The Mahler volume of a convex body K is defined as

$$M(K) = \text{vol}(K)\text{vol}(K^\circ).$$

Lemma 4.8. *Suppose that A is an invertible $n \times n$ matrix and that K is a convex body. Then AK is a convex body with polar $(A^T)^{-1}K^\circ$.*

Theorem 4.9. *Let $K \subseteq \mathbb{R}^n$ be a convex body and let $A \in M_n$ be invertible. Then*

- $M(K) = M(K^\circ)$.
- $M(AK) = M(K)$.

2.5 Prime Numbers

2.5.1 Infinitude of Primes

Theorem 5.1 (Fundamental Theorem of Arithmetic). *Every natural number n can be written uniquely, up to re-ordering of the factors, as a product of primes.*

Theorem 5.2 (Euclid). *There are ∞ -ly many primes. As n tends to infinity, we have*

$$\sum_{p \leq n, p \in \mathbb{P}} \frac{1}{p} \rightarrow \infty.$$

2.5.2 Elementary Estimates for the Growth of $\pi(x)$

Theorem 5.3 (Gauss). *For $x \geq 2$, we have $\pi(x) \geq \log \log x$.*

2.5.3 Statement of the Prime Number Theorem

Theorem 5.4. *There exists a constant $c > 0$, effectively computable such that for $x \geq 2$*

$$\pi(x) = \text{Li}(x) + O \left[x \exp \left(-c \sqrt{\log x} \right) \right],$$

where the implied constant is absolute.

2.6 The Real Numbers

Definition 6.1. A sequence $\{x_n\}$ of rationals is Cauchy if, for all integers j , there exists N_0 such that if $n, m \geq N_0$ then $|x_n - x_m| < 1/j$.

Definition 6.2. A **cut** is a subset r of \mathbb{Q} such that

1. r is nonempty,
2. $r \neq \mathbb{Q}$,
3. if $x, y \in \mathbb{Q}$, if $x < y$ and if $y \in r$, then $x \in r$ too,
4. for every $x \in r$ there exists $y \in r$ with $x < y$.

Definition 6.3. Suppose that \mathbb{F} is an ordered field and suppose that $\emptyset \neq S \subseteq \mathbb{F}$.

1. We say that $L \in \mathbb{F}$ is an **upper bound** for S if $x \leq L$ for all $x \in S$.
2. L is the **least upper bound** for S if L is an upper bound, and if L' is any other upper bound then $L \leq L'$. We call L the **supremum** of S , written $\sup S$.

Definition 6.4. An ordered field has the **least upper bound property** if every nonempty set which has an upper bound, has a least upper bound.

Theorem 6.5. 1. *There is an ordered field with the least upper bound property.*

2. *If \mathbb{F}_1 and \mathbb{F}_2 are ordered fields with the least upper bound property then there is an order preserving isomorphism between them. Informally, \mathbb{F}_1 and \mathbb{F}_2 are the same structures but with different names for the elements.*

2.7 Absolute Values and p -adic Numbers

Definition 7.1. Let \mathbb{F} be a field. A **multiplicative valuation** or **absolute value** on \mathbb{F} is a function $v : \mathbb{F} \rightarrow \mathbb{R}$ satisfying:

1. $v(x) \geq 0$ for all $x \in \mathbb{F}$,
2. $v(x) = 0$ if and only if $x = 0$,
3. $v(xy) = v(x)v(y)$, ($x, y \in \mathbb{F}$),
4. $v(x + y) \leq v(x) + v(y)$, ($x, y \in \mathbb{F}$).

Definition 7.2 (The p -adic valuation on \mathbb{Q}). Fix a prime p . Define $|0|_p = 0$. Any $0 \neq n \in \mathbb{Z}$ can be written as $n = p^a b$ where p doesn't divide b . Define $|n|_p = p^{-a}$. For $x = \frac{n}{m} \in \mathbb{Q}$, define $|x|_p = |n|_p / |m|_p$.

Theorem 7.3. For any prime p , $|\cdot|_p$ is a valuation.

Definition 7.4. Two valuations v and u on \mathbb{F} are **equivalent** if there is some $c > 0$ such that $v(x) = u(x)^c$ for all $x \in \mathbb{F}$.

Definition 7.5. The p -adic numbers \mathbb{Q}_p are the set of equivalence classes of $|\cdot|_p$ Cauchy sequences of rational numbers.

Proposition 7.6. Suppose that $\{x_n\}$ is a $|\cdot|_p$ Cauchy sequence. Then $\{|x_n|_p\}$ converges in \mathbb{R} .

Definition 7.7. The set of p -adic integers \mathbb{Z}_p is a unit disk around 0 in \mathbb{Q}_p . That is

$$\mathbb{Z}_p = \{\alpha \in \mathbb{Q}_p : |\alpha|_p \leq 1\}.$$

Theorem 7.8. \mathbb{Z}_p is a ring.

Theorem 7.9. Every p -adic integer is the limit of a sequence of non-negative integers.

Theorem 7.10. Every p -adic number $\alpha \in \mathbb{Q}_p$ has a unique p -adic expansion

$$\alpha = \sum_{k=-r}^{\infty} \alpha_k p^k$$

with $\alpha_k \in \mathbb{Z}$ and $0 \leq \alpha_k \leq p - 1$. Also, $\alpha \in \mathbb{Z}_p$ if and only if all the coefficients of negative powers of p are zero.

Theorem 7.11. A standard p -adic expansion $\alpha = \sum_{k=-r}^{\infty} \alpha_k p^k$ represents a rational number if and only if it is eventually periodic to the left.

Theorem 7.12. Every infinite sequence of p -adic integers has a convergent subsequence.