

Large-Scale Galactic Magnetic Fields

Jeremiah Lübke

Ruhr-Universität Bochum, July 2019

Table of Contents

- 1 MHD
- 2 Dynamos
- 3 Spatial Structures of Dynamo Solutions
- 4 Observing Magnetic Fields

Magnetohydrodynamics (MHD)

Plasma

continuity:

$$\partial_t \rho + \nabla \cdot (\rho \underline{u}) = 0$$

momentum:

$$\begin{aligned} \partial_t (\rho \underline{u}) + \nabla \cdot (\rho \underline{u} \underline{u}) \\ = \underline{j} \times \underline{B} - \nabla p \end{aligned}$$

adiabatic closure:

$$p = 2nkT$$

Magnetic Field

Ohm's Law:

$$\underline{E} + \underline{u} \times \underline{B} = \eta \underline{j}$$

Faraday's Law:

$$\begin{aligned} \partial_t \underline{B} &= -\nabla \times \underline{E} \\ &= \nabla \times (\underline{u} \times \underline{B}) + \eta \nabla^2 \underline{B} \end{aligned}$$

- Plasma *induces* Magnetic Field
- Magnetic Field *accelerates* Plasma

- Consider the equation governing the magnetic field:

$$\partial_t \underline{B} = \nabla \times (\underline{u} \times \underline{B}) + \eta \nabla^2 \underline{B} \quad (1)$$

- Consider the equation governing the magnetic field:

$$\partial_t \underline{B} = \nabla \times (\underline{u} \times \underline{B}) + \eta \nabla^2 \underline{B} \quad (1)$$

-
- Splitting up the involved quantities into regular average and random fluctuations:

$$X = \langle X \rangle + X' \quad \text{with} \quad \langle \langle X \rangle \rangle = \langle X \rangle, \quad \langle X' \rangle = 0$$

- Consider the equation governing the magnetic field:

$$\partial_t \underline{B} = \nabla \times (\underline{u} \times \underline{B}) + \eta \nabla^2 \underline{B} \quad (1)$$

-
- Splitting up the involved quantities into regular average and random fluctuations:

$$X = \langle X \rangle + X' \quad \text{with} \quad \langle \langle X \rangle \rangle = \langle X \rangle, \quad \langle X' \rangle = 0$$

- And working out (1):

$$\implies \partial_t \langle \underline{B} \rangle = \nabla \times (\langle \underline{u} \rangle \times \langle \underline{B} \rangle) + \eta \nabla^2 \langle \underline{B} \rangle + \underbrace{\nabla \times \langle \underline{u}' \times \underline{B}' \rangle}_{\equiv \mathcal{E}}$$

- Consider the equation governing the magnetic field:

$$\partial_t \underline{B} = \nabla \times (\underline{u} \times \underline{B}) + \eta \nabla^2 \underline{B} \quad (1)$$

-
- Splitting up the involved quantities into regular average and random fluctuations:

$$X = \langle X \rangle + X' \quad \text{with} \quad \langle \langle X \rangle \rangle = \langle X \rangle, \quad \langle X' \rangle = 0$$

- And working out (1):

$$\implies \partial_t \langle \underline{B} \rangle = \nabla \times (\langle \underline{u} \rangle \times \langle \underline{B} \rangle) + \eta \nabla^2 \langle \underline{B} \rangle + \underbrace{\nabla \times \langle \underline{u}' \times \underline{B}' \rangle}_{\equiv \mathcal{E}}$$

- The large-scale average is influenced by the small-scale fluctuations!

- Consider the equation governing the magnetic field:

$$\partial_t \underline{B} = \nabla \times (\underline{u} \times \underline{B}) + \eta \nabla^2 \underline{B} \quad (1)$$

-
- Working out (1) in cylindrical coordinates, with $\underline{u} = r \Omega \underline{e}_\phi$ and some galaxy-related assumptions, one finally arrives at:

- Consider the equation governing the magnetic field:

$$\partial_t \underline{B} = \nabla \times (\underline{u} \times \underline{B}) + \eta \nabla^2 \underline{B} \quad (1)$$

-
- Working out (1) in cylindrical coordinates, with $\underline{u} = r \Omega \underline{e}_\phi$ and some galaxy-related assumptions, one finally arrives at:

$$\mathcal{E} = 0 \implies \begin{cases} \partial_t B_r = \partial_{zz} B_r \\ \partial_t B_\phi = \mathcal{R}_\Omega B_r r \partial_r \Omega + \partial_{zz} B_\phi \end{cases} \quad (2)$$

- Consider the equation governing the magnetic field:

$$\partial_t \underline{B} = \nabla \times (\underline{u} \times \underline{B}) + \eta \nabla^2 \underline{B} \quad (1)$$

-
- Working out (1) in cylindrical coordinates, with $\underline{u} = r \Omega \underline{e}_\phi$ and some galaxy-related assumptions, one finally arrives at:

$$\mathcal{E} = 0 \implies \begin{cases} \partial_t B_r = \partial_{zz} B_r \\ \partial_t B_\phi = \mathcal{R}_\Omega B_r r \partial_r \Omega + \partial_{zz} B_\phi \end{cases} \quad (2)$$

- The magnetic field dilutes and the dynamo dies
(\rightarrow Antidynamo theorem)

- Consider the equation governing the magnetic field:

$$\partial_t \underline{B} = \nabla \times (\underline{u} \times \underline{B}) + \eta \nabla^2 \underline{B} \quad (1)$$

-
- Working out (1) in cylindrical coordinates, with $\underline{u} = r \Omega \underline{e}_\phi$ and some galaxy-related assumptions, one finally arrives at:

$$\mathcal{E} \approx \alpha \langle \underline{B} \rangle \implies \begin{cases} \partial_t B_r = -\mathcal{R}_\alpha \partial_z (\alpha B_\phi) + \partial_{zz} B_r \\ \partial_t B_\phi = \mathcal{R}_\Omega B_r r \partial_r \Omega + \partial_{zz} B_\phi \end{cases} \quad (3)$$

- Consider the equation governing the magnetic field:

$$\partial_t \underline{B} = \nabla \times (\underline{u} \times \underline{B}) + \eta \nabla^2 \underline{B} \quad (1)$$

-
- Working out (1) in cylindrical coordinates, with $\underline{u} = r \Omega \underline{e}_\phi$ and some galaxy-related assumptions, one finally arrives at:

$$\mathcal{E} \approx \alpha \langle \underline{B} \rangle \implies \begin{cases} \partial_t B_r = -\mathcal{R}_\alpha \partial_z (\alpha B_\phi) + \partial_{zz} B_r \\ \partial_t B_\phi = \mathcal{R}_\Omega B_r r \partial_r \Omega + \partial_{zz} B_\phi \end{cases} \quad (3)$$

- small-scale turbulence ($\alpha \neq 0$) sustains the magnetic field
($\rightarrow \alpha$ - and Ω -effect)

- Consider the equation governing the magnetic field:

$$\partial_t \underline{B} = \nabla \times (\underline{u} \times \underline{B}) + \eta \nabla^2 \underline{B} \quad (1)$$

-
- Working out (1) in cylindrical coordinates, with $\underline{u} = r \Omega \underline{e}_\phi$ and some galaxy-related assumptions, one finally arrives at:


$$\mathcal{E} \approx \alpha \langle \underline{B} \rangle \implies \begin{cases} \partial_t B_r = -\mathcal{R}_\alpha \partial_z (\alpha B_\phi) + \partial_{zz} B_r \\ \partial_t B_\phi = \mathcal{R}_\Omega B_r r \partial_r \Omega + \partial_{zz} B_\phi \end{cases} \quad (3)$$

- small-scale turbulence ($\alpha \neq 0$) sustains the magnetic field
($\rightarrow \alpha$ - and Ω -effect)


► Order out of Chaos

Spatial Structures of Dynamo Solutions


Dynamo Mode 0 (Axisymmetric Spiral)



Dynamo Mode 1 (Bisymmetric Spiral)




Dynamo Mode 2 (Quadraxisymmetric Spiral)



Observing Magnetic Fields

- Synchrotron radiation:
polarization \perp field orientation
- Polarization is changed when
passing through magnetized
plasma (*Faraday Rotation*):


$$\beta = \text{RM}(n_e, B_{\parallel}) \lambda^2$$



Observing Magnetic Fields

- Synchrotron radiation:
polarization \perp field orientation
- Polarization is changed when
passing through magnetized
plasma (*Faraday Rotation*):

$$\beta = \text{RM}(n_e, B_{\parallel}) \lambda^2$$



References



S. B. F. Dorch.

Magnetohydrodynamics.

in: *Scholarpedia*, revision #186287.



R. Beck.

Galactic Magnetic Fields.

in: *Scholarpedia*, revision #185129.



U. Klein, A. Fletcher.

Galactic and Intergalactic Magnetic Fields.

Springer Praxis Books, 2015.



R. Beck.

Magnetic Fields in Spiral Galaxies, Part I: Disks.

https://www.mpifr-bonn.mpg.de/1281929/Beck_Galaxies.pdf