

From Group Theory to Representation Theory

Jerome Queck

May to July 2026

Contents

1	Rudiments of Group Theory	2
1	Review	2
2	Automorphisms	21
3	Group Actions	44
2	The General Linear Group	63
4	Basic Structure	63
5	Parabolic Subgroups	86
6	The Special Linear Group	102
3	Local Structure	120
7	Sylow's Theorem	120
8	Finite p -groups	128
9	The Schur-Zassenhaus Theorem	131
4	Normal Structure	134
10	Composition Series	134
11	Solvable Groups	137
5	Semisimple Algebras	142
12	Modules and Representations	142
13	Wedderburn Theory	148
6	Group Representations	154
14	Characters	154
15	The Character Table	159
16	Induction	168
A	Algebraic Integers and Characters	173
17	Algebraic Integers and Characters	173

Chapter 1

Rudiments of Group Theory

1 Review

Group Axioms

A group consists of a non-empty set G and a binary operation on G such that the following axioms hold:

- $\forall g, h, k \in G, (gh)k = g(hk)$ (associativity)
- $\exists e \in G$ s.t. $eg = ge = g \forall g \in G$ (identity)
- $\forall g \in G, \exists g^{-1} \in G$ s.t. $gg^{-1} = g^{-1}g = e$ (inverses)

Properties of Groups

- If $xy = yx$, the commutator $[x, y] = xyx^{-1}y^{-1} = e$
- If $xy = yx \forall x, y \in G$, then G is an **abelian** group.
- A $x \in G$ is of finite order if $\exists n \in \mathbb{N}$ s.t. $x^n = e$
- $|G|$, the number of elements in G is defined to be the order of a finite group G . Every element of a finite group is of finite order
- There are infinite groups that has every element having a finite order, but there are also infinite groups in which the identity is the only element of finite order (torsion-free groups such as $\mathbb{Z}$)

Properties of Subgroups

- H is a **subgroup** of G , denoted $H \leq G$, if $H \subseteq G$ and H is a group under the same operation as G . In particular, $e \in H$ and $h^{-1} \in H \forall h \in H$
- Every subgroup of a finite group is finite, but an infinite group always has both finite and infinite subgroups
- Every subgroup of an abelian group is abelian, but a non-abelian group can have both abelian and non-abelian subgroups

Proposition 1.1. *If $H, K \leq G$, then $H \cap K \leq G$. More generally, $H_i \leq G \Rightarrow \bigcap H_i \leq G$*

Proof. Suppose $H_i \leq G \forall i$

Since each $H_i \leq G$, we have $e \in H_i \forall i$, so $e \in \bigcap H_i$ and $\bigcap H_i \neq \emptyset$. For $x, y \in \bigcap H_i$, we have $x, y \in H_i \forall i$, so by closure of each H_i , $xy \in H_i \forall i$, giving $xy \in \bigcap H_i$. Similarly, $x^{-1} \in H_i \forall i$, so $x^{-1} \in \bigcap H_i$.

$\therefore \bigcap H_i \leq G$ □

Theorem (Lagrange's Theorem). *Let G be a **finite** group, and let $H \leq G$. Then $|H| \mid |G|$.*

Proof. Let $x \in G$. Define $f_x: H \rightarrow xH$ by $f_x(h) = xh$. If $f_x(h_1) = f_x(h_2)$, then $xh_1 = xh_2$, and multiplying on the left by x^{-1} gives $h_1 = h_2$. Also, every element of xH has the form xh for some $h \in H$, so f_x is surjective. Thus f_x is a bijection, and $|xH| = |H|$.

We claim that any two sets of the form xH are either equal or disjoint. Suppose that $xH \cap yH \neq \emptyset$. Then $xh_1 = yh_2$ for some $h_1, h_2 \in H$, so $x = y(h_2h_1^{-1})$. Since $h_2h_1^{-1} \in H$, we have $x = yh$ for some $h \in H$, and so $xH = yhH = yH$. Thus the distinct sets of the form xH partition G .

Since G is finite, there are finitely many such sets, say r . Each has cardinality $|H|$, so $|G| = r|H|$.

$\therefore |H| \mid |G|$. $\square$

If $X \subseteq G$, $\langle X \rangle := \bigcap \{H \leq G \mid X \subseteq H\}$. By Proposition 1.1, $\langle X \rangle \leq G$, and it is the smallest subgroup of G containing X . We call $\langle X \rangle$ the **subgroup generated by X** , and we say that X **generates** $\langle X \rangle$.

Proposition 1.2. *Let $X \subseteq G$. Then $\langle X \rangle = \{e\} \cup \{x_1^{\epsilon_1} \cdots x_r^{\epsilon_r} \mid r \geq 1, x_i \in X, \epsilon_i = \pm 1, 1 \leq i \leq r\}$.*

Proof. Let $S = \{e\} \cup \{x_1^{\epsilon_1} \cdots x_r^{\epsilon_r} \mid r \geq 1, x_i \in X, \epsilon_i = \pm 1, 1 \leq i \leq r\}$.

We first show that $S \leq G$. Clearly $e \in S$. If $a = e$ or $b = e$, then $ab \in S$. Otherwise, if $a = x_1^{\epsilon_1} \cdots x_r^{\epsilon_r} \in S$ and $b = y_1^{\delta_1} \cdots y_s^{\delta_s} \in S$, then $ab = x_1^{\epsilon_1} \cdots x_r^{\epsilon_r} y_1^{\delta_1} \cdots y_s^{\delta_s} \in S$. Also $e^{-1} = e$, and if $a = x_1^{\epsilon_1} \cdots x_r^{\epsilon_r}$, then $a^{-1} = x_r^{-\epsilon_r} \cdots x_1^{-\epsilon_1} \in S$. Hence $S \leq G$.

Let $\mathcal{H}_X = \{H \leq G \mid X \subseteq H\}$. By Proposition 1.1, $\langle X \rangle = \bigcap_{H \in \mathcal{H}_X} H \leq G$. Since $X \subseteq S$, we have $S \in \mathcal{H}_X$, and so $\langle X \rangle = \bigcap_{H \in \mathcal{H}_X} H \subseteq S$.

Conversely, let $H \in \mathcal{H}_X$. Since $x_i \in X \subseteq H$, we have $x_i^{\epsilon_i} \in H$, so $x_1^{\epsilon_1} \cdots x_r^{\epsilon_r} \in H$. Also $e \in H$, hence $S \subseteq H$. Thus $S \subseteq H$ for all $H \in \mathcal{H}_X$, so $S \subseteq \bigcap_{H \in \mathcal{H}_X} H = \langle X \rangle$.

$\therefore S = \langle X \rangle$. $\square$

Cyclic Groups

- A group G is **cyclic** if $G = \langle g \rangle$ for some $g \in G$. By Proposition 1.2, $\langle g \rangle = \{g^n \mid n \in \mathbb{Z}\}$ s.t. $|\langle g \rangle| = n$ and $o(g) = n$
- All cyclic groups are abelian
- If g is not of finite order, then $\langle g \rangle$ is a torsion-free infinite abelian group, and $\langle g \rangle \cong \mathbb{Z}$
- If g is of finite order, then $\langle g \rangle \cong \mathbb{Z}/n\mathbb{Z}$
- If $g \in G$ and $o(g) = n \in \mathbb{Z}$, $\langle g \rangle \leq G$, so by Lagrange's Theorem, $n \mid |G|$. Thus, the order of an element of a finite group must divide the order of that group

If $X, Y \subseteq G$, $XY := \{xy \mid x \in X, y \in Y\} \subseteq G$. Also, $X^{-1} := \{x^{-1} \mid x \in X\} \subseteq G$. If $H \neq \emptyset$, then $H \leq G \iff HH = H$ and $H^{-1} = H$

Proposition 1.3. *Let $H, K \leq G$. Then $HK \leq G \iff HK = KH$*

Proof. ($\Rightarrow$) Suppose $HK \leq G$. Since HK is a subgroup, $(HK)^{-1} = HK$. Also $H^{-1} = H$ and $K^{-1} = K$, since $H, K \leq G$. Hence $(HK)^{-1} = K^{-1}H^{-1} = KH$. Therefore $HK = (HK)^{-1} = KH$.

($\Leftarrow$) Suppose $HK = KH$. Since $e \in H \cap K$, we have $e = ee \in HK$. Let $h_1k_1, h_2k_2 \in HK$. Since $k_1h_2 \in KH = HK$, write $k_1h_2 = h_3k_3$ for some $h_3 \in H, k_3 \in K$. Then $(h_1k_1)(h_2k_2) = h_1(k_1h_2)k_2 = h_1h_3k_3k_2 \in HK$. Also $(hk)^{-1} = k^{-1}h^{-1} \in KH = HK$ for all $h \in H, k \in K$.

$\therefore HK \leq G$. $\square$

Theorem 1.4. *Let $G = \langle g \rangle$ be a cyclic group of order n*

- For each $d \mid n$, there is exactly one subgroup of order d , $\langle g^{\frac{n}{d}} \rangle$
- If $d, e \mid n$, then $\langle g^{\frac{n}{d}} \rangle \cap \langle g^{\frac{n}{e}} \rangle = \langle g^{\frac{n}{\gcd(d, e)}} \rangle$
- If $d, e \mid n$, then $\langle g^{\frac{n}{d}} \rangle \langle g^{\frac{n}{e}} \rangle = \langle g^{\frac{n}{\text{lcm}(d, e)}} \rangle$

Proof. First note that if $m \mid n$, then $o(g^m) = \frac{n}{m}$. Indeed, $(g^m)^{\frac{n}{m}} = g^n = e$, and if $(g^m)^t = e$, then $n \mid mt$, so $\frac{n}{m} \mid t$.

Let $d \mid n$. Taking $m = \frac{n}{d}$, we get $o(g^{\frac{n}{d}}) = d$, so $\langle g^{\frac{n}{d}} \rangle$ is a subgroup of order d .

For uniqueness, let $H \leq G$ with $|H| = d$, and let $a = \min\{m \in \mathbb{N} \mid g^m \in H\}$. This set is non-empty since $g^n = e \in H$. If $g^b \in H$, write $b = qa + r$, $0 \leq r < a$. Then $g^r = g^b(g^a)^{-q} \in H$, so by minimality of

$a, r = 0$. Hence $a \mid b$, so $H \subseteq \langle g^a \rangle$. Since $g^a \in H$, $\langle g^a \rangle \subseteq H$, and hence $H = \langle g^a \rangle$. Applying the same division argument to $n = qa + r$ gives $a \mid n$, so $|H| = |\langle g^a \rangle| = \frac{n}{a} = d$. Thus $a = \frac{n}{d}$, and $H = \langle g^{\frac{n}{d}} \rangle$. Now let $d, e \mid n$. An element $g^b \in \langle g^{\frac{n}{d}} \rangle \cap \langle g^{\frac{n}{e}} \rangle$ iff $\frac{n}{d} \mid b$ and $\frac{n}{e} \mid b$, iff $\text{lcm}(\frac{n}{d}, \frac{n}{e}) = \frac{n}{\gcd(d, e)}$ divides b . Hence $\langle g^{\frac{n}{d}} \rangle \cap \langle g^{\frac{n}{e}} \rangle = \langle g^{\frac{n}{\gcd(d, e)}} \rangle$. Finally, $\langle g^{\frac{n}{d}} \rangle \langle g^{\frac{n}{e}} \rangle = \{g^{\frac{n}{d}r + \frac{n}{e}s} \mid r, s \in \mathbb{Z}\}$. Since $\frac{n}{d}\mathbb{Z} + \frac{n}{e}\mathbb{Z} = \gcd(\frac{n}{d}, \frac{n}{e})\mathbb{Z} = \frac{n}{\text{lcm}(d, e)}\mathbb{Z}$, we get $\langle g^{\frac{n}{d}} \rangle \langle g^{\frac{n}{e}} \rangle = \langle g^{\frac{n}{\text{lcm}(d, e)}} \rangle$.
 $\therefore$ all three assertions hold. $\square$

Cosets

- If $H \leq G$ and $x \in G$, xH is called a left coset of H in G and Hx is called a right coset of H in G .
- There is a bijective correspondence between left and right cosets of H in G , sending xH to its inverse $(xH)^{-1} = Hx^{-1}$.
- Any two cosets are equal or disjoint, with $xH = yH \iff y^{-1}x \in H$.
- A $x \in G$ lies in exactly one coset of H , xH .
- $\forall x \in G, \exists$ a bijective correspondence between H and xH .
- We define the index of H in G , $|G : H|$ to be the number of cosets of H in G .
- The cosets of H in G partition G into $|G : H|$ disjoint sets of cardinality $|H|$, and we have $|G| = |G : H||H|$ (Lagrange's Theorem).
- All subgroups of a finite group are of finite index, while subgroups of an infinite group may be of finite or infinite index.

Theorem 1.5. *Let $G = \langle g \rangle$ be an infinite cyclic group*

- *For each $d \in \mathbb{N}, \exists$ exactly one subgroup of G of index d , $\langle g^d \rangle$. Furthermore, every non-trivial subgroup of G is of finite index*
- *The intersection of the subgroups of indices d and e is the subgroup of index $\text{lcm}(d, e)$*
- *The product of subgroups of indices d and e is the subgroup of index $\gcd(d, e)$*

Proof. As in the proof of Theorem 1.4, let $1 \neq H \leq G$, and choose $a = \min\{m \in \mathbb{N} \mid g^m \in H\}$. This set is non-empty since H contains some $g^b \neq e$, and then either $b > 0$ or $g^{-b} \in H$ with $-b > 0$. If $g^c \in H$, write $c = qa + r$, $0 \leq r < a$. Then $g^r = g^c(g^a)^{-q} \in H$, so minimality of a gives $r = 0$. Hence $a \mid c$, so $H = \langle g^a \rangle$.

For $d \in \mathbb{N}$, the cosets of $\langle g^d \rangle$ are $g^i \langle g^d \rangle$, $0 \leq i < d$. Hence $|G : \langle g^d \rangle| = d$. By the previous paragraph, every non-trivial subgroup is $\langle g^a \rangle$ for some $a \in \mathbb{N}$, so it has finite index a , and $\langle g^d \rangle$ is the unique subgroup of index d .

If $d, e \in \mathbb{N}$, then $g^b \in \langle g^d \rangle \cap \langle g^e \rangle \iff d \mid b$ and $e \mid b \iff \text{lcm}(d, e) \mid b$. Thus $\langle g^d \rangle \cap \langle g^e \rangle = \langle g^{\text{lcm}(d, e)} \rangle$, which has index $\text{lcm}(d, e)$.

Since G is abelian, $\langle g^d \rangle \langle g^e \rangle \leq G$ by Proposition 1.3. Moreover $\langle g^d \rangle \langle g^e \rangle = \{g^{dr+es} \mid r, s \in \mathbb{Z}\} = \langle g^{\gcd(d, e)} \rangle$, since $d\mathbb{Z} + e\mathbb{Z} = \gcd(d, e)\mathbb{Z}$. Hence the product has index $\gcd(d, e)$.

$\therefore$ all three assertions hold. $\square$

Theorem 1.6. *If $K \leq H \leq G$, then $|G : K| = |G : H||H : K|$*

Proof. By the coset-counting identity in Lagrange's Theorem, the coefficient of $|H|$ in $|G|$ is $|G : H|$, and the coefficient of $|K|$ in $|H|$ is $|H : K|$. Hence $|G| = |G : H||H|$ and $|H| = |H : K||K|$.

Substituting, $|G| = |G : H||H : K||K|$. But applying the same identity directly to $K \leq G$ gives $|G| = |G : K||K|$.

$\therefore |G : K| = |G : H||H : K|$. $\square$

Let $H \leq G$, and let $\mathcal{I}$ be an indexing set that is in bijective correspondence with G/H . $T = \{t_i \mid i \in \mathcal{I}\} \subseteq G$ is said to be a (left) transversal for H (or a set of (left) coset representatives of H in G) if the sets $t_i H$ are precisely the distinct cosets of H in G .

Normal Subgroups

- N is a *normal subgroup* of G if $xN = Nx \forall x \in G$, or equivalently if $xNx^{-1} \subseteq N$ for all $x \in G$
- If G is abelian, then every subgroup of G is normal.
- If $\{e\}$ and G are the only normal subgroups of G , then we say that G is *simple*.
- If $H \trianglelefteq G$ and $K \trianglelefteq H$, then it is not necessarily true that $K \trianglelefteq G$. However, it is clearly true that if $K \trianglelefteq G$ and $K \leq H \leq G$, then $K \trianglelefteq H$.

Proposition 1.7. *Let $H, K \leq G$. If $K \trianglelefteq G$, then $HK \leq G$ and $H \cap K \trianglelefteq H$; if also $H \trianglelefteq G$, then $HK \trianglelefteq G$ and $H \cap K \trianglelefteq G$.*

Proof. Let $H, K \leq G$. Suppose that $K \trianglelefteq G$.

Then we have $gK = Kg \forall g \in G$. Since $H \leq G$, we have $hK = Kh \forall h \in H$.

Take $x \in HK$, so $x = hk$ for some $h \in H, k \in K$. Since $hK = Kh, \exists$ some $h' \in H$ such that $x = kh'$, so we have $HK \subseteq KH$. By a similar argument, we have $KH \subseteq HK$, so we conclude $HK = KH$. By Proposition 1.3, we have $HK \leq G$.

Take $y \in H \cap K$ and $h \in H$. Since $y \in H$, we have $hyh^{-1} \in H$. Since $y \in K$ and $K \trianglelefteq G$, we have $hKh^{-1} \subseteq K \Rightarrow hyh^{-1} \in K$. It follows that $hyh^{-1} \in H \cap K$, so $H \cap K \trianglelefteq H$.

Suppose also that $H \trianglelefteq G$. Take $hk \in HK$ and $g \in G$. Then $g(hk)g^{-1} = (ghg^{-1})(gkg^{-1})$, where $ghg^{-1} \in H$ since $H \trianglelefteq G$ and $gkg^{-1} \in K$ since $K \trianglelefteq G$. Hence $g(hk)g^{-1} \in HK$, so $gHKg^{-1} \subseteq HK \Rightarrow HK \trianglelefteq G$.

Take $y \in H \cap K$ and $g \in G$. Since $H \trianglelefteq G$, $gyg^{-1} \in H$, and since $K \trianglelefteq G$, $gyg^{-1} \in K$. So $gyg^{-1} \in H \cap K \Rightarrow H \cap K \trianglelefteq G$. $\square$

Proposition 1.8. *Any subgroup of index 2 is normal*

Proof. Let $H \leq G$ and suppose that $|G : H| = 2$

Then there are 2 left cosets, H and $G - H$

It follows that $x \in H \iff xH = H = Hx$, and $x \notin H \iff xH = G - H = Hx$

$\therefore H \trianglelefteq G$ $\square$

Theorem 1.9. *If $N \trianglelefteq G$, then G/N forms the quotient group under the operation $(xN)(yN) = (xy)N$*

Proof. First, we show that the operation is well-defined. Suppose $xN = x'N$ and $yN = y'N$. Then $x' = xn_1$ and $y' = yn_2$ for some $n_1, n_2 \in N$. Since $N \trianglelefteq G$, $y^{-1}n_1y \in N$, so $n_1y = y(y^{-1}n_1y)$. Hence $x'y' = xn_1yn_2 = xy(y^{-1}n_1y)n_2 \in xyN$, and so $x'y'N = xyN$. Thus $(xN)(yN) = (xy)N$ is well-defined. Associativity follows from associativity in G , since $((xN)(yN))(zN) = ((xy)z)N$ and $(xN)((yN)(zN)) = (x(yz))N$.

The identity is $N = eN$, since $N(xN) = xN = (xN)N$. Also, $(xN)(x^{-1}N) = N = (x^{-1}N)(xN)$, so $x^{-1}N$ is the inverse of xN .

$\therefore G/N$ is a group under the operation $(xN)(yN) = (xy)N$. $\square$

The identity element of G/N is N , and the inverse is $x^{-1}N$. If G is abelian, then G/N is also abelian

Conjugation

- The conjugate of an element x by g is defined as gxg^{-1}
- x and y are conjugate if $\exists g \in G$ s.t. $y = gxg^{-1}$
- No 2 distinct elements of an abelian group can be conjugate
- $N \trianglelefteq G \iff \forall n \in N, gng^{-1} \in N$

Permutations

- A permutation of a set X is a bijective set map from X to X . The set of permutations of X forms a group under composition
- If $X = \{1, \dots, n\}$, then this group is called the symmetric group, S_n
- S_n is a finite group of order $n!$

- $\rho \in S_n$ is called a r -cycle if $\exists$ distinct integers $1 \leq a_1, \dots, a_r \leq n$ s.t. $\rho(a_i) = a_{i+1} \forall 1 \leq i \leq r, \rho(a_r) = a_1, \rho(b) = b \forall 1 \leq b \leq n$ which is not equal to some a_i
- Two cycles are disjoint if there is no number that is moved by both cycles
- Every element of S_n can be written as a product of disjoint cycles
- The cycle structure refers to the partition of n consisting of the lengths of the cycles that appear in a disjoint cycle decomposition of ρ

Proposition 1.10. *Let $n \in \mathbb{N}$. Then two elements of S_n are conjugate iff they have the same cycle structure*

Proof. Let $\alpha, \beta \in S_n$.

($\Rightarrow$) Suppose that α and β are conjugate, so $\beta = \sigma\alpha\sigma^{-1}$ for some $\sigma \in S_n$. We show that $\sigma\alpha\sigma^{-1}$ is the permutation with the same cycle structure as α obtained by applying σ to the symbols in α . Let π be this permutation.

Suppose that α fixes some i , so $\sigma(i)$ resides in a 1-cycle of π , and π fixes $\sigma(i)$. Then $\sigma\alpha\sigma^{-1}(\sigma(i)) = \sigma\alpha(i) = \sigma(i)$, so $\sigma\alpha\sigma^{-1}$ fixes $\sigma(i)$ as well.

Suppose that α moves i , and let $\alpha(i) = j$. Take the complete factorization of α to be

$$\alpha = \gamma_1\gamma_2 \cdots (\cdots i j \cdots) \cdots \gamma_t$$

Take $\sigma(i) = k$ and $\sigma(j) = l$, so $\pi : k \mapsto l$. Since $\sigma\alpha\sigma^{-1} : k \mapsto i \mapsto j \mapsto l$, we have $\sigma\alpha\sigma^{-1}(k) = \pi(k)$. So π and $\sigma\alpha\sigma^{-1}$ agree on all symbols of the form $k = \sigma(i)$. Since σ is a surjection, it follows that $\pi = \sigma\alpha\sigma^{-1} = \beta$, so α and β have the same cycle structure.

($\Leftarrow$) Suppose that α and β have the same cycle structure. Place the complete factorization of α over that of β so that cycles of the same length correspond, and let $\gamma \in S_n$ be the function sending the top to the bottom. For example, if

$$\begin{aligned}\alpha &= \gamma_1\gamma_2 \cdots (\cdots i j \cdots) \cdots \gamma_t \\ \beta &= \delta_1\delta_2 \cdots (\cdots k l \cdots) \cdots \delta_t\end{aligned}$$

then $\gamma(i) = k, \gamma(j) = l$, etc. γ is a permutation, since every i between 1 and n occurs exactly once in a complete factorization. By the forward direction, $\gamma\alpha\gamma^{-1} = \beta$, so α and β are conjugate. $\square$

- A **transposition** in S_n is a 2-cycle
- Every element of S_n can be written as a (not necessarily disjoint) product of transpositions in many ways. However, any two such expressions use the same number, modulo 2, of transpositions
- A permutation is **even** (resp. **odd**) if it is a product of an even (resp. odd) number of transpositions; every permutation is exactly one of these. Since an r -cycle is a product of $r - 1$ transpositions, an r -cycle is even iff r is odd
- The subset of S_n consisting of all even permutations is a subgroup of index 2, hence is normal in S_n by Proposition 1.8; this is the **alternating group** of degree n , denoted A_n

Homomorphisms

- Let G and H be groups. A **homomorphism** is a map $\varphi : G \rightarrow H$ satisfying $\varphi(xy) = \varphi(x)\varphi(y)$ for all $x, y \in G$
- If φ is a homomorphism, then $\varphi(1) = 1$ and $\varphi(x^{-1}) = \varphi(x)^{-1}$ for any $x \in G$
- The **trivial homomorphism** from G to H sends every element of G to the identity of H
- φ is a **monomorphism** if injective, an **epimorphism** if surjective, and an **isomorphism** if bijective. If φ is an isomorphism, then so is $\varphi^{-1} : H \rightarrow G$
- A homomorphism $\varphi : G \rightarrow G$ is an **endomorphism** of G ; a bijective endomorphism is an **automorphism**
- If there is an isomorphism $\varphi : G \rightarrow H$, we say G and H are **isomorphic** and write $G \cong H$. Isomorphism is an equivalence relation on groups, so we can speak of the *isomorphism class* of a group.

Standard examples of homomorphisms and isomorphisms:

- Let $G = \langle g \rangle$ and $H = \langle h \rangle$ be cyclic groups of order n . The map $\varphi : G \rightarrow H$ by $\varphi(g^a) = h^a$ is an isomorphism. Hence any two finite cyclic groups of the same order are isomorphic.
- Let G be a group, $H \leq G$, $g \in G$. The **conjugate** of H by g is $gHg^{-1} = \{ghg^{-1} \mid h \in H\}$, and $gHg^{-1} \leq G$. The map $\varphi : H \rightarrow gHg^{-1}$ by $\varphi(h) = ghg^{-1}$ is an isomorphism, so conjugate subgroups are isomorphic. (However, isomorphic subgroups need not be conjugate)
- Let $X = \{x_1, \dots, x_n\}$ and let S_X be the group of permutations of X . The map $\varphi : S_n \rightarrow S_X$ by $\varphi(\rho)(x_i) = x_{\rho(i)}$ is an isomorphism
- Let G be a group, $N \trianglelefteq G$. The **natural map** $\eta : G \rightarrow G/N$ defined by $\eta(x) = xN$ is an epimorphism

Kernel and image:

- If $\varphi : G \rightarrow H$ is a homomorphism, the **kernel** of φ is $\ker \varphi = \{g \in G \mid \varphi(g) = 1\}$, and the **image** is $\operatorname{im} \varphi = \{\varphi(g) \mid g \in G\}$
- We use $\varphi(K) = \{\varphi(k) \mid k \in K\}$ for the image of $K \leq G$
- E.g., if $N \trianglelefteq G$ and $\eta : G \rightarrow G/N$ is the natural map, then $\ker \eta = N$ and $\eta(K) = KN/N$ for any $K \leq G$. ($\eta(K) = K/N$ if K contains N .)

Proposition 1.11. *Let G and H be groups, and let $\varphi : G \rightarrow H$ be a homomorphism. Then $\ker \varphi \trianglelefteq G$, and $\varphi(K) \leq H$ for any $K \leq G$.*

Proof. First, $\varphi(1) = 1$, so $1 \in \ker \varphi$. If $x, y \in \ker \varphi$, then $\varphi(xy^{-1}) = \varphi(x)\varphi(y)^{-1} = 1$, so $xy^{-1} \in \ker \varphi$. Hence $\ker \varphi \leq G$.

Let $x \in \ker \varphi$ and $g \in G$. Then $\varphi(gxg^{-1}) = \varphi(g)\varphi(x)\varphi(g^{-1}) = \varphi(g)\varphi(g)^{-1} = 1$. Thus $gxg^{-1} \in \ker \varphi$, so $\ker \varphi \trianglelefteq G$.

Now let $K \leq G$. Since $1 \in K$, we have $1 = \varphi(1) \in \varphi(K)$. If $a, b \in \varphi(K)$, then $a = \varphi(k_1)$ and $b = \varphi(k_2)$ for some $k_1, k_2 \in K$. Hence $ab^{-1} = \varphi(k_1)\varphi(k_2)^{-1} = \varphi(k_1k_2^{-1}) \in \varphi(K)$.

$\therefore \varphi(K) \leq H$. □

Theorem (Fundamental Theorem on Homomorphisms). *If G and H are groups and $\varphi : G \rightarrow H$ is a homomorphism, then there is an isomorphism $\psi : G/K \rightarrow \varphi(G)$ such that $\varphi = \psi \circ \eta$, where $K = \ker \varphi$ and $\eta : G \rightarrow G/K$ is the natural map; moreover, the map ψ is uniquely determined.*

Proof. Define $\psi : G/K \rightarrow \varphi(G)$ by $\psi(xK) = \varphi(x)$. This is well-defined: if $xK = yK$, then $y^{-1}x \in K$, so $1 = \varphi(y^{-1}x) = \varphi(y)^{-1}\varphi(x)$, and hence $\varphi(x) = \varphi(y)$.

The map ψ is a homomorphism, since $\psi((xK)(yK)) = \psi(xyK) = \varphi(xy) = \varphi(x)\varphi(y) = \psi(xK)\psi(yK)$.

It is surjective by definition of $\varphi(G)$: if $h \in \varphi(G)$, then $h = \varphi(x) = \psi(xK)$ for some $x \in G$. It is injective since $\psi(xK) = 1 \Rightarrow \varphi(x) = 1 \Rightarrow x \in K \Rightarrow xK = K$. Hence ψ is an isomorphism.

For all $x \in G$, $(\psi \circ \eta)(x) = \psi(xK) = \varphi(x)$, so $\varphi = \psi \circ \eta$.

Finally, suppose $\theta : G/K \rightarrow \varphi(G)$ also satisfies $\varphi = \theta \circ \eta$. For any $xK \in G/K$, $\theta(xK) = \theta(\eta(x)) = \varphi(x) = \psi(xK)$. Thus $\theta = \psi$, so ψ is uniquely determined. □

Theorem (First Isomorphism Theorem). *Let G be a group. If $N \trianglelefteq G$ and $H \leq G$, then $HN/N \cong H/(H \cap N)$.*

Proof. By Proposition 1.7, $HN \leq G$ and $H \cap N \trianglelefteq H$. Since $N \trianglelefteq G$ and $HN \leq G$, we also have $N \trianglelefteq HN$, so HN/N is a group.

Let $\eta : G \rightarrow G/N$ be the natural map $\eta(g) = gN$, and let $\varphi = \eta|_H : H \rightarrow G/N$. Then φ is a homomorphism, and $\ker \varphi = \{h \in H \mid hN = N\} = H \cap N$.

For the natural map, $\eta(K) = KN/N$ for any $K \leq G$. Taking $K = H$, we get $\varphi(H) = \eta(H) = HN/N$, so we may regard φ as a homomorphism $H \rightarrow HN/N$.

By the Fundamental Theorem on Homomorphisms, $H/\ker \varphi \cong \varphi(H)$. Thus $H/(H \cap N) \cong HN/N$.

$\therefore HN/N \cong H/(H \cap N)$. □

Theorem (Correspondence Theorem). *Let G and H be groups, and let $\varphi : G \rightarrow H$ be an epimorphism having kernel N . Then there is a bijective correspondence given by φ between the set of subgroups of G that contain N and the set of subgroups of H . If K is a subgroup of G containing N , then this correspondence sends K to $\varphi(K)$; if L is a subgroup of H , then the subgroup of G sent to L under this correspondence is $\varphi^{-1}(L) = \{x \in G \mid \varphi(x) \in L\}$. Moreover, if K_1 and K_2 are subgroups of G containing N , then:*

- $K_2 \leq K_1$ iff $\varphi(K_2) \leq \varphi(K_1)$, and in this case we have $|K_1 : K_2| = |\varphi(K_1) : \varphi(K_2)|$.
- $K_2 \trianglelefteq K_1$ iff $\varphi(K_2) \trianglelefteq \varphi(K_1)$, and in this case the map from K_1/K_2 to $\varphi(K_1)/\varphi(K_2)$ sending xK_2 to $\varphi(x)\varphi(K_2)$ is an isomorphism.

Proof. Let $\mathcal{S} = \{K \leq G \mid N \leq K\}$ and $\mathcal{T} = \{L \leq H\}$. For $K \in \mathcal{S}$, Proposition 1.11 gives $\varphi(K) \leq H$. If $L \in \mathcal{T}$, then $\varphi^{-1}(L) \leq G$ by the subgroup test, and $N \leq \varphi^{-1}(L)$, so $\varphi^{-1}(L) \in \mathcal{S}$.

Define $\Phi : \mathcal{S} \rightarrow \mathcal{T}$ by $\Phi(K) = \varphi(K)$, and define $\Psi : \mathcal{T} \rightarrow \mathcal{S}$ by $\Psi(L) = \varphi^{-1}(L)$. Since φ is surjective, $\Phi(\Psi(L)) = \varphi(\varphi^{-1}(L)) = L$. If $K \in \mathcal{S}$, then $K \subseteq \Psi(\Phi(K))$. Conversely, if $x \in \Psi(\Phi(K))$, then $\varphi(x) = \varphi(k)$ for some $k \in K$, so $k^{-1}x \in \ker \varphi = N \leq K$, and hence $x \in K$. Thus $\Psi(\Phi(K)) = K$. Hence Φ and Ψ are inverse bijections.

Let $K_1, K_2 \in \mathcal{S}$. If $K_2 \leq K_1$, then $\varphi(K_2) \leq \varphi(K_1)$. Conversely, if $\varphi(K_2) \leq \varphi(K_1)$, then $K_2 = \Psi(\Phi(K_2)) \leq \Psi(\Phi(K_1)) = K_1$.

Now suppose $K_2 \leq K_1$. Define $\alpha : K_1/K_2 \rightarrow \varphi(K_1)/\varphi(K_2)$ by $\alpha(xK_2) = \varphi(x)\varphi(K_2)$. If $xK_2 = yK_2$, then $y^{-1}x \in K_2$, so $\varphi(y)^{-1}\varphi(x) \in \varphi(K_2)$, and $\alpha(xK_2) = \alpha(yK_2)$. Thus α is well-defined. It is surjective by definition. If $\alpha(xK_2) = \alpha(yK_2)$, then $\varphi(y)^{-1}\varphi(x) = \varphi(k)$ for some $k \in K_2$. Hence $k^{-1}y^{-1}x \in N \leq K_2$, so $y^{-1}x \in K_2$, and $xK_2 = yK_2$. Thus α is bijective, so $|K_1 : K_2| = |\varphi(K_1) : \varphi(K_2)|$. If $K_2 \trianglelefteq K_1$, then for $x \in K_1, k \in K_2$, we have $\varphi(x)\varphi(k)\varphi(x)^{-1} = \varphi(xkx^{-1}) \in \varphi(K_2)$, so $\varphi(K_2) \trianglelefteq \varphi(K_1)$. Conversely, suppose $\varphi(K_2) \trianglelefteq \varphi(K_1)$. If $x \in K_1, k \in K_2$, then $\varphi(xkx^{-1}) \in \varphi(K_2)$, so $xkx^{-1} \in \Psi(\Phi(K_2)) = K_2$. Hence $K_2 \trianglelefteq K_1$.

In this case, α is a homomorphism, since $\alpha((xK_2)(yK_2)) = \alpha(xyK_2) = \varphi(xy)\varphi(K_2) = \varphi(x)\varphi(y)\varphi(K_2) = \alpha(xK_2)\alpha(yK_2)$. Since α is already bijective, it is an isomorphism.

$\therefore$ the correspondence has all the stated properties. $\square$

As a special case of the Correspondence Theorem, if G is a group and $N \trianglelefteq G$, then every subgroup of G/N has the form K/N for some subgroup $K \leq G$ with $N \leq K$. Here we take φ to be the natural map $G \rightarrow G/N$.

Theorem (Second Isomorphism Theorem). *Let H and K be normal subgroups of a group G . If H contains K , then $G/H \cong (G/K)/(H/K)$.*

Proof. Let $\eta : G \rightarrow G/K$ be the natural map. Then $\ker \eta = K$. Since $K \leq H \leq G$, the Correspondence Theorem applies to $H \leq G$.

Since $H \trianglelefteq G$, applying the normality and quotient-isomorphism parts with $K_1 = G$ and $K_2 = H$ gives $\eta(H) \trianglelefteq \eta(G)$ and $G/H \cong \eta(G)/\eta(H)$.

Now $\eta(G) = G/K$ and $\eta(H) = H/K$. Hence $G/H \cong (G/K)/(H/K)$.

$\therefore G/H \cong (G/K)/(H/K)$. $\square$

Exercises

Exercise 1.1. Prove, or complete the sketched proof of, each result in this section.

Solution. The sketched proofs are completed above in:

- Proposition 1.1, Lagrange's Theorem, Proposition 1.2, and Proposition 1.3
- Theorem 1.4, Theorem 1.5, and Theorem 1.6
- Proposition 1.7, Proposition 1.8, Theorem 1.9, Proposition 1.10, and Proposition 1.11
- Fundamental Theorem on Homomorphisms, First Isomorphism Theorem, Correspondence Theorem, and Second Isomorphism Theorem

$\square$

Exercise 1.2. We say that a group G has *exponent* e if e is the smallest positive integer such that $x^e = 1$ for every $x \in G$. Show that if G has exponent 2, then G is abelian. For what integers e is a group having exponent e necessarily abelian?

Solution. First suppose that G has exponent 2. Let $x, y \in G$. Then $x^2 = y^2 = (xy)^2 = 1$, so $xyxy = 1$. Multiplying on the left by x and on the right by y , and using $x^2 = y^2 = 1$, gives $yx = xy$. Since x, y were arbitrary, G is abelian.

If $e = 1$, then $x = 1$ for all $x \in G$, so $G = \{1\}$, which is abelian. We claim that these are the only possibilities.

Let $e \geq 3$. If e is even, take the dihedral group

$$D_e = \langle r, s \mid r^e = s^2 = 1, srs^{-1} = r^{-1} \rangle.$$

This group is non-abelian, since $srs^{-1} = r^{-1} \neq r$. Its rotations have order dividing e , and its reflections have order 2. Since e is even, the exponent is $\text{lcm}(e, 2) = e$.

Now suppose that $e \geq 3$ is odd. Let

$$U_e = \left\{ M(a, b, c) = \begin{pmatrix} 1 & a & c \\ 0 & 1 & b \\ 0 & 0 & 1 \end{pmatrix} \mid a, b, c \in \mathbb{Z}/e\mathbb{Z} \right\},$$

under matrix multiplication, with entries computed in $\mathbb{Z}/e\mathbb{Z}$. Direct multiplication gives

$$M(a, b, c)M(a', b', c') = M(a + a', b + b', c + c' + ab').$$

Hence U_e is closed under multiplication. The identity is $M(0, 0, 0)$, and

$$M(a, b, c)^{-1} = M(-a, -b, ab - c).$$

Associativity follows from associativity of matrix multiplication, so U_e is a group. It is non-abelian, since

$$M(1, 0, 0)M(0, 1, 0) = M(1, 1, 1), \quad M(0, 1, 0)M(1, 0, 0) = M(1, 1, 0).$$

For $n \in \mathbb{N}$, induction using the multiplication formula gives

$$M(a, b, c)^n = M\left(na, nb, nc + \binom{n}{2}ab\right).$$

Taking $n = e$, we get

$$M(a, b, c)^e = M\left(0, 0, \binom{e}{2}ab\right).$$

Since e is odd, $\binom{e}{2} = e(e-1)/2$ is divisible by e , so $M(a, b, c)^e = M(0, 0, 0)$ for every $M(a, b, c) \in U_e$. Thus every element has order dividing e . However $M(1, 0, 0)^n = M(n, 0, 0)$, so $M(1, 0, 0)$ has order e . Hence U_e is a non-abelian group of exponent e .

$\therefore$ a group of exponent e is necessarily abelian iff $e = 1$ or $e = 2$. $\square$

Exercise 1.3. Let G be a finite group, and suppose that the map $\varphi : G \rightarrow G$ defined by $\varphi(x) = x^3$ for $x \in G$ is a homomorphism. Show that if 3 does not divide $|G|$, then G must be abelian. (See [2] for a generalization.)

Solution. Let $x, y \in G$. Since φ is a homomorphism, we have

$$x^3y^3 = \varphi(x)\varphi(y) = \varphi(xy) = (xy)^3 = xyxyxy.$$

Cancelling x on the left and y on the right gives

$$(yx)^2 = x^2y^2.$$

Swapping x and y , we also have

$$(xy)^2 = y^2x^2.$$

Using this in the original equation,

$$x^3y^3 = (xy)^3 = (xy)(xy)^2 = xy(y^2x^2) = xy^3x^2.$$

Cancelling x on the left gives

$$y^3x^2 = x^2y^3.$$

Hence every cube commutes with every square.

Now let $z \in \ker \varphi$. Then $z^3 = 1$, so $o(z) \mid 3$. Then we also have $o(z) \mid |G|$. Since $3 \nmid |G|$, we must have $o(z) = 1$, and hence $z = 1$. Thus $\ker \varphi = \{1\}$, so φ is injective. Since G is finite, φ is also surjective. Therefore every element of G is a cube.

Let $g \in G$. Since every element of G is a cube, write $g = y^3$ for some $y \in G$. Then for every $x \in G$, the relation above gives

$$gx^2 = y^3x^2 = x^2y^3 = x^2g.$$

Thus every square is central. In particular, x^2 and y^2 commute for all $x, y \in G$. Since $(xy)^2 = y^2x^2$, it follows that

$$(xy)^2 = x^2y^2.$$

Hence $xyxy = xxyy$. Cancelling x on the left and y on the right gives $yx = xy$. Since x, y were arbitrary, G is abelian. $\square$

Exercise 1.4. Let g be an element of a group G , and suppose that $|G| = mn$ where m and n are coprime. Show that there are unique elements x and y of G such that $xy = g = yx$ and $x^m = 1 = y^n$. (In the case where m is a power of some prime p , we call x the p -part of g and y the p' -part of g ; more generally, if π is a set of primes which includes all prime divisors of m and no prime divisors of n , then x and y are called the π -part and π' -part, respectively, of g .)

Solution. Since $\gcd(m, n) = 1$, choose $a, b \in \mathbb{Z}$ such that $am + bn = 1$. Since $\langle g \rangle \leq G$, Lagrange's Theorem gives $o(g) = |\langle g \rangle| \mid |G| = mn$, so $g^{mn} = 1$. Define

$$x = g^{bn}, \quad y = g^{am}.$$

Then x and y commute, since both are powers of g . Moreover,

$$xy = g^{bn}g^{am} = g^{am+bn} = g,$$

and similarly $yx = g$. Also,

$$x^m = g^{bmn} = (g^{mn})^b = 1, \quad y^n = g^{amn} = (g^{mn})^a = 1.$$

Thus such elements $x, y \in G$ exist.

We now show that they are unique. Suppose that $x', y' \in G$ also satisfy

$$x'y' = g = y'x', \quad (x')^m = 1, \quad (y')^n = 1.$$

Since x' and y' commute, we may take powers of $x'y'$ termwise. Thus

$$g^{bn} = (x'y')^{bn} = (x')^{bn}(y')^{bn}.$$

But $(y')^{bn} = ((y')^n)^b = 1$, so

$$g^{bn} = (x')^{bn}.$$

Since $bn = 1 - am$, we get

$$(x')^{bn} = (x')^{1-am} = x'((x')^m)^{-a} = x'.$$

Therefore $x' = g^{bn} = x$. Similarly,

$$g^{am} = (x'y')^{am} = (x')^{am}(y')^{am} = (y')^{am},$$

and since $am = 1 - bn$,

$$(y')^{am} = (y')^{1-bn} = y'((y')^n)^{-b} = y'.$$

Therefore $y' = g^{am} = y$. Hence the required elements x and y exist and are unique. $\square$

Exercise 1.5. Let r, s , and t be positive integers greater than 1. Show that there is a finite group G having elements x and y such that x has order r , y has order s , and xy has order t .

Solution. First approach: a linear construction. Let

$$N = 2rst.$$

Choose a prime $p \nmid N$. By taking a suitable power $q = p^d$, we may arrange that

$$q \equiv 1 \pmod{N},$$

since p is invertible modulo N . Thus $2r, 2s, 2t$ all divide $q-1$. Since $\mathbb{F}_q^\times$ is cyclic, we may choose elements $u, v, w \in \mathbb{F}_q^\times$ of orders $2r, 2s, 2t$, respectively. Define

$$A = \begin{pmatrix} u & 1 \\ 0 & u^{-1} \end{pmatrix}, \quad B = \begin{pmatrix} v & 0 \\ \lambda & v^{-1} \end{pmatrix},$$

where

$$\lambda = w + w^{-1} - uv - u^{-1}v^{-1}.$$

Then $A, B \in \mathrm{SL}(2, \mathbb{F}_q)$. The characteristic polynomial of A is

$$X^2 - (u + u^{-1})X + 1,$$

with distinct roots u, u^{-1} , since u has order $2r > 2$. Hence A is conjugate over $\mathbb{F}_q$ to $\mathrm{diag}(u, u^{-1})$, and so A has order $2r$. Similarly, B has characteristic polynomial

$$X^2 - (v + v^{-1})X + 1$$

with distinct roots v, v^{-1} , and therefore B has order $2s$. Finally,

$$AB = \begin{pmatrix} uv + \lambda & v^{-1} \\ u^{-1}\lambda & u^{-1}v^{-1} \end{pmatrix},$$

so

$$\mathrm{tr}(AB) = uv + \lambda + u^{-1}v^{-1} = w + w^{-1}.$$

Since $\det(AB) = 1$, the characteristic polynomial of AB is

$$X^2 - (w + w^{-1})X + 1,$$

with distinct roots w, w^{-1} . Thus AB has order $2t$.

Now pass to the finite quotient

$$G = \mathrm{SL}(2, \mathbb{F}_q) / \{\pm I\}.$$

The element $-I$ is the unique element of order 2 in $\mathrm{SL}(2, \mathbb{F}_q)$: indeed, if $C^2 = I$, then C is diagonalizable with eigenvalues in $\{\pm 1\}$, and the condition $\det C = 1$ forces either $C = I$ or $C = -I$. Hence, if an element $g \in \mathrm{SL}(2, \mathbb{F}_q)$ has order $2n$, its image modulo $\{\pm I\}$ has order n . For g^n is the unique element of order 2, hence $g^n = -I$, so the image has order dividing n ; and if a smaller positive power became trivial in the quotient, then $g^m = \pm I$ for some $m < n$. The case $g^m = I$ contradicts $o(g) = 2n$, while $g^m = -I = g^n$ gives $g^{n-m} = I$, again impossible. Therefore the images x, y of A, B in G have orders r, s , respectively, and xy , the image of AB , has order t .

Second approach: finite quotients of triangle groups. We use one standard fact about a particular group given by generators and relations. For $a, b, c > 1$, the triangle group $\Delta(a, b, c)$ is the abstract group

$$\Delta(a, b, c) = \langle U, V \mid U^a = 1, V^b = 1, (UV)^c = 1 \rangle$$

generated by two formal elements U and V , subject to the relations $U^a = 1, V^b = 1$, and $(UV)^c = 1$. The word “triangle” is only the standard name for this presentation; no geometry is needed here. Equivalently, $\Delta(a, b, c)$ is the universal group with these three relations: whenever a group H has elements $\alpha, \beta \in H$ satisfying

$$\alpha^a = 1, \quad \beta^b = 1, \quad (\alpha\beta)^c = 1,$$

there is a homomorphism $\Delta(a, b, c) \rightarrow H$ sending U to α and V to β .

The standard fact we use is that U, V, UV have exact orders a, b, c , respectively, in $\Delta(a, b, c)$, and that $\Delta(a, b, c)$ is residually finite. Here residually finite means that if $g \neq 1$, then there is a homomorphism

from $\Delta(a, b, c)$ to some finite group in which the image of g is not 1. Now apply this fact with $a = r$, $b = s$, and $c = t$. Put

$$\Gamma = \langle X, Y \mid X^r = 1, Y^s = 1, (XY)^t = 1 \rangle.$$

In Γ , the elements X, Y, XY have exact orders r, s, t . Hence every element of the finite set

$$\mathcal{S} = \{X^i \mid 1 \leq i < r\} \cup \{Y^j \mid 1 \leq j < s\} \cup \{(XY)^k \mid 1 \leq k < t\}$$

is non-identity. By residual finiteness, for each $g \in \mathcal{S}$, there is a finite group Q_g and a homomorphism

$$\varphi_g: \Gamma \rightarrow Q_g$$

such that $\varphi_g(g) \neq 1$. Now define

$$Q = \prod_{g \in \mathcal{S}} Q_g, \quad \Phi: \Gamma \rightarrow Q, \quad \Phi(h) = (\varphi_g(h))_{g \in \mathcal{S}}.$$

Let $G = \Phi(\Gamma) \leq Q$, and set

$$x = \Phi(X), \quad y = \Phi(Y).$$

Since Q is finite, G is finite. Also, since $X^r = 1$, $Y^s = 1$, and $(XY)^t = 1$ in Γ , we have

$$x^r = 1, \quad y^s = 1, \quad (xy)^t = 1.$$

Thus $o(x) \mid r$, $o(y) \mid s$, and $o(xy) \mid t$.

It remains to show that no smaller positive powers become trivial in this finite quotient. Suppose $1 \leq i < r$. Since $X^i \in \mathcal{S}$, the X^i -coordinate of $\Phi(X^i)$ is $\varphi_{X^i}(X^i)$, which is not 1. Hence $\Phi(X^i) \neq 1$, so $x^i \neq 1$. Therefore $o(x) = r$.

Similarly, if $1 \leq j < s$, then the Y^j -coordinate of $\Phi(Y^j)$ is $\varphi_{Y^j}(Y^j) \neq 1$, so $y^j \neq 1$. Hence $o(y) = s$. Finally, if $1 \leq k < t$, then the $(XY)^k$ -coordinate of $\Phi((XY)^k)$ is $\varphi_{(XY)^k}((XY)^k) \neq 1$, so $(xy)^k \neq 1$. Hence $o(xy) = t$.

Third approach: Miller's permutation construction. We now give Miller's construction in modern notation. Miller's "substitutions" are permutations, and his "circular substitutions" are cycles. The proof is based on the following explicit overlap calculation. Let two cycles share a consecutive block

$$a_1, a_2, \dots, a_c$$

and have no other letters in common. If $c = 2h + 1$ is odd, arrange the two cycles as

$$S_1 = (a_1 \cdots a_{2j} a_{2j+1} \cdots a_{2h+1} b_1 \cdots b_p),$$

$$S_2 = (a_{2j+1} \cdots a_1 a_{2j+2} \cdots a_{2h+1} d_1 \cdots d_q).$$

Tracing the product gives

$$S_1 S_2 = (a_{2j+1} a_{2j+3} \cdots a_{2h+1} b_1 \cdots b_p a_{2j+2} a_{2j+4} \cdots a_{2h} d_1 \cdots d_q),$$

so $S_1 S_2$ is one cycle and the omitted common letters are exactly

$$a_1, \dots, a_{2j}.$$

Thus an odd overlap lets us omit any prescribed even number of common letters. Similarly, if $c = 2h$ is even and the same calculation is made with the first omitted block $a_1, \dots, a_{2j-1}$, then the product is one cycle omitting exactly $2j - 1$ common letters. Thus an even overlap lets us omit any prescribed odd number of common letters.

If the required parity is opposite to the parity supplied by the two-cycle calculation, Miller uses one cycle together with a permutation of the form

$$(\text{one transposition})(\text{one disjoint cycle}).$$

The same trace shows that this changes the parity of the number of omitted common letters. This is harmless whenever the corresponding order is even, since the least common multiple of 2 and an even cycle length is still the required even order. When both relevant orders are odd, no transposition is

inserted; instead, one uses two adjacent odd cycles and performs the same overlap twice. The first overlap changes the parity, and the second restores the required odd order. This is the only extra bookkeeping in Miller's all-odd case.

We now use this calculation to build permutations L, M, N of prescribed orders $\ell, m, n > 1$ with $LM = N$. Since

$$LM = N \iff MN^{-1} = L^{-1} \iff N^{-1}L = M^{-1},$$

we may construct the triple after arranging the desired orders as $\ell \leq m \leq n$, and then translate back to the original order. Write

$$n = am + b, \quad 0 \leq b < m.$$

Take M to be a product of a disjoint m -cycles,

$$M = (a_1 \cdots a_m)(a_{m+1} \cdots a_{2m}) \cdots (a_{(a-1)m+1} \cdots a_{am}).$$

The permutation L is chosen so that its cycles join these m -cycles end-to-end. If $a = 1$, this is exactly the overlap calculation above, since $m \leq n < m + \ell$. For instance, when ℓ is even and $a > 1$, the joining part may be written schematically as

$$\begin{aligned} L = & (a_1 a_{am+1})(a_2 a_{am+2}) \cdots (a_b a_{am+b}) \\ & \cdot (a_m a_{m+1})(a_{2m} a_{2m+1}) \cdots \\ & \cdot (a_{(a-1)m} a_{(a-1)m+1} \cdots a_{(a-1)m+\ell-1}). \end{aligned}$$

where the first row is omitted if $b = 0$. Here the displayed transpositions perform the ordinary joins, and the final ℓ -cycle makes $o(L) = \ell$. A direct trace explains the construction: starting at a_1 , the product LM first follows the first m -cycle of M ; when it reaches the end of that cycle, the next factor of L sends it to the beginning of the next m -cycle; the same process repeats until the trace reaches the final block. At the final block the overlap calculation above omits exactly $m - b$ common letters, so the resulting product has precisely

$$am + b = n$$

letters. Hence LM is an n -cycle.

If ℓ is odd, the same threading is done with ℓ -cycles rather than transpositions. The only issue is parity of the number of omitted common letters in the final overlap. When m is even, the transposition-plus-cycle variant above supplies the missing parity while keeping $o(M) = m$. When both ℓ and m are odd, two neighboring odd overlaps are used: the first overlap changes the parity of the omitted block and the second overlap joins the next m -cycle back into the same long trace. Since every component of L is then an ℓ -cycle and every component of M is an m -cycle, the orders remain

$$o(L) = \ell, \quad o(M) = m.$$

The same trace as before shows that the product is still a single n -cycle. Thus in all cases we have constructed permutations L, M, N such that

$$o(L) = \ell, \quad o(M) = m, \quad o(N) = n, \quad LM = N.$$

We now return to r, s, t . If t is the largest of the three orders, take $\ell = r$, $m = s$, and $n = t$, and set $x = L$, $y = M$. If r is the largest, take $\ell = s$, $m = t$, and $n = r$. Then $LM = N$, so $N^{-1}L = M^{-1}$; setting $x = N^{-1}$ and $y = L$ gives $o(x) = r$, $o(y) = s$, and $o(xy) = t$. If s is the largest, take $\ell = t$, $m = r$, and $n = s$. Then $LM = N$, so $MN^{-1} = L^{-1}$; setting $x = M$ and $y = N^{-1}$ gives $o(x) = r$, $o(y) = s$, and $o(xy) = t$. In all cases the permutations act on a finite set Ω . Let

$$G = \langle x, y \rangle \leq \text{Sym}(\Omega).$$

Then G is finite, and by construction $o(x) = r$, $o(y) = s$, and $o(xy) = t$. $\therefore$ the required finite group exists. $\square$

Exercise 1.6. Let X and Y be subsets of a group G . Are $\langle X \rangle \cap \langle Y \rangle$ and $\langle X \cap Y \rangle$ necessarily equal? Are $\langle \langle X \rangle \cup \langle Y \rangle \rangle$ and $\langle X \cup Y \rangle$ necessarily equal?

Solution. The first pair need not be equal. Let $G = C_3 = \langle g \rangle$, $X = \{1, g\}$, and $Y = \{1, g^2\}$. Then $\langle X \rangle = G$, since $g \in X$, and $\langle Y \rangle = G$, since g^2 also generates C_3 . Hence

$$\langle X \rangle \cap \langle Y \rangle = G.$$

However, $X \cap Y = \{1\}$, so

$$\langle X \cap Y \rangle = \{1\}.$$

Thus $\langle X \rangle \cap \langle Y \rangle$ and $\langle X \cap Y \rangle$ are not necessarily equal.

The second pair is always equal. Since $X \subseteq \langle X \rangle$ and $Y \subseteq \langle Y \rangle$, we have

$$X \cup Y \subseteq \langle X \rangle \cup \langle Y \rangle.$$

Therefore

$$\langle X \cup Y \rangle \leq \langle \langle X \rangle \cup \langle Y \rangle \rangle.$$

Conversely, since $X \subseteq X \cup Y$ and $Y \subseteq X \cup Y$, we have

$$\langle X \rangle \leq \langle X \cup Y \rangle, \quad \langle Y \rangle \leq \langle X \cup Y \rangle.$$

Hence

$$\langle X \rangle \cup \langle Y \rangle \subseteq \langle X \cup Y \rangle.$$

Since $\langle X \cup Y \rangle$ is a subgroup containing $\langle X \rangle \cup \langle Y \rangle$, it contains the subgroup generated by that union, so

$$\langle \langle X \rangle \cup \langle Y \rangle \rangle \leq \langle X \cup Y \rangle.$$

$$\therefore \langle \langle X \rangle \cup \langle Y \rangle \rangle = \langle X \cup Y \rangle. \quad \square$$

Exercise 1.7. Let G be a finite group and let $H \leq G$. Show that there is a subset T of G which is simultaneously a left transversal for H and a right transversal for H .

Solution. We use the following elementary matching fact. Suppose $A_1, \dots, A_n$ are finite sets of possible choices. If, for every $I \subseteq \{1, \dots, n\}$,

$$\left| \bigcup_{i \in I} A_i \right| \geq |I|,$$

then we can choose distinct representatives $a_i \in A_i$.

Let

$$L_1, \dots, L_n$$

be the distinct left cosets of H , and let

$$R_1, \dots, R_n$$

be the distinct right cosets of H . There are the same number n of each, since both left and right cosets partition G into subsets of size $|H|$. For each i , define

$$A_i = \{j \in \{1, \dots, n\} \mid L_i \cap R_j \neq \emptyset\}.$$

Thus A_i is the set of right cosets which the left coset L_i meets. We check the matching condition. Choose any m left cosets, say those indexed by I , where $|I| = m$. Suppose these m left cosets meet q right cosets. Since distinct left cosets are disjoint,

$$\left| \bigcup_{i \in I} L_i \right| = m|H|.$$

This union is contained in the union of the q right cosets it meets, whose total size is $q|H|$. Hence

$$m|H| \leq q|H|,$$

so $m \leq q$. Therefore any m left cosets meet at least m right cosets. In terms of the sets A_i , this says exactly that

$$\left| \bigcup_{i \in I} A_i \right| \geq |I|.$$

By the matching fact, choose distinct $j_1, \dots, j_n$ such that $j_i \in A_i$ for every i . Since $j_i \in A_i$,

$$L_i \cap R_{j_i} \neq \emptyset,$$

so choose

$$t_i \in L_i \cap R_{j_i}.$$

Set

$$T = \{t_1, \dots, t_n\}.$$

Because $t_i \in L_i$, the set T contains one element from each left coset. Since the L_i are disjoint, it contains exactly one element from each left coset, so T is a left transversal for H .

Because the indices $j_1, \dots, j_n$ are distinct, the right cosets $R_{j_1}, \dots, R_{j_n}$ are all distinct. There are n of them, and there are only n right cosets in total, so they are all the right cosets. Since $t_i \in R_{j_i}$, the set T also contains exactly one element from each right coset. Hence T is a right transversal for H .

$\therefore$ there exists a subset $T \subseteq G$ which is simultaneously a left transversal and a right transversal for H . $\square$

Exercise 1.8. Suppose that $\mathcal{C}$ is a family of subsets of a group G which forms a partition of G , and suppose further that $gC \in \mathcal{C}$ for any $g \in G$ and $C \in \mathcal{C}$. (Recall that a *partition* of a set S is a collection $\mathcal{S}$ of subsets of S with the property that every element of S lies in exactly one member of $\mathcal{S}$.) Show that $\mathcal{C}$ is the set of cosets of some subgroup of G .

Solution. Since $\mathcal{C}$ is a partition of G , there is a unique member $H \in \mathcal{C}$ such that $1 \in H$. We first show that $H \leq G$.

Let $h \in H$. By the hypothesis, $hH \in \mathcal{C}$. Also, since $1 \in H$, we have $h = h1 \in hH$. Thus both H and hH are members of $\mathcal{C}$ containing h . Since $\mathcal{C}$ is a partition, we must have

$$hH = H.$$

Hence for any $h, k \in H$, we have $hk \in hH = H$, so H is closed under multiplication. Also, since $1 \in H = hH$, there exists $u \in H$ such that $hu = 1$. Thus $u = h^{-1}$, so $h^{-1} \in H$. Therefore $H \leq G$.

Now let $C \in \mathcal{C}$. Choose $g \in C$. By the hypothesis, $gH \in \mathcal{C}$, and since $1 \in H$, we have $g \in gH$. Thus both C and gH are members of $\mathcal{C}$ containing g . Since $\mathcal{C}$ is a partition, $C = gH$. Therefore every member of $\mathcal{C}$ is a left coset of H . Conversely, for every $g \in G$, the set gH belongs to $\mathcal{C}$ by the hypothesis.

$\therefore \mathcal{C} = \{gH \mid g \in G\}$, the set of left cosets of the subgroup H . $\square$

Exercise 1.9. Suppose that $\mathcal{C}$ is a family of subsets of a group G which forms a partition of G , and suppose further that $XY \in \mathcal{C}$ for any $X, Y \in \mathcal{C}$. Show that exactly one of the sets belonging to $\mathcal{C}$ is a subgroup of G , that this subgroup is normal in G , and that $\mathcal{C}$ consists of its cosets.

Solution. Since $\mathcal{C}$ is a partition of G , there is a unique member $H \in \mathcal{C}$ such that $1 \in H$. We first show that $H \leq G$.

By the hypothesis, $HH \in \mathcal{C}$. Since $1 \in H$, we have $1 = 1 \cdot 1 \in HH$. Thus H and HH are both members of $\mathcal{C}$ containing 1, so $HH = H$. Hence H is closed under multiplication.

Now let $h \in H$. Let $K \in \mathcal{C}$ be the unique member containing h^{-1} . By the hypothesis, $HK \in \mathcal{C}$. Since $h \in H$ and $h^{-1} \in K$, we have $1 = hh^{-1} \in HK$. Thus HK and H are both members of $\mathcal{C}$ containing 1, so $HK = H$. Since $1 \in H$ and $h^{-1} \in K$, it follows that $h^{-1} = 1h^{-1} \in HK = H$. Therefore $H \leq G$.

We now show that every member of $\mathcal{C}$ is a left coset of H . Let $X \in \mathcal{C}$, and choose $x \in X$. By the hypothesis, $XH \in \mathcal{C}$. Since $1 \in H$, we have $x = x1 \in XH$. Thus XH and X are both members of $\mathcal{C}$ containing x , so $XH = X$. Hence $xH \subseteq XH = X$.

For the reverse inclusion, let $X' \in \mathcal{C}$ be the unique member containing x^{-1} . By the hypothesis, $X'X \in \mathcal{C}$. Since $x^{-1}x = 1 \in X'X$, we have $X'X = H$. If $y \in X$, then $x^{-1}y \in X'X = H$, and hence

$$y = x(x^{-1}y) \in xH.$$

Thus $X \subseteq xH$. Therefore $X = xH$, and so every member of $\mathcal{C}$ is a left coset of H . Conversely, if $g \in G$, and $X \in \mathcal{C}$ contains g , then $X = gH$. Hence

$$\mathcal{C} = \{gH \mid g \in G\}.$$

The subgroup H is the only member of $\mathcal{C}$ which is a subgroup, since it is the only member containing 1. It remains to show that H is normal. Let $g \in G$. Since $gH, g^{-1}H \in \mathcal{C}$, the product $(gH)(g^{-1}H)$ is a member of $\mathcal{C}$. It contains 1, since $g \in gH$, $g^{-1} \in g^{-1}H$, and $gg^{-1} = 1$. Thus

$$(gH)(g^{-1}H) = H.$$

Now let $h \in H$. Since $gh \in gH$ and $g^{-1} \in g^{-1}H$, we have

$$ghg^{-1} \in (gH)(g^{-1}H) = H.$$

Thus $gHg^{-1} \subseteq H$. Replacing g by g^{-1} , we also have $g^{-1}Hg \subseteq H$, and conjugating this inclusion by g gives $H \subseteq gHg^{-1}$. Hence $gHg^{-1} = H$ for all $g \in G$. Therefore $H \trianglelefteq G$.

$\therefore$ exactly one member of $\mathcal{C}$ is a subgroup, this subgroup is normal in G , and $\mathcal{C}$ consists of its cosets. $\square$

Exercise 1.10. Prove the following generalization of Proposition 1.8: If G is a finite group and $H \leq G$ is such that $|G : H|$ is equal to the smallest prime divisor of $|G|$, then $H \trianglelefteq G$.

Solution. Let $p = |G : H|$. By hypothesis, p is the smallest prime divisor of $|G|$. Let

$$\Omega = \{xH \mid x \in G\}$$

be the set of left cosets of H . Then $|\Omega| = p$. Define

$$\varphi: G \rightarrow S_\Omega$$

by

$$\varphi(g)(xH) = gxH$$

for all $g, x \in G$. This is well-defined: if $xH = yH$, then $gxH = gyH$. For each $g \in G$, the map $\varphi(g)$ is a permutation of Ω , with inverse $\varphi(g^{-1})$. Also,

$$\varphi(g_1g_2)(xH) = g_1g_2xH = \varphi(g_1)(g_2xH) = (\varphi(g_1)\varphi(g_2))(xH),$$

so φ is a homomorphism.

Let $K = \ker \varphi$. Then $K \trianglelefteq G$. We claim that $K \leq H$. If $k \in K$, then $\varphi(k)$ fixes every left coset, so in particular

$$H = \varphi(k)(H) = kH.$$

Thus $k \in H$, and hence $K \leq H$.

By the Fundamental Theorem on Homomorphisms, $G/K \cong \text{im } \varphi \leq S_\Omega$. Hence

$$|G : K| = |\text{im } \varphi| \mid |S_\Omega| = p!.$$

Also, by Lagrange's Theorem, $|G : K| \mid |G|$. Therefore every prime divisor of $|G : K|$ is a prime divisor of $|G|$, and hence is at least p . On the other hand, since $|G : K| \mid p!$, every prime divisor of $|G : K|$ is at most p . Thus the only possible prime divisor of $|G : K|$ is p .

Since $K \leq H \leq G$, we have

$$|G : K| = |G : H| |H : K| = p |H : K|.$$

Thus $p \mid |G : K|$. Since the only possible prime divisor of $|G : K|$ is p , we may write $|G : K| = p^a$ for some $a \geq 1$. But $|G : K| \mid p!$, and $p^2 \nmid p!$, so $a = 1$. Hence

$$|G : K| = p.$$

It follows that $p \mid |H : K| = p$, so $|H : K| = 1$. Therefore $H = K$. Since $K \trianglelefteq G$, we conclude that $H \trianglelefteq G$. $\square$

Further Exercises

If $K \trianglelefteq H \leq G$, then H/K is called a *section* of G . We say that two sections H_1/K_1 and H_2/K_2 of G are *incident* if every coset of K_1 in H_1 has non-empty intersection with exactly one coset of K_2 in H_2 , and vice versa. (In other words, two sections are incident if the relation of non-empty intersection gives a bijective correspondence between their elements.)

Exercise 1.11. Show that incident sections are isomorphic.

Solution. Let H_1/K_1 and H_2/K_2 be incident sections of G . Define

$$\varphi: H_1/K_1 \rightarrow H_2/K_2$$

by letting $\varphi(xK_1)$ be the unique coset yK_2 such that

$$xK_1 \cap yK_2 \neq \emptyset.$$

This is well-defined by the definition of incidence. We first show that φ is bijective. If $\varphi(xK_1) = \varphi(x'K_1) = yK_2$, then yK_2 has non-empty intersection with both xK_1 and $x'K_1$. Since incidence also says that each coset of K_2 in H_2 has non-empty intersection with exactly one coset of K_1 in H_1 , we have $xK_1 = x'K_1$. Hence φ is injective. If $yK_2 \in H_2/K_2$, then, again by incidence, yK_2 has non-empty intersection with some coset $xK_1 \in H_1/K_1$, and then $\varphi(xK_1) = yK_2$. Hence φ is surjective.

It remains to show that φ is a homomorphism. Suppose

$$\varphi(xK_1) = yK_2 \quad \text{and} \quad \varphi(x'K_1) = y'K_2.$$

Choose

$$a \in xK_1 \cap yK_2 \quad \text{and} \quad b \in x'K_1 \cap y'K_2.$$

Then $a = xk$ and $b = x'k'$ for some $k, k' \in K_1$. Hence

$$ab = xkx'k' = xx'(x'^{-1}kx')k' \in xx'K_1,$$

because $K_1 \trianglelefteq H_1$ and $x' \in H_1$. Similarly, writing $a = y\ell$ and $b = y'\ell'$ with $\ell, \ell' \in K_2$, we get

$$ab = y\ell y'\ell' = yy'(y'^{-1}\ell y')\ell' \in yy'K_2,$$

because $K_2 \trianglelefteq H_2$ and $y' \in H_2$. Therefore

$$xx'K_1 \cap yy'K_2 \neq \emptyset.$$

By the uniqueness in the definition of incidence, this implies

$$\varphi(xx'K_1) = yy'K_2.$$

On the other hand,

$$\varphi(xK_1)\varphi(x'K_1) = (yK_2)(y'K_2) = yy'K_2.$$

Thus

$$\varphi((xK_1)(x'K_1)) = \varphi(xK_1)\varphi(x'K_1).$$

Therefore φ is a bijective homomorphism, so $H_1/K_1 \cong H_2/K_2$. $\square$

Exercise 1.12. (cont.) Suppose that $N \trianglelefteq G$ and $H \leq G$. Show that HN/N and $H/(H \cap N)$ are incident. (Exercises 1.11 and 1.12 provide an alternate proof of the First Isomorphism Theorem.)

Solution. Let $K = H \cap N$. By Proposition 1.7, $HN \leq G$ and $K \trianglelefteq H$. Since $N \trianglelefteq G$ and $HN \leq G$, we also have $N \trianglelefteq HN$, so HN/N and H/K are sections of G . We show that they are incident.

First let $xN \in HN/N$. Since $x \in HN$, we may write $x = hn$ for some $h \in H$ and $n \in N$. Hence

$$xN = hnN = hN.$$

Since $1 \in K$, we have $h \in hN \cap hK$, so $hN \cap hK \neq \emptyset$. Thus every coset of N in HN has non-empty intersection with at least one coset of K in H . We now prove uniqueness. Suppose that

$$hN \cap h'K \neq \emptyset$$

for some $h' \in H$. Choose $z \in hN \cap h'K$. Then $z = hn_1 = h'k$ for some $n_1 \in N$ and $k \in K$, and hence

$$h^{-1}h' = n_1k^{-1}.$$

Since $n_1 \in N$ and $k \in K \leq N$, we have $n_1k^{-1} \in N$. Since also $h, h' \in H$, we have $h^{-1}h' \in H$. Hence $h^{-1}h' \in H \cap N = K$, so $hK = h'K$. Therefore each coset of N in HN intersects exactly one coset of K in H .

Conversely, let $hK \in H/K$. Since $h \in hK \cap hN$, the coset hK has non-empty intersection with at least one coset of N in HN . Suppose that $hK \cap xN \neq \emptyset$ for some $xN \in HN/N$. As above, we may write $xN = h'N$ for some $h' \in H$. Choose $z \in hK \cap h'N$. Then $z = hk = h'n_2$ for some $k \in K$ and $n_2 \in N$, and hence

$$h^{-1}h' = kn_2^{-1}.$$

Since $k \in K \leq N$ and $n_2 \in N$, we get $h^{-1}h' \in N$. Therefore $h'N = hN$. Thus each coset of K in H intersects exactly one coset of N in HN .

$\therefore HN/N$ and $H/(H \cap N)$ are incident. $\square$

If L/M is a section of G and $H \leq G$, then the *projection* of H on L/M is the subset of L/M consisting of those cosets of M in L which contain elements of H .

Exercise 1.13. (cont.) Show that the projection of H on L/M is the subgroup $(L \cap H)M/M$ of L/M .

Solution. Let

$$P = \{lM \in L/M \mid lM \cap H \neq \emptyset\}$$

be the projection of H on L/M . Since L/M is a section of G , we have $M \trianglelefteq L$. Let $\eta: L \rightarrow L/M$ be the natural map. By Proposition 1.11,

$$\eta(L \cap H) = (L \cap H)M/M$$

is a subgroup of L/M . It remains to show that $P = \eta(L \cap H)$.

First let $lM \in P$. Then $lM \cap H \neq \emptyset$, so we may choose $h \in lM \cap H$. Since $h \in lM \subseteq L$, and also $h \in H$, we have $h \in L \cap H$. Moreover $h \in lM$, so $hM = lM$. Hence

$$lM = hM \in \eta(L \cap H).$$

Thus $P \subseteq \eta(L \cap H)$.

Conversely, let $xM \in \eta(L \cap H)$. Then $xM = hM$ for some $h \in L \cap H$. Since $h \in H$ and $h \in hM = xM$, we have $xM \cap H \neq \emptyset$. Hence $xM \in P$. Thus $\eta(L \cap H) \subseteq P$.

$\therefore P = \eta(L \cap H) = (L \cap H)M/M$, so the projection of H on L/M is the subgroup $(L \cap H)M/M$ of L/M . $\square$

Let H_1/K_1 and H_2/K_2 be sections of a group G .

Exercise 1.14. (cont.) Show that the projection of K_2 on H_1/K_1 is a normal subgroup of the projection of H_2 on H_1/K_1 . The quotient group obtained thereby is called the projection of H_2/K_2 on H_1/K_1 .

Solution. By Exercise 1.13, the projection of K_2 on H_1/K_1 is

$$(H_1 \cap K_2)K_1/K_1,$$

and the projection of H_2 on H_1/K_1 is

$$(H_1 \cap H_2)K_1/K_1.$$

We first show that $H_1 \cap K_2 \trianglelefteq H_1 \cap H_2$. Let $x \in H_1 \cap K_2$ and $y \in H_1 \cap H_2$. Since $x, y \in H_1$, we have $xyx^{-1} \in H_1$. Since $x \in K_2$, $y \in H_2$, and $K_2 \trianglelefteq H_2$, we also have $xyx^{-1} \in K_2$. Hence

$$xyx^{-1} \in H_1 \cap K_2.$$

Therefore $H_1 \cap K_2 \trianglelefteq H_1 \cap H_2$.

Now take

$$xK_1 \in (H_1 \cap K_2)K_1/K_1 \quad \text{and} \quad yK_1 \in (H_1 \cap H_2)K_1/K_1,$$

where $x \in H_1 \cap K_2$ and $y \in H_1 \cap H_2$. Then

$$(yK_1)(xK_1)(yK_1)^{-1} = (yxy^{-1})K_1.$$

By the first paragraph, $yxy^{-1} \in H_1 \cap K_2$, so

$$(yxy^{-1})K_1 \in (H_1 \cap K_2)K_1/K_1.$$

Thus

$$(H_1 \cap K_2)K_1/K_1 \trianglelefteq (H_1 \cap H_2)K_1/K_1.$$

$\therefore$ the projection of K_2 on H_1/K_1 is a normal subgroup of the projection of H_2 on H_1/K_1 . $\square$

Exercise 1.15. (cont.) Show that the projection of H_1/K_1 on H_2/K_2 and the projection of H_2/K_2 on H_1/K_1 are incident. Deduce that, if $H_1, H_2 \leq G$, $K_1 \trianglelefteq H_1$, and $K_2 \trianglelefteq H_2$, then

$$(H_1 \cap H_2)K_1 / (H_1 \cap K_2)K_1 \cong (H_1 \cap H_2)K_2 / (K_1 \cap H_2)K_2.$$

Solution. Let

$$A = H_1 \cap H_2, \quad B_1 = (H_1 \cap K_2)K_1, \quad B_2 = (K_1 \cap H_2)K_2.$$

By Exercise 1.13 and Exercise 1.14, the projection of H_2/K_2 on H_1/K_1 is

$$AK_1/B_1,$$

and the projection of H_1/K_1 on H_2/K_2 is

$$AK_2/B_2.$$

Every coset of B_1 in AK_1 has the form aB_1 for some $a \in A$, since $K_1 \leq B_1$. Similarly, every coset of B_2 in AK_2 has the form aB_2 for some $a \in A$, since $K_2 \leq B_2$. For each $a \in A$, the two cosets aB_1 and aB_2 have non-empty intersection, since both contain a .

We now prove uniqueness. Suppose that

$$aB_1 \cap bB_2 \neq \emptyset$$

for some $a, b \in A$. Choose $z \in aB_1 \cap bB_2$. Then $z = as = bt$ for some $s \in B_1$ and $t \in B_2$. Write

$$s = ck, \quad c \in H_1 \cap K_2, \quad k \in K_1,$$

and

$$t = d\ell, \quad d \in K_1 \cap H_2, \quad \ell \in K_2.$$

From $ack = bdl$, we get

$$k = c^{-1}a^{-1}bd\ell.$$

Since $a, b \in A \leq H_2$, $c \in K_2 \leq H_2$, $d \in H_2$, and $\ell \in K_2 \leq H_2$, it follows that $k \in H_2$. Thus $k \in K_1 \cap H_2$. Since $c \in K_2$ and $K_2 \trianglelefteq H_2$, we have

$$s = ck = k(k^{-1}ck) \in (K_1 \cap H_2)K_2 = B_2.$$

Since $t \in B_2$, the equality $as = bt$ gives

$$b^{-1}a = ts^{-1} \in B_2.$$

Hence $aB_2 = bB_2$. Therefore each coset of B_1 in AK_1 intersects at most one coset of B_2 in AK_2 . Conversely, suppose that

$$aB_2 \cap bB_1 \neq \emptyset$$

for some $a, b \in A$. Choose $z \in aB_2 \cap bB_1$. Then $z = at = bs$ with $t \in B_2$ and $s \in B_1$. Write

$$t = d\ell, \quad d \in K_1 \cap H_2, \quad \ell \in K_2,$$

and

$$s = ck, \quad c \in H_1 \cap K_2, \quad k \in K_1.$$

From $ad\ell = bck$, we get

$$\ell = d^{-1}a^{-1}bck.$$

Since $a, b \in A \leq H_1$, $d \in K_1 \leq H_1$, $c \in H_1$, and $k \in K_1 \leq H_1$, it follows that $\ell \in H_1$. Thus $\ell \in H_1 \cap K_2$. Since $d \in K_1$ and $K_1 \trianglelefteq H_1$, we have

$$t = d\ell = \ell(\ell^{-1}d\ell) \in (H_1 \cap K_2)K_1 = B_1.$$

Since $s \in B_1$, the equality $at = bs$ gives

$$b^{-1}a = st^{-1} \in B_1.$$

Hence $aB_1 = bB_1$. Therefore each coset of B_2 in AK_2 intersects at most one coset of B_1 in AK_1 . Thus the projections AK_1/B_1 and AK_2/B_2 are incident. $\square$

Theorem (Third Isomorphism Theorem). *Let $H_1, H_2 \leq G$, let $K_1 \trianglelefteq H_1$, and let $K_2 \trianglelefteq H_2$. Then*

$$(H_1 \cap H_2)K_1 / (H_1 \cap K_2)K_1 \cong (H_1 \cap H_2)K_2 / (K_1 \cap H_2)K_2.$$

Proof. By Exercise 1.15, the sections

$$(H_1 \cap H_2)K_1 / (H_1 \cap K_2)K_1 \quad \text{and} \quad (H_1 \cap H_2)K_2 / (K_1 \cap H_2)K_2$$

are incident. By Exercise 1.11, incident sections are isomorphic. Hence

$$(H_1 \cap H_2)K_1 / (H_1 \cap K_2)K_1 \cong (H_1 \cap H_2)K_2 / (K_1 \cap H_2)K_2.$$

□

(This result is also called the fourth isomorphism theorem, or Zassenhaus' lemma — after its discoverer, who proved it as a student at the age of 21 — or even the butterfly lemma. The last name refers to the shape of the diagram showing the inclusion relations between the many subgroups involved.)

2 Automorphisms

The Automorphism Group

- The set of automorphisms of a group G is denoted $\text{Aut}(G)$.
- If $\varphi, \rho \in \text{Aut}(G)$, then $\varphi \circ \rho \in \text{Aut}(G)$, so composition is a binary operation on $\text{Aut}(G)$.
- Under composition, $\text{Aut}(G)$ is a group: identity is the trivial automorphism, and the inverse of φ is its set-map inverse φ^{-1} .
- Call $\text{Aut}(G)$ with this operation the *automorphism group* of G ; we may write $\varphi\rho$ for $\varphi \circ \rho$.

Inner Automorphisms, $\text{Inn}(G)$, and the Center

- Each $g \in G$ defines a conjugation homomorphism $\varphi_g : G \rightarrow G$ by $\varphi_g(x) = gxg^{-1}$.
- Each φ_g is an automorphism: for $x \in G$, $x = \varphi_g(g^{-1}xg)$, and if $\varphi_g(x) = \varphi_g(y)$ then $x = y$ by cancellation. These are the *inner automorphisms* of G .
- $\varphi_g\varphi_h = \varphi_{gh}$ for $g, h \in G$, since $g(hxh^{-1})g^{-1} = (gh)x(gh)^{-1}$. Hence there is a homomorphism $G \rightarrow \text{Aut}(G)$ sending g to φ_g .
- Its image is the *inner automorphism group* of G , denoted $\text{Inn}(G)$; its kernel is the *center* of G , denoted $Z(G)$. Observe

$$Z(G) = \{g \in G \mid \varphi_g(x) = x \text{ for all } x \in G\} = \{g \in G \mid gx = xg \text{ for all } x \in G\},$$

so $Z(G)$ consists of those elements of G which commute with every element of G .

- Clearly G is abelian iff $Z(G) = G$.

Outer Automorphisms, $\text{Out}(G)$

- If $\sigma \in \text{Aut}(G)$ and $\varphi_g \in \text{Inn}(G)$, then $\sigma\varphi_g\sigma^{-1} = \varphi_{\sigma(g)}$. Hence $\text{Inn}(G) \trianglelefteq \text{Aut}(G)$.
- The quotient $\text{Aut}(G)/\text{Inn}(G)$ is the *outer automorphism group* of G , denoted $\text{Out}(G)$.
- “Outer automorphism” refers to automorphisms of G which are not inner and which thus have non-trivial image in $\text{Out}(G)$.
- If G is abelian, then all non-trivial automorphisms of G are outer in this sense, since $\text{Inn}(G) = 1$.

Automorphism Groups of Cyclic Groups

- Let $G = \langle x \rangle \cong \mathbb{Z}$, and let φ be an automorphism of G . Then $\varphi(x)$ must generate G ; the only generators are x and x^{-1} . So φ either fixes each element or sends each to its inverse, and $\text{Aut}(G) \cong \mathbb{Z}_2$.
- Let $n \in \mathbb{N}$ and $G = \langle x \rangle \cong \mathbb{Z}_n$. Suppose φ is an endomorphism of G . Then $\varphi(x) = x^m$ for some $0 \leq m < n$, so φ sends every element to its m th power. Hence G has exactly n endomorphisms, namely the m th power maps σ_m for $0 \leq m < n$.

Proposition 2.1. *Let $G = \langle x \rangle \cong \mathbb{Z}_n$ for $n \in \mathbb{N}$, and for each $0 \leq m < n$ let σ_m be the endomorphism of G sending x to x^m . Then $\text{Aut}(G)$ consists precisely of those σ_m for which $m \neq 0$ and $\gcd(m, n) = 1$. Furthermore, $\text{Aut}(G)$ is abelian and is isomorphic with the group $(\mathbb{Z}/n\mathbb{Z})^\times$ of units of the ring $\mathbb{Z}/n\mathbb{Z}$.*

Proof. The map σ_0 has trivial image, so it is not an automorphism. Let $1 \leq m < n$ and consider σ_m . Suppose first that $\gcd(m, n) = 1$. Then there exist $a, b \in \mathbb{Z}$ such that $am + bn = 1$, and hence

$$\sigma_m(x^a) = x^{am} = x^{1-bn} = x(x^n)^{-b} = x.$$

Thus $x \in \text{im } \sigma_m$, so σ_m is surjective. Since G is finite, σ_m is also injective, and therefore $\sigma_m \in \text{Aut}(G)$. Conversely, suppose that $\sigma_m \in \text{Aut}(G)$. Since σ_m is surjective, there is some $a \in \mathbb{Z}$ such that $x = \sigma_m(x^a) = x^{am}$. Hence $x^{am-1} = 1$, so $am - 1 = bn$ for some $b \in \mathbb{Z}$. It follows that any common divisor

of m and n divides 1, and therefore $\gcd(m, n) = 1$. This proves the first assertion.

Now let $1 \leq m_1, m_2 < n$ with $\gcd(m_1, n) = \gcd(m_2, n) = 1$. If t is chosen with $1 \leq t < n$ and $m_1 m_2 \equiv t \pmod{n}$, then

$$(\sigma_{m_1} \sigma_{m_2})(x) = \sigma_{m_1}(x^{m_2}) = x^{m_1 m_2} = x^t.$$

Hence $\sigma_{m_1} \sigma_{m_2} = \sigma_t = \sigma_{m_2} \sigma_{m_1}$, so $\text{Aut}(G)$ is abelian. Finally, the map

$$\text{Aut}(G) \rightarrow (\mathbb{Z}/n\mathbb{Z})^\times, \quad \sigma_m \mapsto m + n\mathbb{Z}$$

is a bijection by the first assertion, and the computation above shows that it preserves multiplication. Hence $\text{Aut}(G) \cong (\mathbb{Z}/n\mathbb{Z})^\times$. $\square$

The Totient and Primitive Roots

- The *totient* of $n \in \mathbb{N}$ is the number of positive integers less than n and coprime to n (the value at n of the Euler phi-function).
- If $n = p_1^{a_1} \cdots p_r^{a_r}$ with the p_i distinct primes, then the totient of n is $(p_1^{a_1} - p_1^{a_1-1}) \cdots (p_r^{a_r} - p_r^{a_r-1})$. To see this, note that m is coprime to n iff no p_i divides m . Among $1, \dots, n$, exactly n/p_i integers are divisible by p_i , exactly $n/(p_i p_j)$ are divisible by both p_i and p_j , and so on. Hence inclusion-exclusion gives

$$\varphi(n) = n \prod_{i=1}^r \left(1 - \frac{1}{p_i}\right) = \prod_{i=1}^r (p_i^{a_i} - p_i^{a_i-1}).$$

- By Proposition 2.1, the order of $\text{Aut}(\mathbb{Z}_n)$ is the totient of n . In particular, $|\text{Aut}(\mathbb{Z}_p)| = p - 1$ when p is prime.

Proposition 2.2. *Let p be a prime. Then $\text{Aut}(\mathbb{Z}_p) \cong \mathbb{Z}_{p-1}$.*

Proof. Let $F = \mathbb{F}_p$. By Proposition 2.1, $\text{Aut}(\mathbb{Z}_p)$ is isomorphic with $F^\times = (\mathbb{Z}/p\mathbb{Z})^\times$. Thus it suffices to show that $F^\times$ is cyclic of order $p - 1$.

For each divisor d of $p - 1$, define

$$f_d = |\{x \in F^\times \mid o(x) = d\}|, \quad z_d = |\{x \in \mathbb{Z}_{p-1} \mid o(x) = d\}|.$$

We first show that $f_d \leq z_d$ for every $d \mid (p - 1)$. If $x \in F^\times$ has order dividing d , then $x^d = 1$, so x is a root of $T^d - 1 \in F[T]$. Since this polynomial has degree d , it has at most d roots in F . Now suppose $F^\times$ has an element x of order d . Then the d elements of $\langle x \rangle$ are all roots of $T^d - 1$, so they are all the roots. Hence every element of $F^\times$ of order d lies in $\langle x \rangle \cong \mathbb{Z}_d$. Therefore either $f_d = 0$, or f_d is the number of elements of order d in a cyclic group of order d . By Theorem 1.4, the cyclic group $\mathbb{Z}_{p-1}$ has a unique subgroup of order d ; every element of order d generates a subgroup of order d , so all elements of order d in $\mathbb{Z}_{p-1}$ lie in this unique subgroup. This subgroup is cyclic of order d , and hence $f_d \leq z_d$.

Since every element of $F^\times$ has order dividing $|F^\times| = p - 1$, and every element of $\mathbb{Z}_{p-1}$ has order dividing $p - 1$, we have

$$\sum_{d \mid (p-1)} f_d = |F^\times| = p - 1 = |\mathbb{Z}_{p-1}| = \sum_{d \mid (p-1)} z_d.$$

Since $f_d \leq z_d$ for every $d \mid (p - 1)$ and the sums are equal, we must have $f_d = z_d$ for every $d \mid (p - 1)$. In particular, $f_{p-1} = z_{p-1}$. But $\mathbb{Z}_{p-1}$ has an element of order $p - 1$, so $z_{p-1} > 0$, and hence $f_{p-1} > 0$. Therefore $F^\times$ has an element of order $p - 1$, so $F^\times \cong \mathbb{Z}_{p-1}$.

$\therefore \text{Aut}(\mathbb{Z}_p) \cong \mathbb{Z}_{p-1}$. $\square$

Primitive Roots Modulo n

- Let $G = \langle x \rangle \cong \mathbb{Z}_n$, $1 \leq m \leq n$, $\gcd(m, n) = 1$. Induction gives $(\sigma_m)^k(x) = x^{m^k}$.
- The order of σ_m is the least positive k such that $x^{m^k} = x$, i.e. the smallest $k \in \mathbb{N}$ with $m^k \equiv 1 \pmod{n}$.
- If the order of σ_m equals the totient of n , we call m a *primitive root modulo n* .
- $\text{Aut}(\mathbb{Z}_n)$ is cyclic iff there is a primitive root modulo n .

- For composite n , the structure of $\text{Aut}(\mathbb{Z}_n)$ belongs to number theory more than group theory.

Theorem 2.3. *For $n > 1$, $\text{Aut}(\mathbb{Z}_n)$ is cyclic iff $n = 2$ or 4 , or $n = p^k$ or $2p^k$ for some odd prime p and some $k \in \mathbb{N}$.*

Proof. By Proposition 2.1,

$$\text{Aut}(\mathbb{Z}_n) \cong (\mathbb{Z}/n\mathbb{Z})^\times.$$

Under this isomorphism, σ_m corresponds to the unit $m + n\mathbb{Z}$, and the order of σ_m is the least $k \in \mathbb{N}$ such that $m^k \equiv 1 \pmod{n}$. Hence $\text{Aut}(\mathbb{Z}_n)$ is cyclic iff there is a primitive root modulo n . It remains to prove the number-theoretic classification of those $n > 1$ which have primitive roots.

We use Euler's theorem in the following form: if $\gcd(a, q) = 1$, then $a^{\varphi(q)} \equiv 1 \pmod{q}$. First suppose $q = 2^s$ with $s \geq 3$. If a is odd, then $a^2 \equiv 1 \pmod{8}$. We claim that

$$a^{2^{s-2}} \equiv 1 \pmod{2^s}$$

for all $s \geq 3$. The case $s = 3$ is the congruence just observed. If $a^{2^{s-2}} = 1 + b2^s$, then

$$a^{2^{s-1}} = (1 + b2^s)^2 \equiv 1 \pmod{2^{s+1}},$$

so the claim follows by induction. But $\varphi(2^s) = 2^{s-1}$, so every odd residue modulo 2^s has order at most $2^{s-2} < \varphi(2^s)$. Thus 2^s has no primitive root for $s \geq 3$.

Next suppose $\gcd(u, v) = 1$ with $u, v > 2$. Let a be coprime to uv , and put

$$h = \text{lcm}(\varphi(u), \varphi(v)).$$

Since $u, v > 2$, both $\varphi(u)$ and $\varphi(v)$ are even (the reduced residue classes pair as b and $-b$), so

$$h \leq \frac{\varphi(u)\varphi(v)}{2} = \frac{\varphi(uv)}{2}.$$

Euler's theorem gives $a^h \equiv 1 \pmod{u}$ and $a^h \equiv 1 \pmod{v}$; since $\gcd(u, v) = 1$, this implies $a^h \equiv 1 \pmod{uv}$. Hence no unit modulo uv can have order $\varphi(uv)$, and uv has no primitive root.

These two obstructions leave only 2 , 4 , odd prime powers, and twice odd prime powers. Indeed, if

$$n = 2^a p_1^{e_1} \cdots p_t^{e_t}$$

is the prime factorization of $n > 1$, then two distinct odd prime factors are ruled out by the previous paragraph, as is $2^a p^e$ with $a \geq 2$ and p odd. The pure powers 2^a with $a \geq 3$ were ruled out in the paragraph before that. Therefore any $n > 1$ with a primitive root must be one of 2 , 4 , p^e , or $2p^e$ with p odd.

It remains to see that all numbers in this list actually have primitive roots. Clearly 1 is a primitive root modulo 2 , and 3 is a primitive root modulo 4 . Now let p be an odd prime. By Proposition 2.2, there is a primitive root r modulo p . We may choose such an r with

$$r^{p-1} \not\equiv 1 \pmod{p^2}.$$

For if a chosen primitive root r satisfies $r^{p-1} \equiv 1 \pmod{p^2}$, then $r + p$ is still primitive modulo p , and

$$(r + p)^{p-1} \equiv r^{p-1} + (p-1)pr^{p-2} \equiv 1 - pr^{p-2} \not\equiv 1 \pmod{p^2}.$$

We claim that this r is primitive modulo p^e for every $e \in \mathbb{N}$. For $e = 1$ this is how r was chosen. For $e \geq 2$, we first show by induction that

$$r^{p^{e-2}(p-1)} \not\equiv 1 \pmod{p^e}.$$

The case $e = 2$ is our choice of r . If the assertion holds for e , then Euler's theorem modulo p^{e-1} gives

$$r^{p^{e-2}(p-1)} = 1 + cp^{e-1}$$

for some integer c not divisible by p . Raising both sides to the p th power gives

$$r^{p^{e-1}(p-1)} = (1 + cp^{e-1})^p \equiv 1 + cp^e \not\equiv 1 \pmod{p^{e+1}},$$

proving the induction step. Now let d be the order of r modulo p^e . Then $d \mid \varphi(p^e) = p^{e-1}(p-1)$, and reducing modulo p shows $p-1 \mid d$. Hence $d = p^j(p-1)$ for some $0 \leq j \leq e-1$. If $j < e-1$, then $d \mid p^{e-2}(p-1)$, contradicting the congruence just proved. Thus $d = p^{e-1}(p-1) = \varphi(p^e)$, so r is primitive modulo p^e .

Finally, let r be primitive modulo p^e . Replacing r by $r + p^e$ if necessary, we may assume that r is odd; this does not change its residue modulo p^e . Let d be the order of r modulo $2p^e$. Since $r^d \equiv 1 \pmod{2p^e}$ implies $r^d \equiv 1 \pmod{p^e}$, we have $\varphi(p^e) \mid d$. Also $d \mid \varphi(2p^e) = \varphi(p^e)$. Therefore $d = \varphi(2p^e)$, and r is primitive modulo $2p^e$.

We have proved that primitive roots exist exactly for $n = 2, 4, p^e$, and $2p^e$ with p odd. Therefore $\text{Aut}(\mathbb{Z}_n)$ is cyclic exactly for the same values of n . $\square$

The number-theoretic primitive-root classification used in this proof follows [9, Section 8.3].

Fixed and Characteristic Subgroups

- Let $\varphi \in \text{Aut}(G)$ and $H \leq G$. We say H is *fixed by* φ if $\varphi(H) = H$; in this case the restriction is an automorphism of H .
- If $L \leq \text{Aut}(G)$, H is *fixed by* L if H is fixed by every $\varphi \in L$.
- H is normal in G iff H is fixed by $\text{Inn}(G)$.
- H is a *characteristic subgroup* of G (or H *characteristic* in G) if H is fixed by $\text{Aut}(G)$. (Some authors write $H \text{ char } G$.)
- Example: $Z(G)$ is always characteristic. For if $x \in Z(G)$ and $\varphi \in \text{Aut}(G)$, then $\varphi(x)y = \varphi(xy) = \varphi(yx) = \varphi(y)\varphi(x)$ for any $y \in G$, showing $\varphi(x) \in Z(G)$.
- Characteristic subgroups are normal, but the converse fails. In fact, an infinite abelian group need not have any non-trivial proper characteristic subgroups (Exercise 2.5).

Lemma 2.4. *If K is a characteristic subgroup of H and H is a characteristic subgroup of G , then K is a characteristic subgroup of G .*

Proof. Let $\varphi \in \text{Aut}(G)$. We need to show that $\varphi(K) = K$. Since H is characteristic in G , we have $\varphi(H) = H$. Hence the restriction $\varphi|_H: H \rightarrow H$ is an automorphism of H : it is still a homomorphism, it is injective because φ is injective, and it is surjective onto H because $\varphi(H) = H$.

Now K is characteristic in H , so every automorphism of H fixes K . Applying this to $\varphi|_H$, we obtain $(\varphi|_H)(K) = K$. But $K \leq H$, so $(\varphi|_H)(K) = \varphi(K)$. Therefore $\varphi(K) = K$. Since $\varphi \in \text{Aut}(G)$ was arbitrary, K is characteristic in G . $\square$

The Derived Group

- Normality is not transitive: if $N \trianglelefteq G$, the restriction to N of an element of $\text{Inn}(G)$ lies in $\text{Aut}(N)$ but need not lie in $\text{Inn}(N)$.
- For $x, y \in G$ the *commutator* is $[x, y] = xyx^{-1}y^{-1}$.
- The *derived group* G' of G is the subgroup generated by all commutators: $G' = \langle \{[x, y] \mid x, y \in G\} \rangle$.
- G is abelian iff $G' = 1$.
- If $H \leq G$, then $H' \leq G'$.
- In general, G' contains more than just the commutators of elements of G .
- Since $[x, y]^{-1} = (xyx^{-1}y^{-1})^{-1} = yxy^{-1}x^{-1} = [y, x]$, by Proposition 1.2 an arbitrary element of G' is a finite product of commutators of elements of G .

Lemma 2.5. *Let G be a group. Then G' is characteristic in G .*

Proof. Let $\varphi \in \text{Aut}(G)$. For any $x, y \in G$, we have

$$\varphi([x, y]) = \varphi(xy x^{-1} y^{-1}) = \varphi(x) \varphi(y) \varphi(x)^{-1} \varphi(y)^{-1} = [\varphi(x), \varphi(y)].$$

Thus φ sends each commutator of elements of G to another commutator of elements of G . Now let $g \in G'$. By Proposition 1.2 and the preceding observation about inverses of commutators, we may write

$$g = [x_1, y_1] \cdots [x_r, y_r]$$

for some $x_i, y_i \in G$. Applying φ , we obtain

$$\varphi(g) = [\varphi(x_1), \varphi(y_1)] \cdots [\varphi(x_r), \varphi(y_r)] \in G'.$$

Hence $\varphi(G') \leq G'$. Applying the same argument to φ^{-1} gives $\varphi^{-1}(G') \leq G'$, and therefore $G' \leq \varphi(G')$. Thus $\varphi(G') = G'$. Since φ was arbitrary, G' is characteristic in G . $\square$

Proposition 2.6. *Let G be a group, and let $N \trianglelefteq G$. Then G/N is abelian iff $G' \leq N$.*

Proof. By Lemma 2.5, G' is characteristic in G , and hence $G' \trianglelefteq G$. We first observe that G/G' is abelian. Indeed, for any $x, y \in G$,

$$[xG', yG'] = [x, y]G' = G',$$

since $[x, y] \in G'$. Thus every commutator in G/G' is trivial, so G/G' is abelian.

Suppose first that $G' \leq N$. Since $G' \leq N$, the set N/G' is a subgroup of G/G' . Moreover, it is normal in G/G' : if $g \in G$ and $n \in N$, then

$$(gG')(nG')(gG')^{-1} = gng^{-1}G' \in N/G',$$

because $N \trianglelefteq G$. Thus $N/G' \trianglelefteq G/G'$, and by Second Isomorphism Theorem,

$$G/N \cong (G/G')/(N/G').$$

Since G/G' is abelian, every quotient of G/G' is abelian. Hence G/N is abelian.

Conversely, suppose that G/N is abelian. For any $x, y \in G$, the commutator of xN and yN is trivial, so

$$N = [xN, yN] = [x, y]N.$$

Therefore $[x, y] \in N$ for all $x, y \in G$. Since N contains every commutator of elements of G , it contains the subgroup generated by all such commutators. Hence $G' \leq N$. $\square$

External Direct Products

- Let $n \in \mathbb{N}$ and let $G_1, \dots, G_n$ be groups. The Cartesian product $G_1 \times \cdots \times G_n$ becomes a group under componentwise multiplication $(g_1, \dots, g_n)(g'_1, \dots, g'_n) = (g_1g'_1, \dots, g_ng'_n)$. The identity is $(1, \dots, 1)$ and inverses are $(g_1^{-1}, \dots, g_n^{-1})$.
- This is the (*external*) *direct product* of $G_1, \dots, G_n$.
- Order of factors is irrelevant: $G_1 \times \cdots \times G_n \cong G_{\rho(1)} \times \cdots \times G_{\rho(n)}$ for any $\rho \in S_n$.
- For $G = G_1 \times \cdots \times G_n$:
 - Each $H_i = \{(1, \dots, g_i, \dots, 1) \mid g_i \in G_i\}$ is a normal subgroup of G isomorphic with G_i . Moreover G/H_i is isomorphic with the direct product of the remaining G_j .
 - Every $g \in G$ has a unique expression $g = h_1 \cdots h_n$ with $h_i \in H_i$; if $g = (g_1, \dots, g_n)$ then $h_i = (1, \dots, g_i, \dots, 1)$.
 - If the G_i are finite, then $|G| = |G_1| \cdots |G_n|$.

Internal Direct Products

- Suppose G has subgroups $H_1, \dots, H_n$ such that:
 1. $H_i \trianglelefteq G$ for each $1 \leq i \leq n$.
 2. Every $g \in G$ has a unique expression $g = h_1 \cdots h_n$ where $h_i \in H_i$.

Conditions (1) and (2) imply:

3. $G = H_1 \cdots H_n$.
 4. $H_i \cap H_1 \cdots H_{i-1} H_{i+1} \cdots H_n = 1$ for each i .
 5. If $i \neq j$, then elements of H_i commute with elements of H_j .
 6. If $g = h_1 \cdots h_n$ and $g' = h'_1 \cdots h'_n$ with $h_i, h'_i \in H_i$, then $gg' = (h_1 h'_1) \cdots (h_n h'_n)$.
- Under these conditions, there is a unique isomorphism from G to the external direct product $H_1 \times \cdots \times H_n$ sending H_i to $1 \times \cdots \times H_i \times \cdots \times 1$. Call G the (*internal*) *direct product* of $H_1, \dots, H_n$ and write $G = H_1 \times \cdots \times H_n$ (slight abuse of notation).
 - If (1) holds, then (2) holds iff both (3) and (4) hold; so to check a direct product, it suffices to verify either (1) and (2) or (1), (3), and (4). (For $n = 2$, (4) reduces to $H_1 \cap H_2 = 1$.)

Lemma 2.7. *Let G be a group having normal subgroups H and K such that $G = HK$. Then $G/H \cap K \cong H/H \cap K \times K/H \cap K$.*

Proof. Put $M = H \cap K$. By Proposition 1.7, we have $M \trianglelefteq G$, and hence G/M , H/M , and K/M are quotient groups. By Correspondence Theorem, the subgroups H/M and K/M are normal in G/M . We first compute their intersection. Suppose that $xM \in (H/M) \cap (K/M)$. Since $xM \in H/M$, there is some $h \in H$ such that $xM = hM$, and because $M \leq H$ this implies $x \in H$. Similarly, $x \in K$. Thus $x \in H \cap K = M$, and so $xM = M$. Therefore

$$(H/M) \cap (K/M) = 1.$$

Next let $gM \in G/M$. Since $G = HK$, we may write $g = hk$ with $h \in H$ and $k \in K$. Then

$$gM = hkM = (hM)(kM),$$

so $(G/M) = (H/M)(K/M)$. Hence G/M is the internal direct product of H/M and K/M , and therefore is isomorphic with the corresponding external direct product. Thus

$$G/(H \cap K) \cong H/(H \cap K) \times K/(H \cap K).$$

□

Lemma 2.8. *Let $n \in \mathbb{N}$, and write $n = p_1^{a_1} \cdots p_r^{a_r}$ where the p_i are distinct primes and the a_i are positive integers. Then we have $\mathbb{Z}_n \cong \mathbb{Z}_{p_1^{a_1}} \times \cdots \times \mathbb{Z}_{p_r^{a_r}}$.*

Proof. If $n = 1$, there is nothing to prove, so assume $n > 1$. For each i , let

$$P_i = \langle x_i \rangle \cong \mathbb{Z}_{p_i^{a_i}},$$

and put $P = P_1 \times \cdots \times P_r$. Then

$$|P| = |P_1| \cdots |P_r| = p_1^{a_1} \cdots p_r^{a_r} = n.$$

Let $x = (x_1, \dots, x_r) \in P$. For $k \in \mathbb{N}$, we have

$$x^k = (1, \dots, 1)$$

iff $x_i^k = 1$ for every i , iff $p_i^{a_i} \mid k$ for every i . Since the prime powers $p_i^{a_i}$ are pairwise coprime, this is equivalent to $n \mid k$. Hence $o(x) = n$.

Therefore $\langle x \rangle$ has n elements and lies in P , which also has n elements. Thus $\langle x \rangle = P$, so P is cyclic of order n . It follows that $P \cong \mathbb{Z}_n$, and substituting the groups $P_i \cong \mathbb{Z}_{p_i^{a_i}}$ gives

$$\mathbb{Z}_n \cong \mathbb{Z}_{p_1^{a_1}} \times \cdots \times \mathbb{Z}_{p_r^{a_r}}.$$

□

Corollary 2.9. *If $\gcd(a, b) = 1$, then $\mathbb{Z}_{ab} \cong \mathbb{Z}_a \times \mathbb{Z}_b$.*

Proof. If $a = 1$ or $b = 1$, the assertion is immediate because $\mathbb{Z}_1$ is the trivial group. Thus assume $a, b > 1$. Write

$$a = p_1^{\alpha_1} \cdots p_s^{\alpha_s}, \quad b = q_1^{\beta_1} \cdots q_t^{\beta_t}$$

as prime-power factorizations. Since $\gcd(a, b) = 1$, the primes $p_1, \dots, p_s$ are distinct from the primes $q_1, \dots, q_t$. Hence

$$ab = p_1^{\alpha_1} \cdots p_s^{\alpha_s} q_1^{\beta_1} \cdots q_t^{\beta_t}$$

is the prime-power factorization of ab . Applying Lemma 2.8 to ab gives

$$\mathbb{Z}_{ab} \cong \mathbb{Z}_{p_1^{\alpha_1}} \times \cdots \times \mathbb{Z}_{p_s^{\alpha_s}} \times \mathbb{Z}_{q_1^{\beta_1}} \times \cdots \times \mathbb{Z}_{q_t^{\beta_t}}.$$

Applying Lemma 2.8 separately to a and to b , and regrouping the factors, gives

$$\mathbb{Z}_{ab} \cong \mathbb{Z}_a \times \mathbb{Z}_b.$$

□

Proposition 2.10. *Suppose that a finite group G is the direct product of its subgroups $H_1, \dots, H_n$, where the orders $|H_i|$ are pairwise coprime. Then any subgroup L of G is the direct product of $L \cap H_1, \dots, L \cap H_n$.*

Proof. We first prove the case $n = 2$. Write $G = H \times K$, where $\gcd(|H|, |K|) = 1$, and let $L \leq G$. Put

$$A = L \cap H, \quad B = L \cap K.$$

Since $H, K \trianglelefteq G$, we have $A, B \trianglelefteq L$. Also $A \cap B \leq H \cap K = 1$, so $A \cap B = 1$. It remains to show that $L = AB$.

Let $g \in L$. Since $G = H \times K$, we may write $g = hk$ with $h \in H$ and $k \in K$, and h and k commute. By Lagrange's Theorem, $o(h) \mid |H|$ and $o(k) \mid |K|$, so $o(h)$ and $o(k)$ are coprime. Put $a = o(h)$ and $b = o(k)$. If $(hk)^m = 1$, then, since h and k commute,

$$h^m = k^{-m} \in H \cap K = 1.$$

Hence $a \mid m$ and $b \mid m$, and because $\gcd(a, b) = 1$, we have $ab \mid m$. Conversely, $(hk)^{ab} = 1$, so $o(hk) = ab$. Now $\langle hk \rangle \leq \langle h \rangle \langle k \rangle$. The subgroup $\langle h \rangle \langle k \rangle$ is the internal direct product of $\langle h \rangle$ and $\langle k \rangle$, so it has order ab . Since $\langle hk \rangle$ also has order ab , we have

$$\langle hk \rangle = \langle h \rangle \langle k \rangle.$$

Thus $h, k \in \langle hk \rangle = \langle g \rangle \leq L$. Therefore $h \in A$ and $k \in B$, and so $g = hk \in AB$. This proves $L \leq AB$, while $AB \leq L$ is immediate. Hence $L = AB$. Since A and B are normal in L , have trivial intersection, and commute with one another, $L = A \times B$.

We now argue by induction on n . The case $n = 1$ is immediate, and the case $n = 2$ has just been proved. Suppose $n > 2$ and that the result is known for direct products with $n - 1$ factors. Let

$$M = H_1 \times \cdots \times H_{n-1}.$$

Then $G = M \times H_n$, and pairwise coprimeness of the orders $|H_i|$ gives $\gcd(|M|, |H_n|) = 1$. Applying the $n = 2$ case to $G = M \times H_n$ gives

$$L = (L \cap M) \times (L \cap H_n).$$

Inside M , the subgroup $L \cap M$ is a subgroup of the direct product $H_1 \times \cdots \times H_{n-1}$, whose factor orders are still pairwise coprime. By the induction hypothesis,

$$L \cap M = (L \cap H_1) \times \cdots \times (L \cap H_{n-1}).$$

Substituting this into the preceding display gives

$$L = (L \cap H_1) \times \cdots \times (L \cap H_n),$$

as required. □

Internal Semidirect Products

- Suppose G has subgroup H and normal subgroup N with $G = NH$ and $N \cap H = 1$. Then G is the (*internal*) *semidirect product* of N by H , written $G = N \rtimes H$. (Notation common but not standard; alternatives include $N \rtimes H$, $H \ltimes N$, no notation.)
- If in addition $H \trianglelefteq G$, then G is the direct product of N and H .
- Example: $G = S_3$, $N = A_3$, $H = \langle (1\ 2) \rangle$. Then $G = N \rtimes H$, but H is not normal, so G is not the direct product.
- Properties of $G = N \rtimes H$:
 - $H \cong H/(N \cap H) \cong NH/N = G/N$ by First Isomorphism Theorem; if G is finite, $|G| = |N||G : N| = |N||H|$.
 - Each $x \in G$ has a unique expression $x = nh$ with $n \in N$, $h \in H$.
 - If $x = n_1h_1$, $y = n_2h_2$, then $xy = n_1h_1n_2h_1^{-1} \cdot h_1h_2 = n'h'$ with $n' = n_1h_1n_2h_1^{-1} \in N$ and $h' = h_1h_2 \in H$.
 - For $h \in H$, conjugation by h maps N to N , so $\varphi_h : N \rightarrow N$, $\varphi_h(n) = hnh^{-1}$, is an automorphism of N , and $\varphi_h \circ \varphi_{h'} = \varphi_{hh'}$. Hence the *conjugation homomorphism* $\varphi : H \rightarrow \text{Aut}(N)$, $\varphi(h) = \varphi_h$.
 - $(n_1h_1)(n_2h_2) = n_1\varphi(h_1)(n_2) \cdot h_1h_2$, so the multiplication of G is determined by the operations of N and H together with φ .
 - If φ is trivial, then $hnh^{-1} = n$ for all $n \in N$, $h \in H$, so $H \trianglelefteq G$ and $G = N \times H$. Conversely, if $G = N \times H$, then H commutes with N , so φ is trivial.
 - If φ is non-trivial, G is non-abelian: there exist $h \in H$, $n \in N$ with $hnh^{-1} = \varphi(h)(n) \neq n$.

External Semidirect Products

- Given groups N , H and a homomorphism $\varphi : H \rightarrow \text{Aut}(N)$, define a binary operation on $N \times H$ by

$$(n_1, h_1)(n_2, h_2) = (n_1\varphi(h_1)(n_2), h_1h_2).$$
 This makes $N \times H$ a group with identity $(1, 1)$ and inverse $(n, h)^{-1} = (\varphi(h^{-1})(n^{-1}), h^{-1})$.
- Call this the (*external*) *semidirect product* of N by H corresponding to φ , denoted $G = N \rtimes_{\varphi} H$.
- This generally differs from the direct product unless φ is trivial.
- $\mathcal{N} = N \times 1$ and $\mathcal{H} = 1 \times H$ in G are isomorphic copies of N , H respectively. Computation shows $\mathcal{N} \trianglelefteq G$, $G = \mathcal{N}\mathcal{H}$, $\mathcal{N} \cap \mathcal{H} = 1$, so G is the internal semidirect product of $\mathcal{N}$ by $\mathcal{H}$.
- Identifying $\mathcal{N}$ with N and $\mathcal{H}$ with H , the conjugation homomorphism from $\mathcal{H}$ to $\text{Aut}(\mathcal{N})$ corresponds to the original φ .
- Write $N \rtimes_{\varphi} H = \{nh \mid n \in N, h \in H\}$ with multiplication $(n_1h_1)(n_2h_2) = n_1\varphi(h_1)(n_2) \cdot h_1h_2$. Note $hnh^{-1} = \varphi(h)(n)$.
- Distinct $\varphi, \psi : H \rightarrow \text{Aut}(N)$ need not yield isomorphic semidirect products. The next two propositions are partial converses.

Proposition 2.11. *Let H be a cyclic group and let N be an arbitrary group. If φ and ψ are monomorphisms from H to $\text{Aut}(N)$ such that $\varphi(H) = \psi(H)$, then we have $N \rtimes_{\varphi} H \cong N \rtimes_{\psi} H$.*

Proof. Write $H = \langle x \rangle$. Since $\varphi(H) = \psi(H)$, the automorphisms $\varphi(x)$ and $\psi(x)$ generate the same cyclic subgroup of $\text{Aut}(N)$. Hence there exist $a, b \in \mathbb{Z}$ such that

$$\varphi(x)^a = \psi(x), \quad \psi(x)^b = \varphi(x).$$

It follows that, for every $h \in H$,

$$\varphi(h^a) = \psi(h), \quad \psi(h^b) = \varphi(h).$$

Indeed, if $h = x^m$, then $\varphi(h^a) = \varphi(x^{am}) = \varphi(x)^{am} = \psi(x)^m = \psi(h)$, and the second equality is similar. Define

$$\tau: N \rtimes_{\psi} H \rightarrow N \rtimes_{\varphi} H, \quad \tau(nh) = nh^a.$$

We show that τ is a homomorphism. Let $n_1, n_2 \in N$ and $h_1, h_2 \in H$. Multiplication in $N \rtimes_{\psi} H$ gives

$$(n_1 h_1)(n_2 h_2) = n_1 \psi(h_1)(n_2) h_1 h_2.$$

Since H is cyclic, it is abelian, so $(h_1 h_2)^a = h_1^a h_2^a$. Therefore

$$\tau((n_1 h_1)(n_2 h_2)) = n_1 \psi(h_1)(n_2)(h_1 h_2)^a = n_1 \varphi(h_1^a)(n_2) h_1^a h_2^a.$$

But this is exactly the product of $n_1 h_1^a$ and $n_2 h_2^a$ in $N \rtimes_{\varphi} H$, so

$$\tau((n_1 h_1)(n_2 h_2)) = \tau(n_1 h_1) \tau(n_2 h_2).$$

Thus τ is a homomorphism. Similarly, the map

$$\lambda: N \rtimes_{\varphi} H \rightarrow N \rtimes_{\psi} H, \quad \lambda(nh) = nh^b$$

is a homomorphism.

It remains to check that these two homomorphisms are inverse to one another. Since

$$\varphi(x) = \psi(x)^b = (\varphi(x)^a)^b = \varphi(x^{ab}),$$

and φ is injective, we have $x = x^{ab}$. Hence $h = h^{ab}$ for every $h \in H$. Similarly, using the injectivity of ψ , we get $h = h^{ba}$ for every $h \in H$. Therefore

$$(\tau \circ \lambda)(nh) = \tau(nh^b) = nh^{ba} = nh, \quad (\lambda \circ \tau)(nh) = \lambda(nh^a) = nh^{ab} = nh.$$

Thus τ and λ are mutually inverse isomorphisms, and so $N \rtimes_{\varphi} H \cong N \rtimes_{\psi} H$. $\square$

Proposition 2.12. *Let N and H be groups, let $\psi: H \rightarrow \text{Aut}(N)$ be a homomorphism, and let $f \in \text{Aut}(N)$. If $\hat{f}$ is the inner automorphism of $\text{Aut}(N)$ induced by f , then $N \rtimes_{\hat{f} \circ \psi} H \cong N \rtimes_{\psi} H$.*

Proof. By definition, the inner automorphism $\hat{f}$ of $\text{Aut}(N)$ sends $\alpha \in \text{Aut}(N)$ to $f \circ \alpha \circ f^{-1}$. Define

$$\theta: N \rtimes_{\psi} H \rightarrow N \rtimes_{\hat{f} \circ \psi} H, \quad \theta(nh) = f(n)h.$$

We show that θ is a homomorphism. Let $n_1, n_2 \in N$ and $h_1, h_2 \in H$. Multiplication in $N \rtimes_{\psi} H$ gives

$$(n_1 h_1)(n_2 h_2) = n_1 \psi(h_1)(n_2) h_1 h_2.$$

Hence

$$\theta((n_1 h_1)(n_2 h_2)) = f(n_1 \psi(h_1)(n_2)) h_1 h_2 = f(n_1) f(\psi(h_1)(n_2)) h_1 h_2.$$

On the other hand, multiplication in $N \rtimes_{\hat{f} \circ \psi} H$ gives

$$\theta(n_1 h_1) \theta(n_2 h_2) = (f(n_1) h_1)(f(n_2) h_2) = f(n_1) (\hat{f} \circ \psi)(h_1)(f(n_2)) h_1 h_2.$$

Since

$$(\hat{f} \circ \psi)(h_1)(f(n_2)) = (f \circ \psi(h_1) \circ f^{-1})(f(n_2)) = f(\psi(h_1)(n_2)),$$

the two expressions are equal. Thus θ is a homomorphism.

Finally, the map

$$N \rtimes_{\hat{f} \circ \psi} H \rightarrow N \rtimes_{\psi} H, \quad nh \mapsto f^{-1}(n)h$$

is inverse to θ . Therefore θ is an isomorphism, and so $N \rtimes_{\hat{f} \circ \psi} H \cong N \rtimes_{\psi} H$. $\square$

Dihedral Groups

- Let $N \cong \mathbb{Z}_n$ ($n \in \mathbb{N}$), $H \cong \mathbb{Z}_2$, $\varphi : H \rightarrow \text{Aut}(N)$ send the generator of H to the automorphism sending each element of N to its inverse (so $\varphi(h) = \sigma_{n-1}$).
- The group $N \rtimes_{\varphi} H$ is the *dihedral group* of order $2n$, denoted D_{2n} . (Some authors write D_n .)
- D_{2n} is non-abelian iff $n > 2$; when $n = 2$, φ is trivial and $D_4 \cong \mathbb{Z}_2 \times \mathbb{Z}_2$.
- Geometrically D_{2n} is the symmetry group of a regular n -gon: the generator of N is rotation by $2\pi/n$, the generator of H is reflection through a fixed axis.
- Also: the *infinite dihedral group* $D_{\infty} = N \rtimes_{\varphi} H$ where $N \cong \mathbb{Z}$ and H, φ as above.

Proposition 2.13. *Let s and t be elements of order 2 in a group G . (Such elements are called involutions.) Then $\langle s, t \rangle$ is a dihedral group; in particular, $\langle st \rangle \rtimes \langle s \rangle$.*

Proof. Put

$$L = \langle s, t \rangle, \quad r = st, \quad N = \langle r \rangle, \quad H = \langle s \rangle.$$

Since s and t have order 2, we have $s^{-1} = s$ and $t^{-1} = t$. Therefore

$$srs^{-1} = s(st)s = ts = (st)^{-1} = r^{-1}.$$

Similarly,

$$trt^{-1} = t(st)t = ts = r^{-1}.$$

Thus both generators s and t of L conjugate r to r^{-1} . Hence both generators normalize $N = \langle r \rangle$, and so

$$N \trianglelefteq L.$$

Also $H = \langle s \rangle \leq L$. We next show that $L = NH$. Since $r = st \in N$ and $s \in H$, we have

$$t = sr \in HN.$$

But $N \trianglelefteq L$, so $HN = NH$. Hence both s and t lie in NH , and therefore

$$L = \langle s, t \rangle \leq NH.$$

The reverse inclusion is immediate from $N, H \leq L$, so $L = NH$.

It remains to check that $N \cap H = 1$. Since $H = \langle s \rangle$ and s has order 2, the only possible non-identity element of H is s . Suppose, for contradiction, that $s \in N$. Then s is a power of r . Since $t = sr$, it follows that $t \in N$ as well. Hence $L = \langle s, t \rangle \leq N$, so $L = N$ is cyclic. But a cyclic group has at most one element of order 2, so $s = t$. Then $r = st = 1$, and hence $N = \langle r \rangle = 1$, contradicting $s \in N$. Therefore $s \notin N$, and so $N \cap H = 1$.

We have shown that

$$L = NH, \quad N \trianglelefteq L, \quad N \cap H = 1.$$

Hence L is the internal semidirect product $N \rtimes H$. Finally, the conjugation action of H on N sends the generator r of N to r^{-1} , so it sends every element of N to its inverse. This is precisely the action used in the definition of a dihedral group. Thus $\langle s, t \rangle$ is dihedral; more explicitly, if $r = st$ has finite order n , then $L \cong D_{2n}$, while if r has infinite order, then $L \cong D_{\infty}$. $\square$

Reference note. Reference [7] is Aschbacher's *Finite Group Theory*; the local reference copy is the second edition, and Section 45 is titled "Involutions in finite groups." The point of the reference is that the small fact proved above becomes a major tool once one studies finite simple groups. In a finite group, two involutions x and y generate a dihedral group whose rotation subgroup is $\langle xy \rangle$. This means that the product xy controls the subgroup $\langle x, y \rangle$, and the elementary geometry of dihedral groups gives information about conjugacy among involutions.

Aschbacher begins Section 45 by using this exact observation. If $D = \langle x, y \rangle$ and $n = o(xy)$, then D has order $2n$. When n is odd, all involutions of D are conjugate in D , so x and y are already conjugate inside the small subgroup they generate. When n is even, there is a unique involution in the rotation subgroup $\langle xy \rangle$, and this central involution gives an additional conjugacy possibility. Thus the parity of $o(xy)$ tells us how involutions can fuse, meaning how they may become conjugate in the larger group.

This is important because involutions are unavoidable in non-abelian finite simple groups: by the Feit–Thompson Odd Order Theorem, every non-abelian finite simple group has even order, and therefore has involutions. Section 45 uses dihedral subgroups generated by pairs of involutions to count and control such elements. Two central results in that section are the Brauer–Fowler Theorem and the Thompson Order Formula. The Brauer–Fowler Theorem says, roughly, that fixing the centralizer of an involution leaves only finitely many possible finite simple groups. This suggests that finite simple groups can be attacked by studying the centralizers of their involutions.

The Thompson Order Formula is more explicit. When a finite group has at least two conjugacy classes of involutions, it expresses the order of the group in terms of centralizers of involutions and the fusion of involutions inside those centralizers. In one form, if $x_1, \dots, x_k$ represent the conjugacy classes of involutions and n_i counts certain ordered pairs of involutions whose product-generated subgroup contains x_i , then

$$|G| = |C_G(x_1)| |C_G(x_2)| \sum_{i=1}^k \frac{n_i}{|C_G(x_i)|}.$$

The important point is not the formula itself at this stage, but the method behind it: because pairs of involutions generate dihedral groups, their products and conjugacy relations can be counted very precisely. Thus Proposition 2.13 is not just a cute exercise. It is one of the basic mechanisms behind the local analysis of finite simple groups, where one tries to understand a group by studying centralizers and normalizers of special subgroups.

Exercises

Exercise 2.1. Let H be a subgroup of a cyclic group G . Show that every automorphism of H is the restriction to H of an automorphism of G .

Solution. Let $G = \langle g \rangle$ and let $H \leq G$. Since every subgroup of a cyclic group is cyclic, we may write $H = \langle g^k \rangle$ for some integer k .

First suppose that G is infinite cyclic. If $H = 1$, then the only automorphism of H is the restriction of the identity automorphism of G . Otherwise $H = \langle g^k \rangle$ is infinite cyclic, and every automorphism of H sends the generator g^k either to g^k or to g^{-k} . These two automorphisms are respectively the restrictions of the automorphisms of G given by $g \mapsto g$ and $g \mapsto g^{-1}$. Hence the claim holds when G is infinite cyclic. Now suppose that G is finite of order n , and put $|H| = m$. Let $\psi \in \text{Aut}(H)$. Since g^k generates H , we have

$$\psi(g^k) = (g^k)^a$$

for some integer a with $\gcd(a, m) = 1$. We want to choose an automorphism $\varphi \in \text{Aut}(G)$ whose restriction to H is ψ . By Proposition 2.1, an automorphism of G has the form $\varphi(g) = g^b$ for some integer b with $\gcd(b, n) = 1$. For such a map,

$$\varphi(g^k) = \varphi(g)^k = (g^b)^k = g^{bk} = (g^k)^b.$$

Thus φ will agree with ψ on H provided $b \equiv a \pmod{m}$, since g^k has order m .

It remains to choose b so that $b \equiv a \pmod{m}$ and $\gcd(b, n) = 1$. Let q be the product of the distinct prime divisors of n which do not divide m ; if there are no such primes, take $q = 1$. Then $\gcd(m, q) = 1$, so by the Chinese Remainder Theorem there is an integer b such that

$$b \equiv a \pmod{m}, \quad b \equiv 1 \pmod{q}.$$

If p is a prime divisor of n and $p \mid m$, then $p \nmid a$, since $\gcd(a, m) = 1$, and hence $p \nmid b$ because $b \equiv a \pmod{m}$. If $p \mid n$ but $p \nmid m$, then $p \mid q$, so $b \equiv 1 \pmod{p}$, and again $p \nmid b$. Therefore no prime divisor of n divides b , so $\gcd(b, n) = 1$.

Define $\varphi(g) = g^b$. Then $\varphi \in \text{Aut}(G)$ by Proposition 2.1, and

$$\varphi(g^k) = (g^k)^b = (g^k)^a = \psi(g^k).$$

Since g^k generates H , it follows that $\varphi|_H = \psi$. Thus every automorphism of H is the restriction to H of an automorphism of G . $\square$

Exercise 2.2. Show that $\text{Aut}(\mathbb{Z}_8) \cong \mathbb{Z}_2 \times \mathbb{Z}_2$.

Solution. By Proposition 2.1,

$$\text{Aut}(\mathbb{Z}_8) \cong (\mathbb{Z}/8\mathbb{Z})^\times.$$

The units modulo 8 are precisely the residue classes represented by integers coprime to 8, so

$$(\mathbb{Z}/8\mathbb{Z})^\times = \{1 + 8\mathbb{Z}, 3 + 8\mathbb{Z}, 5 + 8\mathbb{Z}, 7 + 8\mathbb{Z}\}.$$

Each non-identity element has order 2, since

$$3^2 \equiv 5^2 \equiv 7^2 \equiv 1 \pmod{8}.$$

Thus $(\mathbb{Z}/8\mathbb{Z})^\times$ has four elements and every non-identity element has order 2. Therefore it is isomorphic to $\mathbb{Z}_2 \times \mathbb{Z}_2$, and hence

$$\text{Aut}(\mathbb{Z}_8) \cong \mathbb{Z}_2 \times \mathbb{Z}_2.$$

□

Exercise 2.3. Show that $\text{Aut}(\mathbb{Z}_{p^2}) \cong \mathbb{Z}_{p^2-p}$ for p a prime. (HINT: Let m be a primitive root modulo p , and show that either m or $m + p$ be a primitive root modulo p^2 .)

Solution. If $p = 2$, then Proposition 2.1 gives

$$\text{Aut}(\mathbb{Z}_4) \cong (\mathbb{Z}/4\mathbb{Z})^\times = \{1 + 4\mathbb{Z}, 3 + 4\mathbb{Z}\} \cong \mathbb{Z}_2,$$

and this is $\mathbb{Z}_{p^2-p}$. Hence we may assume that p is odd.

By Proposition 2.1,

$$\text{Aut}(\mathbb{Z}_{p^2}) \cong (\mathbb{Z}/p^2\mathbb{Z})^\times.$$

The latter group has order $\varphi(p^2) = p^2 - p = p(p-1)$. It remains to show that it has an element of order $p(p-1)$. By Proposition 2.2, there is a primitive root m modulo p , so m has order $p-1$ modulo p . Consider m as a unit modulo p^2 . If d is its order modulo p^2 , then $d \mid p(p-1)$ by Euler's theorem. Also, reducing $m^d \equiv 1 \pmod{p^2}$ modulo p gives $m^d \equiv 1 \pmod{p}$, so $p-1 \mid d$. Therefore d is either $p-1$ or $p(p-1)$. If $d = p(p-1)$, then m is already a primitive root modulo p^2 , so $(\mathbb{Z}/p^2\mathbb{Z})^\times$ is cyclic of order $p(p-1)$.

It remains to handle the case $d = p-1$. Then

$$m^{p-1} \equiv 1 \pmod{p^2}.$$

We claim that $m + p$ has order $p(p-1)$ modulo p^2 . Since $m + p \equiv m \pmod{p}$, the order of $m + p$ modulo p^2 is again divisible by $p-1$ and divides $p(p-1)$. Thus it is enough to show that $(m + p)^{p-1} \not\equiv 1 \pmod{p^2}$. By the binomial theorem,

$$(m + p)^{p-1} \equiv m^{p-1} + (p-1)pm^{p-2} \pmod{p^2},$$

since all remaining terms contain a factor p^2 . Using $m^{p-1} \equiv 1 \pmod{p^2}$, we obtain

$$(m + p)^{p-1} \equiv 1 + p(p-1)m^{p-2} \pmod{p^2}.$$

Since $p \nmid m$ and $p \nmid (p-1)$, the integer $(p-1)m^{p-2}$ is not divisible by p . Hence $p(p-1)m^{p-2}$ is divisible by p but not by p^2 , and therefore

$$(m + p)^{p-1} \not\equiv 1 \pmod{p^2}.$$

Thus $m + p$ has order $p(p-1)$ modulo p^2 . In either case, $(\mathbb{Z}/p^2\mathbb{Z})^\times$ is cyclic of order $p(p-1) = p^2 - p$. Therefore

$$\text{Aut}(\mathbb{Z}_{p^2}) \cong \mathbb{Z}_{p^2-p}.$$

□

Exercise 2.4. Show that if $H \trianglelefteq G$, then any characteristic subgroup of H is normal in G .

Solution. Let N be a characteristic subgroup of H . We want to show that $N \trianglelefteq G$. Let $g \in G$ be arbitrary, and consider the inner automorphism

$$\varphi: G \rightarrow G, \quad \varphi(x) = gxg^{-1}.$$

Since $H \trianglelefteq G$, we have

$$\varphi(H) = gHg^{-1} = H.$$

Hence the restriction $\varphi|_H: H \rightarrow H$ is an automorphism of H . Since N is characteristic in H , every automorphism of H fixes N , so

$$\varphi|_H(N) = N.$$

But $\varphi|_H(N) = gNg^{-1}$. Therefore $gNg^{-1} = N$. Since $g \in G$ was arbitrary, $N \trianglelefteq G$. $\square$

Exercise 2.5. Let F be a field, and consider F as a group under its additive operation. Show that F has no non-trivial proper characteristic subgroups.

Solution. The subgroups 0 and F are characteristic, so it remains to show that no other characteristic subgroups exist. Let H be a characteristic subgroup of the additive group of F , and suppose that H is non-trivial. Then there is some $h \in H$ with $h \neq 0$.

For each $a \in F^\times$, define

$$\varphi_a: F \rightarrow F, \quad \varphi_a(f) = af.$$

Since

$$\varphi_a(f_1 + f_2) = a(f_1 + f_2) = af_1 + af_2 = \varphi_a(f_1) + \varphi_a(f_2),$$

and $\varphi_{a^{-1}}$ is inverse to φ_a , the map φ_a is an automorphism of the additive group of F . Since H is characteristic, we have $\varphi_a(H) = H$ for every $a \in F^\times$.

Now let $f \in F$. If $f = 0$, then $f \in H$ because H is a subgroup. If $f \neq 0$, then $a = fh^{-1}$ lies in $F^\times$, and hence

$$f = (fh^{-1})h = \varphi_a(h) \in H.$$

Thus every element of F lies in H , so $H = F$. Therefore the only characteristic subgroups of the additive group of F are 0 and F , and hence F has no non-trivial proper characteristic subgroups. $\square$

Exercise 2.6. Verify the claim made on page 19 that if (1) holds, then (2) holds iff (3) and (4) hold.

Solution. Assume throughout that (1) holds, so $H_i \trianglelefteq G$ for every $1 \leq i \leq n$. Suppose first that (2) holds. Then every $g \in G$ has an expression $g = h_1 \cdots h_n$ with $h_i \in H_i$, so $G \subseteq H_1 \cdots H_n$. Conversely, since each $H_i \leq G$, every product $h_1 \cdots h_n$ lies in G . Hence $H_1 \cdots H_n \subseteq G$, and therefore

$$G = H_1 \cdots H_n.$$

Thus (3) holds.

To prove (4), fix i and take

$$x \in H_i \cap H_1 \cdots H_{i-1} H_{i+1} \cdots H_n.$$

Since $x \in H_i$, we have one expression

$$x = 1 \cdots 1 \cdot x \cdot 1 \cdots 1,$$

where x appears in the i th position. Since $x \in H_1 \cdots H_{i-1} H_{i+1} \cdots H_n$, we also have an expression

$$x = h_1 \cdots h_{i-1} \cdot 1 \cdot h_{i+1} \cdots h_n$$

with $h_j \in H_j$ for $j \neq i$. By uniqueness in (2), the i th factors in these two expressions are equal, so $x = 1$. Hence

$$H_i \cap H_1 \cdots H_{i-1} H_{i+1} \cdots H_n = 1$$

for every i , proving (4).

Conversely, suppose that (3) and (4) hold. By (3), every $g \in G$ has at least one expression $g = h_1 \cdots h_n$ with $h_i \in H_i$. It remains to show that this expression is unique. We first observe that elements from different factors commute. Let $i \neq j$, and take $h_i \in H_i$, $h_j \in H_j$. Since $H_j \trianglelefteq G$, we have $h_i h_j h_i^{-1} \in H_j$, and hence

$$[h_i, h_j] = (h_i h_j h_i^{-1}) h_j^{-1} \in H_j.$$

Similarly, since $H_i \trianglelefteq G$, we have $h_j h_i^{-1} h_j^{-1} \in H_i$, and hence

$$[h_i, h_j] = h_i(h_j h_i^{-1} h_j^{-1}) \in H_i.$$

Therefore $[h_i, h_j] \in H_i \cap H_j$. But $H_j \leq H_1 \cdots H_{i-1} H_{i+1} \cdots H_n$, so (4) gives $H_i \cap H_j = 1$. Thus $[h_i, h_j] = 1$, and hence h_i and h_j commute.

Now suppose

$$g = h_1 \cdots h_n = h'_1 \cdots h'_n, \quad h_i, h'_i \in H_i.$$

Fix i . Since elements from different factors commute, we may write

$$g = h_i k_i = h'_i k'_i,$$

where

$$k_i \in H_1 \cdots H_{i-1} H_{i+1} \cdots H_n, \quad k'_i \in H_1 \cdots H_{i-1} H_{i+1} \cdots H_n.$$

Since the factors in $H_1 \cdots H_{i-1} H_{i+1} \cdots H_n$ commute with one another, this product is closed under multiplication and inverses. Also, from $h_i k_i = h'_i k'_i$ we get

$$h_i'^{-1} h_i = k'_i k_i^{-1}.$$

The left hand side lies in H_i , and the right hand side lies in $H_1 \cdots H_{i-1} H_{i+1} \cdots H_n$. Therefore, by (4),

$$h_i'^{-1} h_i = 1.$$

Hence $h_i = h'_i$. Since this holds for every i , the expression of g is unique. Thus (2) holds. $\square$

Exercise 2.7. Verify the claim made on page 19 that (1) and (2) together imply (3) through (6).

Solution. Assume that (1) and (2) hold. By Exercise 2.6, conditions (1) and (2) imply (3) and (4). It remains to prove (5) and (6).

To prove (5), let $i \neq j$, and take $h_i \in H_i$, $h_j \in H_j$. Since $H_j \trianglelefteq G$, we have $h_i h_j h_i^{-1} \in H_j$, and hence

$$[h_i, h_j] = (h_i h_j h_i^{-1}) h_j^{-1} \in H_j.$$

Similarly, since $H_i \trianglelefteq G$, we have $h_j h_i^{-1} h_j^{-1} \in H_i$, and hence

$$[h_i, h_j] = h_i(h_j h_i^{-1} h_j^{-1}) \in H_i.$$

Thus $[h_i, h_j] \in H_i \cap H_j$. But $H_j \leq H_1 \cdots H_{i-1} H_{i+1} \cdots H_n$, so (4) gives $H_i \cap H_j = 1$. Hence $[h_i, h_j] = 1$, and therefore h_i and h_j commute. Since $i \neq j$ was arbitrary, elements of distinct factors commute.

Finally, let

$$g = h_1 \cdots h_n, \quad g' = h'_1 \cdots h'_n,$$

where $h_i, h'_i \in H_i$ for each i . By (5), elements from distinct factors commute, so we may move each h'_i left across all factors h_j with $j \neq i$. Therefore

$$gg' = h_1 \cdots h_n h'_1 \cdots h'_n = (h_1 h'_1) \cdots (h_n h'_n).$$

This proves (6), and hence (1) and (2) together imply (3) through (6). $\square$

Exercise 2.8. Let G_1 and G_2 be groups, and let $H \leq G_1 \times G_2$. Define

$$P_1 = \{x \in G_1 \mid (x, y) \in H \text{ for some } y \in G_2\},$$

$$I_1 = \{x \in G_1 \mid (x, 1) \in H\},$$

and analogously define subsets P_2 and I_2 of G_2 .

(a) Show for $i = 1, 2$ that $P_i \leq G_i$ and $I_i \leq P_i$.

(b) Show that there is a unique isomorphism from P_1/I_1 to P_2/I_2 sending xI_1 to yI_2 , where y is any element of G_2 such that $(x, y) \in H$.

- (c) Prove *Goursat's theorem*: There is a bijective correspondence between subgroups of $G_1 \times G_2$ and triples (S_1, S_2, φ) , where S_i is a section of G_i ($i = 1, 2$) and $\varphi : S_1 \rightarrow S_2$ is an isomorphism. (Recall that a section of a group G is a group L/M , where $M \trianglelefteq L \leq G$.)

Solution.

- (a) Fix $i \in \{1, 2\}$, and let j be the other index. First we show that $P_i \leq G_i$. Since $(1, 1) \in H$, we have $1 \in P_i$, so P_i is non-empty. Let $a, b \in P_i$. By definition of P_i , there are elements $a_j, b_j \in G_j$ such that

$$(a_1, a_2), (b_1, b_2) \in H, \quad a_i = a, \quad b_i = b.$$

Since $H \leq G_1 \times G_2$, we have

$$(a_1, a_2)(b_1, b_2)^{-1} = (a_1 b_1^{-1}, a_2 b_2^{-1}) \in H.$$

The i th coordinate of this element is ab^{-1} , so $ab^{-1} \in P_i$. Hence $P_i \leq G_i$.

Next we show that $I_i \trianglelefteq P_i$. First, $I_i \leq P_i$: if $x \in I_i$, then the element with x in the i th coordinate and 1 in the j th coordinate lies in H , so $x \in P_i$; and the same subgroup test as above, with the j th coordinate always equal to 1, shows that $I_i \leq G_i$.

It remains to prove normality in P_i . Let $a \in P_i$ and $x \in I_i$. Since $a \in P_i$, there is some $a_j \in G_j$ such that, putting $a_i = a$, we have

$$(a_1, a_2) \in H.$$

Since $x \in I_i$, putting $x_i = x$ and $x_j = 1$, we have

$$(x_1, x_2) \in H.$$

Since H is a subgroup,

$$(a_1, a_2)(x_1, x_2)(a_1, a_2)^{-1} = (a_1 x_1 a_1^{-1}, a_2 x_2 a_2^{-1}) \in H.$$

The i th coordinate of this element is axa^{-1} , and the j th coordinate is $a_j 1 a_j^{-1} = 1$. Hence $axa^{-1} \in I_i$. Therefore $I_i \trianglelefteq P_i$.

- (b) Define

$$\varphi: P_1/I_1 \rightarrow P_2/I_2$$

by setting $\varphi(xI_1) = yI_2$, where $y \in G_2$ is chosen so that $(x, y) \in H$. We first check that this is well-defined. Suppose $(x, y), (x, y') \in H$. Since H is a subgroup,

$$(x, y)^{-1}(x, y') = (1, y^{-1}y') \in H.$$

Hence $y^{-1}y' \in I_2$, so $yI_2 = y'I_2$. Thus the choice of y does not matter.

Now suppose $xI_1 = x'I_1$, and choose $y, y' \in G_2$ such that $(x, y), (x', y') \in H$. Since $xI_1 = x'I_1$, we have $x^{-1}x' \in I_1$, so

$$(x^{-1}x', 1) \in H.$$

Also,

$$(x, y)^{-1}(x', y') = (x^{-1}x', y^{-1}y') \in H.$$

Hence

$$(x^{-1}x', 1)^{-1}(x^{-1}x', y^{-1}y') = (1, y^{-1}y') \in H.$$

Therefore $y^{-1}y' \in I_2$, so $yI_2 = y'I_2$. Thus φ is well-defined on cosets.

Next, φ is a homomorphism. Let $(x, y), (x', y') \in H$. Then $(xx', yy') \in H$, and hence

$$\varphi(xI_1 \cdot x'I_1) = \varphi(xx'I_1) = yy'I_2 = yI_2 \cdot y'I_2 = \varphi(xI_1)\varphi(x'I_1).$$

Define

$$\psi: P_2/I_2 \rightarrow P_1/I_1$$

by setting $\psi(yI_2) = xI_1$, where $x \in G_1$ is chosen so that $(x, y) \in H$. By the same argument with the two coordinates interchanged, ψ is a well-defined homomorphism. Moreover, if $(x, y) \in H$, then

$$\psi(\varphi(xI_1)) = \psi(yI_2) = xI_1, \quad \varphi(\psi(yI_2)) = \varphi(xI_1) = yI_2.$$

Thus $\psi \circ \varphi$ and $\varphi \circ \psi$ are the identity maps, so φ is a bijection. Therefore φ is an isomorphism. Finally, suppose $\varphi': P_1/I_1 \rightarrow P_2/I_2$ is another function satisfying the same rule: whenever $(x, y) \in H$, we have $\varphi'(xI_1) = yI_2$. Every coset of P_1/I_1 has the form xI_1 with $x \in P_1$, and by definition of P_1 there is some $y \in G_2$ such that $(x, y) \in H$. Hence

$$\varphi'(xI_1) = yI_2 = \varphi(xI_1).$$

Therefore $\varphi' = \varphi$, proving uniqueness.

(c) Let

$$A = \{H \mid H \leq G_1 \times G_2\},$$

and let

$$B = \{(S_1, S_2, \varphi) \mid S_i \text{ is a section of } G_i, \varphi: S_1 \rightarrow S_2 \text{ is an isomorphism}\}.$$

Define

$$\theta: A \rightarrow B$$

as follows. Given $H \in A$, let P_i and I_i be the subgroups defined above. By (a), $I_i \trianglelefteq P_i \leq G_i$, so P_i/I_i is a section of G_i . By (b), there is a unique isomorphism

$$\varphi_H: P_1/I_1 \rightarrow P_2/I_2$$

such that $\varphi_H(xI_1) = yI_2$ whenever $(x, y) \in H$. Thus we define

$$\theta(H) = (P_1/I_1, P_2/I_2, \varphi_H).$$

Define

$$\theta': B \rightarrow A$$

as follows. Let $(S_1, S_2, \varphi) \in B$. Since S_i is a section of G_i , there are subgroups $M_i \trianglelefteq L_i \leq G_i$ such that

$$S_i = L_i/M_i.$$

Define

$$H_\varphi = \{(x, y) \in L_1 \times L_2 \mid \varphi(xM_1) = yM_2\}.$$

Let $(x, y), (x', y') \in H_\varphi$. Since φ is an isomorphism,

$$\varphi(xM_1) = yM_2, \quad \varphi(x'M_1) = y'M_2,$$

and hence

$$\varphi(xx'^{-1}M_1) = \varphi(xM_1)\varphi(x'M_1)^{-1} = (yM_2)(y'M_2)^{-1} = yy'^{-1}M_2.$$

Therefore $(xx'^{-1}, yy'^{-1}) \in H_\varphi$, so

$$H_\varphi \leq L_1 \times L_2 \leq G_1 \times G_2.$$

Thus $H_\varphi \in A$, and we define

$$\theta'(S_1, S_2, \varphi) = H_\varphi.$$

We now show that θ and θ' are inverse maps. First let $H \in A$, and form $\theta(H) = (P_1/I_1, P_2/I_2, \varphi_H)$. Then

$$\theta'(\theta(H)) = \{(x, y) \in P_1 \times P_2 \mid \varphi_H(xI_1) = yI_2\}.$$

If $(x, y) \in H$, then $\varphi_H(xI_1) = yI_2$, so $(x, y) \in \theta'(\theta(H))$. Conversely, suppose $(x, y) \in \theta'(\theta(H))$. Since $x \in P_1$, there is some $z \in G_2$ such that $(x, z) \in H$. Then $\varphi_H(xI_1) = zI_2$, while $\varphi_H(xI_1) = yI_2$ by assumption. Hence $zI_2 = yI_2$, so $z^{-1}y \in I_2$ and therefore $(1, z^{-1}y) \in H$. Thus

$$(x, z)(1, z^{-1}y) = (x, y) \in H.$$

Hence $\theta'(\theta(H)) = H$.

Conversely, let $(S_1, S_2, \varphi) \in B$, and write $S_i = L_i/M_i$ with $M_i \trianglelefteq L_i \leq G_i$. Form $H_\varphi = \theta'(S_1, S_2, \varphi)$. We compute the subgroups associated to H_φ . If $(x, y) \in H_\varphi$, then $x \in L_1$, so $P_1 \leq L_1$. Conversely, if $x \in L_1$, then $\varphi(xM_1) = yM_2$ for some $y \in L_2$, and so $(x, y) \in H_\varphi$. Thus $P_1 = L_1$. By a similar

argument, using the surjectivity of φ , we have $P_2 = L_2$.

For I_1 , we have

$$x \in I_1 \iff (x, 1) \in H_\varphi \iff \varphi(xM_1) = M_2 \iff xM_1 = M_1 \iff x \in M_1,$$

so $I_1 = M_1$. Similarly, $I_2 = M_2$. Therefore the first two entries recovered by θ are

$$P_1/I_1 = L_1/M_1 = S_1, \quad P_2/I_2 = L_2/M_2 = S_2.$$

Finally, the isomorphism recovered by (b) sends xM_1 to yM_2 whenever $(x, y) \in H_\varphi$. By the definition of H_φ , this means exactly that $yM_2 = \varphi(xM_1)$. Hence, by the uniqueness in (b), the recovered isomorphism is φ . Thus $\theta(\theta'(S_1, S_2, \varphi)) = (S_1, S_2, \varphi)$. Therefore θ and θ' are inverse maps, so θ is a bijection. □

Exercise 2.9. (cont.) Use Exercise 2.8 to give a different proof of Proposition 2.10.

Solution. Suppose that

$$G = H_1 \times \cdots \times H_n$$

where the orders $|H_i|$ are pairwise coprime. We want to show that any subgroup $L \leq G$ is the direct product of $L \cap H_1, \dots, L \cap H_n$. We prove this by induction on n .

First suppose $n = 2$, and write $G = H \times K$ with $\gcd(|H|, |K|) = 1$. Let $L \leq H \times K$. By Exercise 2.8, the subgroup L corresponds to a triple

$$(P_H/I_H, P_K/I_K, \varphi),$$

where P_H/I_H is a section of H , P_K/I_K is a section of K , and

$$\varphi: P_H/I_H \rightarrow P_K/I_K$$

is an isomorphism. Hence

$$|P_H/I_H| = |P_K/I_K|.$$

Since $I_H \trianglelefteq P_H \leq H$, we have $|P_H/I_H| \mid |P_H|$ and $|P_H| \mid |H|$, so $|P_H/I_H| \mid |H|$. Similarly, $|P_K/I_K| \mid |K|$. But $\gcd(|H|, |K|) = 1$, and the two quotient groups have the same order, so

$$|P_H/I_H| = |P_K/I_K| = 1.$$

Therefore P_H/I_H and P_K/I_K are trivial, and hence $P_H = I_H$ and $P_K = I_K$. By the reconstruction in Exercise 2.8, we have

$$L = \{(h, k) \in P_H \times P_K \mid \varphi(hI_H) = kI_K\}.$$

Since both quotient groups are trivial, the condition $\varphi(hI_H) = kI_K$ is automatic, so

$$L = P_H \times P_K.$$

Also $I_H = L \cap H$ and $I_K = L \cap K$, so $P_H = L \cap H$ and $P_K = L \cap K$. Thus

$$L = (L \cap H) \times (L \cap K),$$

proving the case $n = 2$.

Now suppose $n > 2$ and that the result is known for direct products with $n - 1$ factors. Put

$$M = H_1 \times \cdots \times H_{n-1}.$$

Then $G = M \times H_n$. Since the orders $|H_i|$ are pairwise coprime, we have $\gcd(|M|, |H_n|) = 1$. Applying the $n = 2$ case to $G = M \times H_n$ gives

$$L = (L \cap M) \times (L \cap H_n).$$

Now $L \cap M$ is a subgroup of $M = H_1 \times \cdots \times H_{n-1}$, and the orders $|H_1|, \dots, |H_{n-1}|$ are still pairwise coprime. By the induction hypothesis,

$$L \cap M = (L \cap H_1) \times \cdots \times (L \cap H_{n-1}).$$

Substituting this into the preceding expression gives

$$L = (L \cap H_1) \times \cdots \times (L \cap H_n).$$

This proves the result for all n . □

Exercise 2.10. Let $G = N \rtimes H$, and suppose that $N \leq K \leq G$. Show that $K = N \rtimes (K \cap H)$.

Solution. We check the internal semidirect product conditions inside K . Since $G = N \rtimes H$, we have $N \trianglelefteq G$, $G = NH$, and $N \cap H = 1$. Since $N \trianglelefteq G$ and $N \leq K \leq G$, it follows that $N \trianglelefteq K$. Also, since K and H are subgroups of G , their intersection $K \cap H$ is a subgroup of K .

Next,

$$N \cap (K \cap H) = K \cap (N \cap H) = K \cap 1 = 1.$$

It remains to show that $K = N(K \cap H)$. First, since $N \leq K$ and $K \cap H \leq K$, and since K is a subgroup, we have

$$N(K \cap H) \leq K.$$

Conversely, take $k \in K$. Since $G = N \rtimes H$, there are $n \in N$ and $h \in H$ such that

$$k = nh.$$

Since $N \leq K$, we have $n \in K$. Also $k \in K$, so

$$h = n^{-1}k \in K.$$

Since $h \in H$, we have $h \in K \cap H$. Therefore

$$k = nh \in N(K \cap H).$$

Hence $K \leq N(K \cap H)$, and so $K = N(K \cap H)$. Thus K is the internal semidirect product

$$K = N \rtimes (K \cap H).$$

□

Further Exercises

Let N and H be groups. An *extension* of N by H is a group E along with a monomorphism $i : N \rightarrow E$ and an epimorphism $\pi : E \rightarrow H$ such that $i(N) = \ker \pi$ (so that N imbeds in E as a normal subgroup, with the quotient group being isomorphic with H). We shall usually refer to an extension (E, i, π) simply by the group E ; however, the nature of the maps i and π are important in distinguishing between extensions. We identify N with its image under i , and H with the quotient of E by N . As an example, let $\varphi : H \rightarrow \text{Aut}(N)$ be a homomorphism; then the semidirect product $N \rtimes_{\varphi} H$ is an extension of N by H in an obvious way, taking i to be the inclusion map sending $n \in N$ to $(n, 1)$ and π to be the projection map sending (n, h) to h .

Exercise 2.11. We say that an extension E of N by H is a *split extension* if there is a homomorphism $t : H \rightarrow E$ (called a *splitting map* for the extension) such that $\pi \circ t$ is the identity map on H , in which case $t(H)$ will be a transversal for N in E . Show that E is a split extension iff it is a semidirect product of N by H .

Solution. ($\Rightarrow$) Suppose that E is a split extension of N by H . Thus there are homomorphisms

$$i : N \rightarrow E, \quad \pi : E \rightarrow H, \quad t : H \rightarrow E$$

such that i is injective, π is surjective, $i(N) = \ker \pi$, and $\pi \circ t = \text{id}_H$. Following the convention above, identify N with $i(N)$. Then

$$N = \ker \pi \trianglelefteq E.$$

Also $t(H) \leq E$, since t is a homomorphism. We show that

$$E = N \rtimes t(H).$$

First let $x \in N \cap t(H)$. Since $x \in t(H)$, we may write $x = t(h)$ for some $h \in H$. Since $x \in N = \ker \pi$, we have

$$1 = \pi(x) = \pi(t(h)) = (\pi \circ t)(h) = h.$$

Hence $x = t(1) = 1$, so $N \cap t(H) = 1$.

Next let $e \in E$, and put $h = \pi(e)$. Then

$$\pi(et(h)^{-1}) = \pi(e)\pi(t(h))^{-1} = h(\pi \circ t)(h)^{-1} = hh^{-1} = 1.$$

Therefore $et(h)^{-1} \in \ker \pi = N$. Say $et(h)^{-1} = n$ with $n \in N$. Then

$$e = nt(h) \in Nt(H),$$

so $E \leq Nt(H)$. Since $N \leq E$ and $t(H) \leq E$, the reverse inclusion is clear, and hence $E = Nt(H)$. Thus $E = N \rtimes t(H)$. Finally, $t(H) \cong H$ because $\pi \circ t = \text{id}_H$ forces t to be injective. Therefore E is a semidirect product of N by H .

($\Leftarrow$) Conversely, suppose that E is a semidirect product of N by H . Thus, after identifying N and H with their copies inside E , we have

$$E = N \rtimes H.$$

Hence $N \trianglelefteq E$, $H \leq E$, $E = NH$, and $N \cap H = 1$. Every element $e \in E$ has a unique expression $e = nh$ with $n \in N$ and $h \in H$. Define

$$i: N \rightarrow E, \quad i(n) = n,$$

and

$$\pi: E \rightarrow H, \quad \pi(nh) = h.$$

The map i is the inclusion map, so it is injective. We check that π is a homomorphism. Let $e_1 = n_1h_1$ and $e_2 = n_2h_2$. Since $N \trianglelefteq E$, we have $h_1n_2h_1^{-1} \in N$, and hence

$$e_1e_2 = n_1h_1n_2h_2 = n_1(h_1n_2h_1^{-1})h_1h_2.$$

Therefore the H -part of e_1e_2 is h_1h_2 , so

$$\pi(e_1e_2) = h_1h_2 = \pi(e_1)\pi(e_2).$$

Thus π is a homomorphism. It is surjective because $\pi(h) = h$ for all $h \in H$, and

$$\ker \pi = \{nh \in E \mid h = 1\} = N.$$

Hence E is an extension of N by H . Now define

$$t: H \rightarrow E, \quad t(h) = h,$$

the inclusion of H into E . Then t is a homomorphism, and for every $h \in H$,

$$(\pi \circ t)(h) = \pi(h) = h.$$

Therefore $\pi \circ t = \text{id}_H$, so the extension splits. $\square$

Exercise 2.12. (cont.) Let Q be the quaternion group of order 8. (We can consider Q as the set $\{\pm 1, \pm i, \pm j, \pm k\}$ with multiplication given by the rules $i^2 = j^2 = k^2 = -1$ and $ij = k = -ji$.) Show that Q can be realized as a non-trivial extension in four ways—thrice as an extension of $\mathbb{Z}_4$ by $\mathbb{Z}_2$, and once as an extension of $\mathbb{Z}_2$ by $\mathbb{Z}_2 \times \mathbb{Z}_2$ —but that none of these extensions is split. (In other words, Q cannot be written non-trivially as a semidirect product.)

Solution. From Exercise 2.11, an extension of N by H splits exactly when Q can be written as

$$Q = N \rtimes t(H)$$

for some subgroup $t(H) \leq Q$ with $t(H) \cong H$ and $N \cap t(H) = 1$. We first exhibit the four extensions. For the extensions of $\mathbb{Z}_4$ by $\mathbb{Z}_2$, take

$$N = \langle i \rangle, \quad N = \langle j \rangle, \quad N = \langle k \rangle.$$

Each of these subgroups has order 4, since

$$\langle i \rangle = \{1, i, -1, -i\},$$

and similarly for j and k . Hence each is isomorphic to $\mathbb{Z}_4$. Also each has index 2 in Q , so each is normal in Q . Thus for each of these choices of N ,

$$Q/N \cong \mathbb{Z}_2.$$

Therefore Q is an extension of $\mathbb{Z}_4$ by $\mathbb{Z}_2$ in three ways.

For the extension of $\mathbb{Z}_2$ by $\mathbb{Z}_2 \times \mathbb{Z}_2$, take

$$N = Z(Q) = \{1, -1\}.$$

Indeed, -1 commutes with every element of Q , while i, j, k do not commute with one another, for example $ij = k$ but $ji = -k$. The same is true for $-i, -j, -k$, so $Z(Q) = \{1, -1\}$. Thus $N \cong \mathbb{Z}_2$ and $N \trianglelefteq Q$. The quotient Q/N has four cosets:

$$N, \quad iN, \quad jN, \quad kN.$$

Moreover,

$$(iN)^2 = i^2N = (-1)N = N,$$

and the same argument gives $(jN)^2 = N$ and $(kN)^2 = N$. Hence every non-identity element of Q/N has order 2. Therefore

$$Q/N \cong \mathbb{Z}_2 \times \mathbb{Z}_2.$$

Thus Q is an extension of $\mathbb{Z}_2$ by $\mathbb{Z}_2 \times \mathbb{Z}_2$.

It remains to show that none of these extensions splits. The only element of order 2 in Q is -1 : the elements $\pm i, \pm j, \pm k$ all have order 4, and 1 has order 1.

First consider one of the three extensions with

$$N \in \{\langle i \rangle, \langle j \rangle, \langle k \rangle\}.$$

If the extension split, then by Exercise 2.11 there would be a subgroup

$$t(H) \cong \mathbb{Z}_2$$

such that $N \cap t(H) = 1$. But the only subgroup of Q isomorphic to $\mathbb{Z}_2$ is

$$\langle -1 \rangle = \{1, -1\}.$$

Since $-1 \in N$ for each of $N = \langle i \rangle, \langle j \rangle, \langle k \rangle$, we would have

$$N \cap t(H) = \{1, -1\} \neq 1,$$

a contradiction. Hence none of the three extensions of $\mathbb{Z}_4$ by $\mathbb{Z}_2$ splits.

Now consider the extension with $N = Z(Q) = \{1, -1\}$. If this extension split, then there would be a subgroup

$$t(H) \cong \mathbb{Z}_2 \times \mathbb{Z}_2$$

of Q . But $\mathbb{Z}_2 \times \mathbb{Z}_2$ has three non-identity elements, all of order 2, while Q has only one element of order 2, namely -1 . Thus Q has no subgroup isomorphic to $\mathbb{Z}_2 \times \mathbb{Z}_2$. This is again a contradiction. Therefore this extension also does not split.

Hence Q has the four stated non-trivial extensions, but none is split. Equivalently, Q cannot be written non-trivially as a semidirect product. $\square$

If E is an extension of N by H , then we cannot expect to find a homomorphism $t : H \rightarrow E$ such that $t(H)$ will be a transversal for N in E , for if such a t existed then E would be split. However, since $H \cong E/N$, we can always find a set map $t : H \rightarrow E$ whose image is a transversal for N ; such a map is called a *section* of the extension. Moreover, we can always choose t so that $t(1) = 1$, in which case we say that t is *normalized*. (We use normalized sections instead of arbitrary sections in order to keep the notational complexity to a minimum.)

Exercise 2.13. (cont.) Let t be a normalized section of an extension E . Let $\Psi : E \rightarrow \text{Aut}(E)$ be the homomorphism sending an element of E to the corresponding inner automorphism of E . We shall, for $x \in E$, regard $\Psi(x)$ as being an automorphism of N , which is possible since $N \trianglelefteq E$. Define set maps $f : H \times H \rightarrow N$ and $\varphi : H \rightarrow \text{Aut}(N)$ by

$$f(\alpha, \beta) = t(\alpha)t(\beta)t(\alpha\beta)^{-1},$$

$$\varphi(\alpha) = \Psi(t(\alpha)).$$

We call (f, φ) the *factor pair* arising from t . Show that (f, φ) has the following properties:

1 $f(\alpha, 1) = f(1, \alpha) = 1$ for every $\alpha \in H$, and $\varphi(1)$ is the identity in $\text{Aut}(N)$.

2 $\varphi(\alpha)\varphi(\beta) = \Psi(f(\alpha, \beta))\varphi(\alpha\beta)$ for $\alpha, \beta \in H$.

3 $f(\alpha, \beta)f(\alpha\beta, \gamma) = \varphi(\alpha)(f(\beta, \gamma))f(\alpha, \beta\gamma)$ for $\alpha, \beta, \gamma \in H$.

Solution. Let $\pi: E \rightarrow H$ be the epimorphism of the extension, and identify N with $\ker \pi$. Since t is a section, we have

$$\pi(t(\alpha)) = \alpha$$

for every $\alpha \in H$. First note that

$$\pi(f(\alpha, \beta)) = \pi(t(\alpha)t(\beta)t(\alpha\beta)^{-1}) = \alpha\beta(\alpha\beta)^{-1} = 1,$$

so $f(\alpha, \beta) \in \ker \pi = N$. Also, since $N \trianglelefteq E$, conjugation by any element of E restricts to an automorphism of N . Thus $\varphi(\alpha) = \Psi(t(\alpha))$ is indeed an element of $\text{Aut}(N)$.

1 Since t is normalized, $t(1) = 1$. Hence

$$f(\alpha, 1) = t(\alpha)t(1)t(\alpha \cdot 1)^{-1} = t(\alpha)t(\alpha)^{-1} = 1,$$

and similarly

$$f(1, \alpha) = t(1)t(\alpha)t(1 \cdot \alpha)^{-1} = t(\alpha)t(\alpha)^{-1} = 1.$$

Moreover,

$$\varphi(1) = \Psi(t(1)) = \Psi(1).$$

For every $n \in N$, we have

$$\Psi(1)(n) = 1n1^{-1} = n,$$

so $\varphi(1) = \text{id}_N$.

2 By the definition of f , we have

$$t(\alpha)t(\beta) = f(\alpha, \beta)t(\alpha\beta).$$

Since Ψ is a homomorphism,

$$\varphi(\alpha)\varphi(\beta) = \Psi(t(\alpha))\Psi(t(\beta)) = \Psi(t(\alpha)t(\beta)).$$

Using the displayed identity, this becomes

$$\Psi(t(\alpha)t(\beta)) = \Psi(f(\alpha, \beta)t(\alpha\beta)) = \Psi(f(\alpha, \beta))\Psi(t(\alpha\beta)).$$

Therefore

$$\varphi(\alpha)\varphi(\beta) = \Psi(f(\alpha, \beta))\varphi(\alpha\beta).$$

3 Again using

$$t(\alpha)t(\beta) = f(\alpha, \beta)t(\alpha\beta),$$

we compute

$$(t(\alpha)t(\beta))t(\gamma) = f(\alpha, \beta)t(\alpha\beta)t(\gamma) = f(\alpha, \beta)f(\alpha\beta, \gamma)t(\alpha\beta\gamma).$$

On the other hand,

$$t(\alpha)(t(\beta)t(\gamma)) = t(\alpha)f(\beta, \gamma)t(\beta\gamma).$$

Since

$$\varphi(\alpha)(n) = t(\alpha)nt(\alpha)^{-1}$$

for every $n \in N$, we have

$$t(\alpha)f(\beta, \gamma) = \varphi(\alpha)(f(\beta, \gamma))t(\alpha).$$

Thus

$$t(\alpha)(t(\beta)t(\gamma)) = \varphi(\alpha)(f(\beta, \gamma))t(\alpha)t(\beta\gamma) = \varphi(\alpha)(f(\beta, \gamma))f(\alpha, \beta\gamma)t(\alpha\beta\gamma).$$

By associativity in E ,

$$(t(\alpha)t(\beta))t(\gamma) = t(\alpha)(t(\beta)t(\gamma)).$$

Comparing the two expressions and cancelling $t(\alpha\beta\gamma)$ on the right, we obtain

$$f(\alpha, \beta)f(\alpha\beta, \gamma) = \varphi(\alpha)(f(\beta, \gamma))f(\alpha, \beta\gamma).$$

□

Exercise 2.14. (cont.) Just as we were able to externalize the notion of semidirect product, so should we be able to externalize the notion of extension; that is, given groups N and H and appropriate additional data, we should be able to construct an extension of N by H . Using Exercise 2.13 as a guide, formulate such an external construction and prove that it works.

Solution. Suppose that we are given groups N and H , together with set maps

$$f: H \times H \rightarrow N, \quad \varphi: H \rightarrow \text{Aut}(N),$$

satisfying the three identities from Exercise 2.13. Thus, for all $\alpha, \beta, \gamma \in H$,

$$f(\alpha, 1) = f(1, \alpha) = 1, \quad \varphi(1) = \text{id}_N,$$

$$\varphi(\alpha)\varphi(\beta) = \Psi(f(\alpha, \beta))\varphi(\alpha\beta),$$

and

$$f(\alpha, \beta)f(\alpha\beta, \gamma) = \varphi(\alpha)(f(\beta, \gamma))f(\alpha, \beta\gamma),$$

where, for $a \in N$, $\Psi(a)$ denotes conjugation by a on N . We construct a group E whose underlying set is

$$E = N \times H.$$

If $(n, \alpha), (m, \beta) \in N \times H$, where $n, m \in N$ and $\alpha, \beta \in H$, define

$$(n, \alpha)(m, \beta) = (n\varphi(\alpha)(m)f(\alpha, \beta), \alpha\beta).$$

This is a well-defined element of $N \times H$, since $\varphi(\alpha)(m) \in N$, $f(\alpha, \beta) \in N$, and $\alpha\beta \in H$.

We first check that this multiplication makes E into a group. The identity element is $(1, 1)$. Indeed, for $n \in N$ and $\alpha \in H$,

$$(1, 1)(n, \alpha) = (\varphi(1)(n)f(1, \alpha), \alpha) = (n, \alpha),$$

and

$$(n, \alpha)(1, 1) = (n\varphi(\alpha)(1)f(\alpha, 1), \alpha) = (n, \alpha).$$

Next we prove associativity. Let

$$(n, \alpha), \quad (m, \beta), \quad (\ell, \gamma) \in N \times H,$$

where $n, m, \ell \in N$ and $\alpha, \beta, \gamma \in H$. First,

$$((n, \alpha)(m, \beta))(\ell, \gamma)$$

has first coordinate

$$n\varphi(\alpha)(m)f(\alpha, \beta)\varphi(\alpha\beta)(\ell)f(\alpha\beta, \gamma),$$

and second coordinate $\alpha\beta\gamma$. On the other hand,

$$(n, \alpha)((m, \beta)(\ell, \gamma))$$

has first coordinate

$$n\varphi(\alpha)(m\varphi(\beta)(\ell)f(\beta, \gamma))f(\alpha, \beta\gamma).$$

Since $\varphi(\alpha)$ is an automorphism of N , this is

$$n\varphi(\alpha)(m)\varphi(\alpha)\varphi(\beta)(\ell)\varphi(\alpha)(f(\beta, \gamma))f(\alpha, \beta\gamma).$$

By the second identity,

$$\varphi(\alpha)\varphi(\beta)(\ell) = f(\alpha, \beta)\varphi(\alpha\beta)(\ell)f(\alpha, \beta)^{-1}.$$

Therefore the first coordinate of $(n, \alpha)((m, \beta)(\ell, \gamma))$ is

$$n\varphi(\alpha)(m)f(\alpha, \beta)\varphi(\alpha\beta)(\ell)f(\alpha, \beta)^{-1}\varphi(\alpha)(f(\beta, \gamma))f(\alpha, \beta\gamma).$$

By the third identity,

$$f(\alpha, \beta)^{-1}\varphi(\alpha)(f(\beta, \gamma))f(\alpha, \beta\gamma) = f(\alpha\beta, \gamma),$$

so this becomes

$$n\varphi(\alpha)(m)f(\alpha, \beta)\varphi(\alpha\beta)(\ell)f(\alpha\beta, \gamma).$$

This agrees with the first coordinate of $((n, \alpha)(m, \beta))(\ell, \gamma)$, and the second coordinates are both $\alpha\beta\gamma$. Hence multiplication is associative.

Now we find inverses. For $(n, \alpha) \in N \times H$, put

$$(n, \alpha)^{-1} = (f(\alpha^{-1}, \alpha)^{-1}\varphi(\alpha^{-1})(n^{-1}), \alpha^{-1}).$$

Let

$$p = f(\alpha^{-1}, \alpha)^{-1}\varphi(\alpha^{-1})(n^{-1}).$$

Then

$$(p, \alpha^{-1})(n, \alpha) = (p\varphi(\alpha^{-1})(n)f(\alpha^{-1}, \alpha), 1) = (1, 1).$$

It remains only to check the product in the other order. Taking $(\alpha, \beta, \gamma) = (\alpha, \alpha^{-1}, \alpha)$ in the third identity gives

$$f(\alpha, \alpha^{-1}) = \varphi(\alpha)(f(\alpha^{-1}, \alpha)).$$

Also, taking $(\alpha, \beta) = (\alpha, \alpha^{-1})$ in the second identity gives

$$\varphi(\alpha)\varphi(\alpha^{-1}) = \Psi(f(\alpha, \alpha^{-1})).$$

Hence

$$\begin{aligned} (n, \alpha)(p, \alpha^{-1}) &= (n\varphi(\alpha)(p)f(\alpha, \alpha^{-1}), 1) \\ &= (n\varphi(\alpha)(f(\alpha^{-1}, \alpha)^{-1}\varphi(\alpha^{-1})(n^{-1})f(\alpha, \alpha^{-1})), 1) \\ &= (nf(\alpha, \alpha^{-1})^{-1}f(\alpha, \alpha^{-1})n^{-1}f(\alpha, \alpha^{-1})^{-1}f(\alpha, \alpha^{-1}), 1) \\ &= (1, 1). \end{aligned}$$

Thus every element has an inverse, and so $E = N \times H$ is a group under the multiplication above. Finally, define

$$i: N \rightarrow E, \quad i(n) = (n, 1),$$

and

$$\pi: E \rightarrow H, \quad \pi(n, \alpha) = \alpha.$$

The map i is injective, and for $n, m \in N$,

$$i(n)i(m) = (n, 1)(m, 1) = (nm, 1) = i(nm),$$

so i is a monomorphism. Also π is a homomorphism, since

$$\pi((n, \alpha)(m, \beta)) = \pi(n\varphi(\alpha)(m)f(\alpha, \beta), \alpha\beta) = \alpha\beta = \pi(n, \alpha)\pi(m, \beta).$$

It is surjective because $\pi(1, \alpha) = \alpha$ for every $\alpha \in H$. Finally,

$$\ker \pi = \{(n, \alpha) \in E \mid \alpha = 1\} = \{(n, 1) \mid n \in N\} = i(N).$$

Therefore (E, i, π) is an extension of N by H . □

We shall return to these ideas in the further exercises to Section 9.

3 Group Actions

Defining group actions

- A (left) action of G on a set X is a map $G \times X \rightarrow X$, the image of (g, x) denoted gx , satisfying $1x = x$ for every $x \in X$ and $(g_1g_2)x = g_1(g_2x)$ for all $g_1, g_2 \in G, x \in X$.
- If G acts on X , we say X is a G -set; for any $H \leq G$, X is also an H -set by restriction.
- For $H \leq G$, the coset space G/H is a G -set under left multiplication: $(g, xH) \mapsto gxH$.

Proposition 3.1. *There is a natural bijective correspondence between the set of actions of G on a set X and the set of homomorphisms from G to S_X .*

Proof. Suppose first that G acts on X . For each $g \in G$, define a function

$$\sigma_g: X \rightarrow X, \quad \sigma_g(x) = gx.$$

We claim that σ_g is a permutation of X . Indeed, $\sigma_{g^{-1}}$ is its inverse, since for every $x \in X$,

$$\sigma_{g^{-1}}(\sigma_g(x)) = g^{-1}(gx) = (g^{-1}g)x = 1x = x$$

and similarly $\sigma_g(\sigma_{g^{-1}}(x)) = x$. Hence $\sigma_g \in S_X$. Now define

$$\Phi: G \rightarrow S_X, \quad \Phi(g) = \sigma_g.$$

For $g_1, g_2 \in G$ and $x \in X$, the action law gives

$$\Phi(g_1g_2)(x) = \sigma_{g_1g_2}(x) = (g_1g_2)x = g_1(g_2x) = \sigma_{g_1}(\sigma_{g_2}(x)).$$

Thus $\Phi(g_1g_2) = \Phi(g_1)\Phi(g_2)$, so Φ is a homomorphism. Conversely, suppose that $\Phi: G \rightarrow S_X$ is a homomorphism. Define a rule $G \times X \rightarrow X$ by

$$gx = \Phi(g)(x).$$

This rule lands in X because each $\Phi(g)$ is a permutation of X . Also,

$$1x = \Phi(1)(x) = x,$$

since $\Phi(1)$ is the identity permutation of X . For $g_1, g_2 \in G$ and $x \in X$,

$$(g_1g_2)x = \Phi(g_1g_2)(x) = (\Phi(g_1)\Phi(g_2))(x) = \Phi(g_1)(\Phi(g_2)(x)) = g_1(g_2x).$$

Therefore this rule is an action of G on X . Finally, these two constructions undo one another. Starting from an action, the homomorphism sends g to the permutation $x \mapsto gx$, so reconstructing the action gives the same rule. Starting from a homomorphism Φ , the action is defined by $gx = \Phi(g)(x)$, so the permutation attached to g is exactly $\Phi(g)$. Hence the correspondence is bijective. $\square$

Faithful actions and Cayley's Theorem

- If $H < G$ with $|G : H| = n$, then the action of G on G/H gives a non-trivial homomorphism $G \rightarrow S_n$ via Proposition 3.1; if G is simple, this homomorphism must be injective.
- The action of G on X is *faithful* (or G acts *faithfully*) if the corresponding homomorphism $G \rightarrow S_X$ is injective; equivalently, the only $g \in G$ satisfying $gx = x$ for every $x \in X$ is the identity.
- If G acts faithfully on X , we call G a *permutation group* on X , since G is then imbedded isomorphically in S_X .

Theorem (Cayley's Theorem). *G is isomorphic with a subgroup of S_G ; in particular, if G is finite with $|G| = n$, then G is isomorphic with a subgroup of S_n .*

Proof. Let G act on itself by left multiplication. In other words, for $g, x \in G$, define

$$gx$$

to be the product of g and x in the group G . This is an action: the identity satisfies $1x = x$, and associativity in G gives

$$(g_1g_2)x = g_1(g_2x)$$

for all $g_1, g_2, x \in G$. By Proposition 3.1, this action gives a homomorphism

$$\Phi: G \rightarrow S_G.$$

Explicitly, $\Phi(g)$ is the permutation of G given by left multiplication by g :

$$\Phi(g)(x) = gx.$$

We now show that Φ is injective. Suppose $g \in \ker \Phi$. Then $\Phi(g)$ is the identity permutation of G , so

$$gx = x$$

for every $x \in G$. Taking $x = 1$, we get $g1 = 1$, and hence $g = 1$. Thus $\ker \Phi = 1$, so Φ is injective. Therefore $G \cong \text{im } \Phi$, and $\text{im } \Phi \leq S_G$. Hence G is isomorphic with a subgroup of S_G . Finally, if G is finite with $|G| = n$, then S_G is isomorphic with S_n , since both are symmetric groups on sets with n elements. Thus G is isomorphic with a subgroup of S_n . $\square$

G -set homomorphisms and orbits

- If X, Y are G -sets, $\varphi: X \rightarrow Y$ is a G -set homomorphism if it commutes with the actions: $\varphi(gx) = g\varphi(x)$ for all $g \in G, x \in X$. A bijective such φ is a G -set isomorphism; we then write $X \cong Y$.
- For each $x \in X$, the orbit of x is $Gx = \{gx \mid g \in G\} \subseteq X$, and the stabilizer of x is $G_x = \{g \in G \mid gx = x\} \leq G$.
- Gx is itself a G -set under the induced action; a subset of X is a G -set under the induced action iff it is a union of orbits.

Lemma 3.2. *If X is a G -set, then $G_{gx} = gG_xg^{-1}$ for any $g \in G$ and $x \in X$.*

Proof. Let $u \in G$. We show that $u \in G_{gx}$ iff $u \in gG_xg^{-1}$. By definition, $u \in G_{gx}$ means that u fixes the element gx , so

$$u \in G_{gx} \iff u(gx) = gx.$$

Using the action law, this is equivalent to

$$(ug)x = gx.$$

Now act on both sides by g^{-1} . Again using the action law, we get

$$(g^{-1}ug)x = x.$$

This means exactly that $g^{-1}ug \in G_x$. Equivalently, $u \in gG_xg^{-1}$. Therefore $G_{gx} = gG_xg^{-1}$. $\square$

Transitive actions and the orbit decomposition

- The action of G on X is *transitive* (or G acts *transitively* on X) if there is some $x \in X$ such that $Gx = X$, equivalently, if for any $x_1, x_2 \in X$ there exists some $g \in G$ such that $gx_1 = x_2$.
- A subset of X is a transitive G -set under the induced action iff it consists of a single orbit.
- For $H \leq G$, the action of G on G/H is transitive: $(yx^{-1})xH = yH$.

Proposition 3.3. *Any G -set has a unique partition consisting of transitive G -sets, namely its partition into orbits.*

Proof. We first show that the orbits form a partition of X . Every $x \in X$ lies in its own orbit Gx , since $x = 1x$. Now let $x, y \in X$, and suppose that $Gx \cap Gy \neq \emptyset$. Then there are $g_1, g_2 \in G$ such that

$$g_1x = g_2y.$$

Hence

$$y = (g_2^{-1}g_1)x,$$

so $y \in Gx$. It follows that $Gy \subseteq Gx$, since for any $u \in G$,

$$uy = u(g_2^{-1}g_1)x = (ug_2^{-1}g_1)x \in Gx.$$

By symmetry, we also have $Gx \subseteq Gy$. Therefore $Gx = Gy$. Thus any two orbits are either equal or disjoint, so the orbits partition X . Each orbit is a transitive G -set. Indeed, if $y, z \in Gx$, then $y = g_1x$ and $z = g_2x$ for some $g_1, g_2 \in G$, and

$$(g_2g_1^{-1})y = (g_2g_1^{-1})(g_1x) = g_2x = z.$$

Hence any element of the orbit can be moved to any other element of the same orbit. Finally, we prove uniqueness. Suppose X has some partition into transitive G -sets, and let P be one piece of this partition. Choose some $x \in P$. Since P is itself a G -set, it is closed under the action of G . Thus

$$Gx \subseteq P.$$

On the other hand, since P is transitive, every $y \in P$ can be reached from x . That is, for every $y \in P$, there is some $g \in G$ such that $gx = y$. Hence every $y \in P$ lies in Gx , so

$$P \subseteq Gx.$$

Therefore $P = Gx$. Since every transitive piece P is an orbit, the only possible partition into transitive G -sets is the partition into orbits. $\square$

Classification of transitive G -sets

- Since any G -set is a union of transitive G -sets, classifying G -sets reduces to classifying transitive G -sets.
- Coset spaces are examples of transitive G -sets; we now show every transitive G -set is isomorphic to one.

Proposition 3.4. *If X is a transitive G -set, then $X \cong G/G_x$ as G -sets for any $x \in X$.*

Proof. Fix $x \in X$. Define

$$\varphi: G/G_x \rightarrow X, \quad \varphi(gG_x) = gx.$$

We first check that φ is well-defined. Suppose $gG_x = hG_x$. Then $h^{-1}g \in G_x$, so $(h^{-1}g)x = x$. Acting by h on both sides gives

$$gx = hx.$$

Hence $\varphi(gG_x) = \varphi(hG_x)$. We next show that φ is injective. Suppose $\varphi(gG_x) = \varphi(hG_x)$. Then $gx = hx$, so acting by h^{-1} on both sides gives

$$(h^{-1}g)x = x.$$

Thus $h^{-1}g \in G_x$, and therefore $gG_x = hG_x$. Hence φ is injective. Since X is transitive, for every $y \in X$ there exists some $g \in G$ such that $gx = y$. Thus $y = \varphi(gG_x)$, so φ is surjective. Finally, we check that φ is a G -set homomorphism. For $u, g \in G$,

$$\varphi(u(gG_x)) = \varphi(ugG_x) = (ug)x = u(gx) = u\varphi(gG_x).$$

Therefore φ is a bijective G -set homomorphism, and hence $X \cong G/G_x$ as G -sets. $\square$

Corollary 3.5 (Orbit-Stabilizer). *Let X be a G -set. Then $Gx \cong G/G_x$ as G -sets for any $x \in X$; in particular, if G is finite, then $|Gx| = |G : G_x|$.*

Proof. The orbit Gx is a transitive G -set under the induced action. The stabilizer of x for this restricted action is still G_x . Hence, by Proposition 3.4,

$$Gx \cong G/G_x$$

as G -sets. If G is finite, then the number of left cosets of G_x in G is $|G : G_x|$. Since Gx is in bijection with G/G_x , we have

$$|Gx| = |G/G_x| = |G : G_x|.$$

□

When are transitive G -sets isomorphic?

- Having decomposed G -sets into transitive ones and identified each with a coset space, the remaining question is when two transitive G -sets are isomorphic.

Lemma 3.6. *Let $\varphi : X \rightarrow Y$ be a homomorphism of G -sets, and let $x \in X$. Then $G_x \leq G_{\varphi(x)}$, and if φ is an isomorphism, then $G_x = G_{\varphi(x)}$.*

Proof. Let $g \in G_x$. Then $gx = x$. Since φ is a G -set homomorphism, we have

$$g\varphi(x) = \varphi(gx) = \varphi(x).$$

Thus $g \in G_{\varphi(x)}$, and therefore $G_x \leq G_{\varphi(x)}$. Now suppose that φ is an isomorphism. Then $\varphi^{-1} : Y \rightarrow X$ is also a G -set homomorphism. Applying the first part to φ^{-1} and the element $\varphi(x) \in Y$, we obtain

$$G_{\varphi(x)} \leq G_{\varphi^{-1}(\varphi(x))} = G_x.$$

Together with the inclusion $G_x \leq G_{\varphi(x)}$, this gives $G_x = G_{\varphi(x)}$. □

Proposition 3.7. *If H and K are subgroups of G , then the G -sets G/H and G/K are isomorphic iff H and K are conjugate in G .*

Proof. The stabilizer of the coset $H \in G/H$ is exactly H . Since every coset of G/H has the form gH for some $g \in G$, Lemma 3.2 shows that the set of stabilizers arising from the G -set G/H is precisely the set of conjugates of H . Similarly, the set of stabilizers arising from G/K is precisely the set of conjugates of K . Suppose first that $\varphi : G/H \rightarrow G/K$ is a G -set isomorphism. By Lemma 3.6,

$$H = G_H = G_{\varphi(H)}.$$

Now $\varphi(H)$ is some coset $gK \in G/K$, and by Lemma 3.2 its stabilizer is gKg^{-1} . Hence

$$H = G_{\varphi(H)} = G_{gK} = gKg^{-1}.$$

Thus H and K are conjugate in G . Conversely, suppose that $H = gKg^{-1}$ for some $g \in G$. Then H is the stabilizer of the coset $gK \in G/K$. Since G/K is transitive, Proposition 3.4 gives

$$G/K \cong G/H$$

as G -sets. Therefore G/H and G/K are isomorphic as G -sets. □

Doubly transitive actions and maximality

- Let X be a G -set. We say X is *doubly transitive* (and G acts *doubly transitively* on X) if whenever (x_1, x_2) and (y_1, y_2) are elements of $X \times X$ with $x_1 \neq x_2$ and $y_1 \neq y_2$, there exists some $g \in G$ such that $gx_1 = y_1$ and $gx_2 = y_2$.
- The natural action of S_n on $\{1, \dots, n\}$ for $n \geq 2$ is doubly transitive. Any doubly transitive G -set is transitive.
- A proper subgroup H of G is *maximal* if there is no proper subgroup of G that properly contains H . Any subgroup of prime index is necessarily maximal by Theorem 1.6.

Proposition 3.8. *Let G be a group, let X be a doubly transitive G -set, and let $x \in X$. Then G_x is a maximal subgroup of G .*

Proof. By Proposition 3.4, we have $X \cong G/G_x$ as G -sets. Thus we may work with the doubly transitive G -set G/G_x . Suppose, for a contradiction, that G_x is not maximal. Then there exists a subgroup K such that

$$G_x < K < G.$$

Choose $g \in G$ with $g \notin K$, and choose $k \in K$ with $k \notin G_x$. Since $k \notin G_x$, the cosets G_x and kG_x are distinct. Since $g \notin K$ and $G_x \leq K$, the cosets G_x and gG_x are also distinct. Because G/G_x is doubly transitive, there exists some $u \in G$ such that

$$uG_x = G_x \quad \text{and} \quad u(kG_x) = gG_x.$$

The first equality gives $u \in G_x$, and hence $u \in K$. Since $k \in K$, we have $uk \in K$. The second equality says

$$ukG_x = gG_x,$$

so $g^{-1}uk \in G_x \leq K$. Since both uk and $g^{-1}uk$ lie in K , it follows that

$$g = uk(g^{-1}uk)^{-1} \in K,$$

contradicting our choice of $g \notin K$. Therefore no subgroup lies strictly between G_x and G , and so G_x is maximal. $\square$

Blocks and primitivity

- A non-empty subset B of a transitive G -set X is a *block* if B and $gB = \{gx \mid x \in B\}$ are either equal or disjoint for every $g \in G$.
- X itself is always a block, and any one-element subset of X is always a block; the transitive G -set X is *primitive* if these are the only blocks.

Proposition 3.9. *There is a natural bijective correspondence between the set of blocks of a transitive G -set X which contain a given element x and the set of subgroups of G which contain G_x .*

Proof. Let B be a block of X containing x , and define

$$H_B = \{g \in G \mid gx \in B\}.$$

We first show that H_B is a subgroup of G . Since $x \in B$, we have $1x = x \in B$, so $1 \in H_B$. Now let $g, g' \in H_B$. Then $gx \in B$ and $g'x \in B$. Since both x and gx lie in B , the sets B and gB meet, and therefore $gB = B$ because B is a block. Hence

$$(gg')x = g(g'x) \in gB = B,$$

so $gg' \in H_B$. Similarly, if $g \in H_B$, then $gx \in B$. Since $g^{-1}(gx) = x \in B$, the sets $g^{-1}B$ and B meet, and therefore $g^{-1}B = B$. Thus $g^{-1}x \in B$, so $g^{-1} \in H_B$. Therefore $H_B \leq G$. Also, if $u \in G_x$, then $ux = x \in B$, so $u \in H_B$. Hence $G_x \leq H_B$.

Thus every block B containing x gives a subgroup H_B containing G_x . We now show that this assignment is injective. Suppose B and B' are distinct blocks containing x . Choose $y \in B'$ with $y \notin B$. Since X is transitive, there exists some $g \in G$ such that $gx = y$. Then $g \in H_{B'}$, but $g \notin H_B$. Hence $H_B \neq H_{B'}$. Conversely, let H be a subgroup of G containing G_x , and define

$$B_H = Hx = \{hx \mid h \in H\}.$$

Clearly $x \in B_H$, so B_H is non-empty. We claim that B_H is a block. If $g \in H$, then $gB_H = B_H$. Now suppose that $gB_H \cap B_H \neq \emptyset$. Then for some $h_1, h_2 \in H$,

$$gh_1x = h_2x.$$

Hence

$$(h_2^{-1}gh_1)x = x,$$

so $h_2^{-1}gh_1 \in G_x \leq H$. Since $h_1, h_2 \in H$, we get

$$g = h_2(h_2^{-1}gh_1)h_1^{-1} \in H.$$

Therefore, whenever gB_H meets B_H , we must have $g \in H$, and hence $gB_H = B_H$. If gB_H does not meet B_H , then it is disjoint from B_H . Thus B_H is a block.

Finally, we check that this construction gives back the subgroup H . By definition,

$$H_{B_H} = \{g \in G \mid gx \in B_H\}.$$

If $g \in H$, then $gx \in Hx = B_H$, so $g \in H_{B_H}$. Thus $H \leq H_{B_H}$. Conversely, if $g \in H_{B_H}$, then $gx = hx$ for some $h \in H$. Therefore

$$(h^{-1}g)x = x,$$

so $h^{-1}g \in G_x \leq H$. Since $h \in H$, we get $g \in H$. Hence $H_{B_H} = H$, proving that every subgroup containing G_x arises from a unique block containing x . $\square$

Corollary 3.10. *Let X be a transitive G -set. Then X is primitive iff G_x is a maximal subgroup of G for every $x \in X$.*

Proof. Suppose first that X is primitive, and fix $x \in X$. By Proposition 3.9, the blocks of X which contain x correspond exactly to the subgroups of G which contain G_x . Since X is primitive, the only blocks containing x are $\{x\}$ and X . These correspond to the subgroups G_x and G , respectively. Hence there is no subgroup H with

$$G_x < H < G.$$

Therefore G_x is maximal in G .

Conversely, suppose that G_x is maximal in G for every $x \in X$. Let B be a block of X , and choose some $x \in B$. Again by Proposition 3.9, the block B corresponds to a subgroup H_B satisfying

$$G_x \leq H_B \leq G.$$

Since G_x is maximal, either $H_B = G_x$ or $H_B = G$. If $H_B = G_x$, then the corresponding block is $\{x\}$; if $H_B = G$, then the corresponding block is X . Thus every block of X is either a one-element subset or all of X . Hence X is primitive. $\square$

Consequences for primitivity

- If X is a transitive G -set, all stabilizers are conjugate by Lemma 3.2; so if G_x is maximal for some x , it is maximal for every x , and Corollary 3.10 can be modified accordingly.

Corollary 3.11. *Any doubly transitive G -set is primitive.*

Proof. This follows from Proposition 3.8 and Corollary 3.10. Indeed, if X is doubly transitive, then Proposition 3.8 shows that G_x is maximal in G for every $x \in X$. Hence, by Corollary 3.10, the G -set X is primitive. $\square$

Counting via group actions

Proposition 3.12. *If G is finite and $H, K \leq G$, then*

$$|HK| = \frac{|H||K|}{|H \cap K|}.$$

Proof. Let $X = G/K$, and consider X as an H -set by restricting the usual left multiplication action of G on G/K . The orbit of the coset $K \in G/K$ under this action is $\{hK \mid h \in H\}$. The union of the cosets in this orbit is exactly

$$\bigcup_{h \in H} hK = HK.$$

Since distinct left cosets of K are disjoint and each has $|K|$ elements, we have

$$|HK| = |\{hK \mid h \in H\}| |K|.$$

The stabilizer of K under the action of H is

$$H_K = \{h \in H \mid hK = K\}.$$

But $hK = K$ iff $h \in K$, and we already have $h \in H$. Hence

$$H_K = H \cap K.$$

By Corollary 3.5, this orbit has size

$$|\{hK \mid h \in H\}| = |H : H \cap K| = \frac{|H|}{|H \cap K|}.$$

Therefore

$$|HK| = |\{hK \mid h \in H\}| |K| = \frac{|H|}{|H \cap K|} |K| = \frac{|H||K|}{|H \cap K|}.$$

□

Centralizers and conjugacy classes

- For $x \in G$, the *centralizer* of x in G is $C_G(x) = \{g \in G \mid gx = xg\} \leq G$.
- For $S \subseteq G$, $C_G(S) = \{g \in G \mid gx = xg \text{ for all } x \in S\} = \bigcap_{x \in S} C_G(x)$, called the centralizer of S in G .
- $Z(G) = C_G(G)$, and $x \in Z(G)$ iff $C_G(x) = G$.
- The *conjugacy class* of $x \in G$ is $\{gxg^{-1} \mid g \in G\}$. By Proposition 1.10, the conjugacy class of $\rho \in S_n$ consists of all elements of S_n having the same cycle structure as ρ .
- More explicitly, if $\rho \in S_n$ has m_i cycles of length i for each $i \geq 1$, then the size of its conjugacy class in S_n is

$$\frac{n!}{\prod_{i \geq 1} i^{m_i} m_i!}.$$

To see the denominator, first arrange the n symbols in an ordered list and then cut this list into m_i blocks of length i for each i . This gives $n!$ arrangements. However, each i -cycle can be written in i cyclic rotations, and these rotations define the same cycle; this gives a factor of i^{m_i} . Also, the m_i cycles of the same length may be permuted among themselves without changing the resulting disjoint cycle decomposition; this gives a factor of $m_i!$. Dividing by all these repetitions gives the formula [24, p. 47].

Proposition 3.13. *The conjugacy classes of G form a partition of G , and if G is finite then an element $x \in G$ has $|G : C_G(x)|$ conjugates in G .*

Proof. Let G act on itself by conjugation: for $g, x \in G$, define

$$g \cdot x = gxg^{-1}.$$

This is a left action, since $1 \cdot x = x$ and

$$g_1 \cdot (g_2 \cdot x) = g_1(g_2 x g_2^{-1}) g_1^{-1} = (g_1 g_2) x (g_1 g_2)^{-1} = (g_1 g_2) \cdot x.$$

The orbit of $x \in G$ under this action is

$$\{gxg^{-1} \mid g \in G\},$$

which is exactly the conjugacy class of x . Hence the first assertion follows from Proposition 3.3, since the orbits of any group action form a partition. If G is finite, then Corollary 3.5 says that the size of the orbit of x is the index of the stabilizer of x . The stabilizer is

$$G_x = \{g \in G \mid gxg^{-1} = x\}.$$

But $gxg^{-1} = x$ iff $gx = xg$, so

$$G_x = C_G(x).$$

Therefore the conjugacy class of x has size $|G : C_G(x)|$. □

Normalizers and conjugates of subgroups

- A subgroup of G is normal iff it is a union of conjugacy classes; such a union is in fact disjoint.
- The order of a normal subgroup of a finite group G must be a sum of orders of conjugacy classes of G .
- For $H \leq G$, the *normalizer* of H in G is $N_G(H) = \{g \in G \mid gHg^{-1} = H\}$. We have $N_G(H) \leq G$ and $H \trianglelefteq N_G(H)$; $N_G(H)$ is the largest subgroup of G in which H is normal, so $N_G(H) = G$ iff $H \trianglelefteq G$.

Proposition 3.14. *A subgroup H of a finite group G has exactly $|G : N_G(H)|$ conjugates in G . In particular, the number of conjugates of H in G divides $|G : H|$ and is equal to 1 iff $H \trianglelefteq G$.*

Proof. Let $\mathcal{P}(G)$ be the set of subsets of G , and let G act on $\mathcal{P}(G)$ by conjugation:

$$g \cdot S = gSg^{-1}$$

for $g \in G$ and $S \subseteq G$. This is a left action, since $1S1^{-1} = S$ and

$$g_1 \cdot (g_2 \cdot S) = g_1(g_2Sg_2^{-1})g_1^{-1} = (g_1g_2)S(g_1g_2)^{-1} = (g_1g_2) \cdot S.$$

The orbit of H under this action is

$$\{gHg^{-1} \mid g \in G\},$$

which is precisely the set of conjugates of H in G . The stabilizer of H is

$$\{g \in G \mid gHg^{-1} = H\} = N_G(H).$$

Therefore, by Corollary 3.5, the number of conjugates of H in G is

$$|G : N_G(H)|.$$

Since $H \leq N_G(H) \leq G$, we have

$$|G : H| = |G : N_G(H)| |N_G(H) : H|.$$

Hence the number of conjugates of H divides $|G : H|$. Finally, H has exactly one conjugate iff $|G : N_G(H)| = 1$, iff $N_G(H) = G$. This is equivalent to $gHg^{-1} = H$ for every $g \in G$, which is exactly the condition that $H \trianglelefteq G$. $\square$

Double cosets

- Let G be a group and let H, K be subgroups of G . A *double coset* of H and K in G is a set $HxK = \{h x k \mid h \in H, k \in K\}$ for some $x \in G$.
- Any two double cosets are either disjoint or equal, as is the case with ordinary cosets.

Proposition 3.15. *If G is finite and $H, K \leq G$, then for any $x \in G$ we have*

$$|HxK| = \frac{|H||K|}{|H \cap xKx^{-1}|} = \frac{|H||K|}{|x^{-1}Hx \cap K|}.$$

Proof. Consider G/K as an H -set by restricting the usual left multiplication action. The orbit of $xK \in G/K$ under this action is

$$\{hxK \mid h \in H\}.$$

The union in G of the cosets in this orbit is exactly HxK . Since distinct left cosets of K are disjoint and each has $|K|$ elements, we have

$$|HxK| = |\{hxK \mid h \in H\}| |K|.$$

The stabilizer of xK under the action of H is

$$\{h \in H \mid hxK = xK\}.$$

Now

$$hxK = xK \iff x^{-1}hx \in K \iff h \in xKx^{-1}.$$

Thus the stabilizer is $H \cap xKx^{-1}$. By Corollary 3.5,

$$|\{hxK \mid h \in H\}| = |H : H \cap xKx^{-1}| = \frac{|H|}{|H \cap xKx^{-1}|}.$$

Therefore

$$|HxK| = \frac{|H|}{|H \cap xKx^{-1}|} |K| = \frac{|H||K|}{|H \cap xKx^{-1}|}.$$

Finally, conjugating the denominator by x^{-1} gives

$$x^{-1}(H \cap xKx^{-1})x = x^{-1}Hx \cap K.$$

Conjugation is a bijection, so these two intersections have the same order. Hence

$$\frac{|H||K|}{|H \cap xKx^{-1}|} = \frac{|H||K|}{|x^{-1}Hx \cap K|}.$$

□

Exercises

Exercise 3.1. Show that a finite simple group whose order is at least $r!$ cannot have a proper subgroup of index r .

Solution. Suppose, for a contradiction, that $H < G$ and $|G : H| = r$. Let G act on the coset space G/H by left multiplication. Since G/H has r elements, Proposition 3.1 gives a homomorphism

$$\Phi : G \rightarrow S_{G/H} \cong S_r.$$

This is the same permutation-representation idea as in Cayley's Theorem, with the coset space G/H in place of G itself. Now $\ker \Phi \trianglelefteq G$. Since G is simple, either $\ker \Phi = G$ or $\ker \Phi = 1$. If $\ker \Phi = G$, then every $g \in G$ fixes every coset. In particular,

$$gH = H$$

for every $g \in G$. Hence every $g \in G$ lies in H , so $G = H$, contradicting that H is proper. Therefore $\ker \Phi \neq G$, and so $\ker \Phi = 1$. Thus Φ is injective, so G is isomorphic to a subgroup of S_r . Hence

$$|G| \leq |S_r| = r!.$$

By hypothesis $|G| \geq r!$, so $|G| = r!$. Therefore $\Phi(G)$ is a subgroup of S_r with the same order as S_r , and hence $\Phi(G) = S_r$. Thus $G \cong S_r$. For $r \geq 3$, this is impossible because $A_r \trianglelefteq S_r$ is a non-trivial proper normal subgroup. Hence no such subgroup H exists. As stated, the exercise also needs the condition $r \geq 3$: for $r = 2$, the group C_2 is simple, has order $2 = 2!$, and has the proper subgroup 1 of index 2 . □

Exercise 3.2. Show that a group G acts doubly transitively on a set X iff G_x acts transitively on $X - \{x\}$ for every $x \in X$. (Here we must assume that X has more than two elements.)

Solution. Suppose first that G acts doubly transitively on X , and fix $x \in X$. We show that G_x acts transitively on $X - \{x\}$. First, G_x preserves $X - \{x\}$. Indeed, if $g \in G_x$ and $y \neq x$, then $gy \neq x$; otherwise $gy = x$, and applying g^{-1} gives $y = g^{-1}x = x$, a contradiction. Now let $y, z \in X - \{x\}$. Then (x, y) and (x, z) are ordered pairs of distinct elements of X . Since G acts doubly transitively on X , there exists $g \in G$ such that

$$gx = x \quad \text{and} \quad gy = z.$$

The first equality says that $g \in G_x$, and the second says that this element of G_x sends y to z . Hence G_x acts transitively on $X - \{x\}$. Conversely, suppose that G_x acts transitively on $X - \{x\}$ for every $x \in X$. Let (x_1, x_2) and (y_1, y_2) be ordered pairs of distinct elements of X . We must find some $g \in G$ such that $gx_1 = y_1$ and $gx_2 = y_2$. We first find an element $a \in G$ such that $ax_1 = y_1$. If $x_1 = y_1$, take $a = 1$. If $x_1 \neq y_1$, choose

$$w \in X - \{x_1, y_1\},$$

which is possible since X has more than two elements. By hypothesis, G_w acts transitively on $X - \{w\}$. Since $x_1, y_1 \in X - \{w\}$, there exists $a \in G_w$ such that

$$ax_1 = y_1.$$

Thus, in either case, we have some $a \in G$ with $ax_1 = y_1$. Since a acts as a bijection on X and $x_2 \neq x_1$, we have $ax_2 \neq ax_1 = y_1$. Hence $ax_2 \in X - \{y_1\}$. Also $y_2 \in X - \{y_1\}$, since $y_2 \neq y_1$. By hypothesis, G_{y_1} acts transitively on $X - \{y_1\}$, so there exists $b \in G_{y_1}$ such that

$$b(ax_2) = y_2.$$

Since $b \in G_{y_1}$, we also have $by_1 = y_1$. Now set $g = ba$. Then

$$gx_1 = b(ax_1) = by_1 = y_1$$

and

$$gx_2 = b(ax_2) = y_2.$$

Therefore G acts doubly transitively on X . □

Exercise 3.3. Show directly from the definitions (that is, without reference to Propositions 3.8 and 3.9) that a doubly transitive G -set is primitive.

Solution. Let B be a block of X . We show that B is either a one-element subset of X or all of X . If B has exactly one element, then B is one of the trivial blocks. Suppose instead that B has at least two elements, and choose distinct elements $x, y \in B$. We will show that $B = X$. Let $z \in X$. If $z = x$, then $z \in B$. So suppose that $z \neq x$. Since $x \neq y$ and $x \neq z$, the ordered pairs

$$(x, y) \quad \text{and} \quad (x, z)$$

are pairs of distinct elements of X . Since G acts doubly transitively on X , there exists $g \in G$ such that

$$gx = x \quad \text{and} \quad gy = z.$$

Now $x \in B$ and $gx = x \in B$, so gB meets B . Since B is a block, this forces $gB = B$. But $y \in B$, so $gy \in gB$. Therefore

$$z = gy \in gB = B.$$

Hence every $z \in X$ lies in B , so $B = X$. Thus every block of X is either a one-element subset or all of X . Therefore X is primitive. □

Exercise 3.4. Let G be the subgroup of S_5 generated by $(1\ 2\ 3\ 4\ 5)$, and let G act on $X = \{1, 2, 3, 4, 5\}$ in the canonical way. Show that this action is primitive, but not doubly transitive.

Solution. Let $\sigma = (1\ 2\ 3\ 4\ 5)$, so $G = \langle \sigma \rangle$ has order 5. First we show that the action is primitive. The action is transitive, since powers of σ move any element of X to any other element of X . Fix $x \in X$. No non-identity power of σ fixes x , since σ^k moves x by k steps around the 5-cycle for $1 \leq k \leq 4$. Hence

$$G_x = 1.$$

Since G has prime order, its only subgroups are 1 and G . Therefore $G_x = 1$ is maximal in G . By Corollary 3.10, the action of G on X is primitive. We now show that the action is not doubly transitive. If it were doubly transitive, then there would exist some $g \in G$ sending the ordered pair $(1, 2)$ to the ordered pair $(1, 3)$; that is,

$$g1 = 1 \quad \text{and} \quad g2 = 3.$$

But the only element of G which fixes 1 is the identity, and the identity sends 2 to 2, not to 3. Thus no such g exists. Therefore the action is not doubly transitive. □

Exercise 3.5. Let $N \trianglelefteq G$, and let $y \in N$. Show that the conjugacy class of y in G is a union of conjugacy classes of the group N . Show further that there is a bijective correspondence between the conjugacy classes of N which comprise the conjugacy class of y in G and the cosets of $NC_G(y)$ in G .

Solution. Let

$$\mathcal{C} = \{gyg^{-1} \mid g \in G\}$$

be the conjugacy class of y in G . Since $N \trianglelefteq G$ and $y \in N$, every element gyg^{-1} lies in N . Hence $\mathcal{C} \subseteq N$. Now let $z \in \mathcal{C}$. By definition of $\mathcal{C}$, there is some $g \in G$ such that

$$z = gyg^{-1}.$$

If $n \in N$, then

$$nzn^{-1} = n(gyg^{-1})n^{-1} = (ng)y(ng)^{-1}.$$

Since $ng \in G$, this is again an element of $\mathcal{C}$. Therefore, whenever $\mathcal{C}$ contains one element z , it contains every N -conjugate of z . Thus $\mathcal{C}$ is a union of conjugacy classes of N . For the second assertion, set $C = C_G(y)$. Since $N \trianglelefteq G$, the product NC is a subgroup of G . We define a map from the left cosets of NC in G to the N -conjugacy classes contained in $\mathcal{C}$ by

$$\Psi: G/NC \rightarrow \{N\text{-conjugacy classes contained in } \mathcal{C}\}, \quad \Psi(gNC) = \{n(gyg^{-1})n^{-1} \mid n \in N\}.$$

In words, the coset gNC is sent to the N -conjugacy class of gyg^{-1} . We first check that this is well-defined. Suppose $hNC = gNC$. Then $h = gmc$ for some $m \in N$ and $c \in C$. Since $c \in C_G(y)$, we have $cyc^{-1} = y$. Hence

$$hyh^{-1} = gmcyc^{-1}m^{-1}g^{-1} = gmy m^{-1}g^{-1}.$$

But

$$gmy m^{-1}g^{-1} = (gmg^{-1})(gyg^{-1})(gmg^{-1})^{-1}.$$

Since $N \trianglelefteq G$, we have $gmg^{-1} \in N$. Therefore hyh^{-1} is N -conjugate to gyg^{-1} . Thus hNC and gNC give the same N -conjugacy class, so Ψ is well-defined. The map Ψ is onto. Indeed, take any N -conjugacy class contained in $\mathcal{C}$. It contains some element gyg^{-1} with $g \in G$, so it is exactly the image of the coset gNC . Finally, we show that Ψ is one-to-one. Suppose

$$\Psi(gNC) = \Psi(hNC).$$

Then gyg^{-1} and hyh^{-1} are N -conjugate, so there exists $n \in N$ such that

$$hyh^{-1} = n(gyg^{-1})n^{-1} = (ng)y(ng)^{-1}.$$

This means that h and ng conjugate y to the same element. Equivalently,

$$((ng)^{-1}h)y((ng)^{-1}h)^{-1} = y.$$

Hence $(ng)^{-1}h \in C_G(y) = C$. So $h = ngc$ for some $c \in C$. Since N is normal in G , we have $g^{-1}ng \in N$, and therefore

$$h = ngc = g(g^{-1}ng)c \in gNC.$$

Thus $hNC = gNC$. Therefore Ψ is injective. Hence the N -conjugacy classes which make up the G -conjugacy class of y are in bijective correspondence with the cosets of $NC_G(y)$ in G . $\square$

Exercise 3.6. Let G be a finite group, and let r be the number of conjugacy classes of G . Show that $|\{(a, b) \in G \times G \mid ab = ba\}| = r|G|$.

Solution. We count the same set in a slower way. Fix $a \in G$. The elements $b \in G$ such that $ab = ba$ are exactly the elements of the centralizer $C_G(a)$. Hence the number of commuting pairs is

$$\sum_{a \in G} |C_G(a)|.$$

We now group this sum by conjugacy classes. Let K be one conjugacy class of G , and choose a representative $x \in K$. If $a \in K$, then $a = gxg^{-1}$ for some $g \in G$. Conjugation by g gives a bijection

$$C_G(x) \rightarrow C_G(a), \quad u \mapsto gug^{-1},$$

so $|C_G(a)| = |C_G(x)|$. Thus, as a runs through the elements of K , the number $|C_G(a)|$ is always the same number $|C_G(x)|$. Therefore the part of the sum coming from this one conjugacy class is

$$\sum_{a \in K} |C_G(a)| = \underbrace{|C_G(x)| + \cdots + |C_G(x)|}_{|K| \text{ times}} = |K| |C_G(x)|.$$

By Proposition 3.13,

$$|K| = |G : C_G(x)|.$$

Since G is finite, this means

$$|K| |C_G(x)| = |G : C_G(x)| |C_G(x)| = |G|.$$

Hence each conjugacy class contributes exactly $|G|$ to the total sum. Since there are r conjugacy classes, we get

$$|\{(a, b) \in G \times G \mid ab = ba\}| = r|G|.$$

□

Exercise 3.7. Show that $C_G(gxg^{-1}) = gC_G(x)g^{-1}$ for any elements g and x of a group G .

Solution. Let G act on itself by conjugation, so that $u \cdot x = uxu^{-1}$ for $u, x \in G$. Under this action, the stabilizer of x is

$$G_x = \{u \in G \mid uxu^{-1} = x\}.$$

But $uxu^{-1} = x$ iff $ux = xu$, so

$$G_x = C_G(x).$$

Also, the image of x under g is

$$g \cdot x = gxg^{-1}.$$

Therefore Lemma 3.2, applied to this conjugation action, gives

$$C_G(gxg^{-1}) = G_{g \cdot x} = gG_xg^{-1} = gC_G(x)g^{-1}.$$

□

Exercise 3.8. Let $n \geq 5$, and assume that A_n is simple. (A proof of this fact is outlined in Exercise 7.8.) Use Exercise 3.1 to show that S_n has no proper subgroup of index less than n other than A_n .

Solution. Let $H < S_n$ be a proper subgroup with

$$|S_n : H| < n.$$

We show that $H = A_n$. Suppose, for a contradiction, that $H \neq A_n$, and set

$$K = H \cap A_n.$$

We first check that K is a proper subgroup of A_n . If $K = A_n$, then $A_n \leq H$. Since $H \neq A_n$, the subgroup H must contain some odd permutation. But then H contains all even permutations and at least one odd permutation. Multiplying that odd permutation by the elements of A_n gives all odd permutations, so $H = S_n$, contradicting that H is proper. Hence

$$K < A_n.$$

We now show that K has index less than n in A_n . There are two cases. First suppose that $H \leq A_n$. Then $K = H$, and

$$|S_n : H| = |S_n : A_n| |A_n : H| = 2|A_n : K|.$$

Therefore $|A_n : K| < n$. Now suppose that $H \not\leq A_n$. Then H contains an odd permutation, so

$$HA_n = S_n.$$

By Proposition 3.12,

$$|S_n| = |HA_n| = \frac{|H||A_n|}{|H \cap A_n|} = \frac{|H||A_n|}{|K|}.$$

Dividing by $|H|$ gives

$$|S_n : H| = |A_n : K|.$$

Hence again $|A_n : K| < n$. Thus, in either case, K is a proper subgroup of A_n of index

$$s = |A_n : K| < n.$$

Since $s < n$, we have $s \leq n - 1$, and hence

$$s! \leq (n-1)! \leq \frac{n!}{2} = |A_n|$$

because $n \geq 5$. If $s = 2$, then K is a subgroup of index 2 in A_n , hence normal in A_n , contradicting the simplicity of A_n . If $s \geq 3$, then Exercise 3.1 applies to the simple group A_n and rules out a proper subgroup of index s . This contradiction shows that our assumption $H \neq A_n$ was false. Therefore the only proper subgroup of S_n of index less than n is A_n . $\square$

Further Exercises

Let X be a transitive G -set. A *system of imprimitivity* of X is a partition of X which is permuted by the action of G . Note that a system of imprimitivity of a G -set is itself a G -set.

Exercise 3.9. Show that there is a bijective correspondence between the set of blocks which contain a given element of X and the set of systems of imprimitivity of X . (Observe that when X is finite, this implies that any two elements of X lie in the same number of blocks.)

Solution. Fix $x \in X$. We construct maps in both directions. First let B be a block of X with $x \in B$. Define

$$\mathcal{P}_B = \{gB \mid g \in G\}.$$

We show that $\mathcal{P}_B$ is a system of imprimitivity of X . The sets in $\mathcal{P}_B$ are either equal or disjoint. Indeed, suppose

$$gB \cap hB \neq \emptyset.$$

Applying g^{-1} gives

$$B \cap g^{-1}hB \neq \emptyset.$$

Since B is a block, this forces $B = g^{-1}hB$. Multiplying by g , we get $gB = hB$. The sets in $\mathcal{P}_B$ also cover X . Let $y \in X$. Since X is transitive, there exists $g \in G$ such that $gx = y$. Since $x \in B$, we have $y = gx \in gB$. Hence every element of X lies in some member of $\mathcal{P}_B$. Thus $\mathcal{P}_B$ is a partition of X . Moreover, G permutes the pieces, because if $u \in G$, then

$$u(gB) = (ug)B,$$

which is again an element of $\mathcal{P}_B$. Hence $\mathcal{P}_B$ is a system of imprimitivity. Conversely, let $\mathcal{P}$ be a system of imprimitivity of X . Since $\mathcal{P}$ is a partition, there is a unique piece $B \in \mathcal{P}$ such that $x \in B$. We claim that B is a block. Let $g \in G$. Since $\mathcal{P}$ is permuted by G , the set gB is also a piece of $\mathcal{P}$. Since $\mathcal{P}$ is a partition, the two pieces B and gB are either equal or disjoint. Therefore B is a block containing x . These two constructions undo one another. Starting with a block B containing x , the system $\mathcal{P}_B$ contains the piece $B = 1B$. Since $\mathcal{P}_B$ is a partition and $x \in B$, this is the unique piece of $\mathcal{P}_B$ containing x . Thus passing from B to $\mathcal{P}_B$ and then taking the piece containing x returns the original block B . Conversely, start with a system of imprimitivity $\mathcal{P}$, and let B be the unique piece of $\mathcal{P}$ containing x . We claim that the system obtained from this block is the original system:

$$\mathcal{P} = \{gB \mid g \in G\}.$$

Since $\mathcal{P}$ is permuted by G , every set gB is a piece of $\mathcal{P}$. Conversely, let $P \in \mathcal{P}$. Choose $y \in P$. Since X is transitive, there exists $g \in G$ such that $gx = y$. Now $x \in B$, so $y = gx \in gB$. Also gB is a piece of $\mathcal{P}$. Thus P and gB are two pieces of the partition $\mathcal{P}$ containing y , so they must be equal. Hence $P = gB$, and every piece of $\mathcal{P}$ is a translate of B . Therefore the blocks containing x are in bijective correspondence with the systems of imprimitivity of X . If X is finite, then for each chosen element $x \in X$, the blocks containing x are in bijection with the same set of systems of imprimitivity. Hence the number of blocks containing x is independent of x , so any two elements of X lie in the same number of blocks. $\square$

Exercise 3.10. Suppose that the G -set Y is an epimorphic image of a G -set X . Show that there is a G -set isomorphism between Y and some system of imprimitivity of X .

Solution. Since Y is an epimorphic image of X , there is a surjective G -set homomorphism

$$\varphi: X \rightarrow Y.$$

For each $y \in Y$, let

$$X_y = \varphi^{-1}(y) = \{x \in X \mid \varphi(x) = y\}.$$

Since φ is surjective, each X_y is non-empty. The fibers of φ form a partition of X , so set

$$\mathcal{P} = \{X_y \mid y \in Y\}.$$

We show that $\mathcal{P}$ is a system of imprimitivity. Let $g \in G$ and $y \in Y$. We claim that

$$gX_y = X_{gy}.$$

Indeed, if $x \in X_y$, then $\varphi(x) = y$, and since φ is a G -set homomorphism,

$$\varphi(gx) = g\varphi(x) = gy.$$

Thus $gx \in X_{gy}$, so $gX_y \subseteq X_{gy}$. Conversely, if $z \in X_{gy}$, then $\varphi(z) = gy$. Since $g^{-1}z \in X$, we have

$$\varphi(g^{-1}z) = g^{-1}\varphi(z) = g^{-1}(gy) = y.$$

Hence $g^{-1}z \in X_y$, and therefore $z \in gX_y$. Thus $X_{gy} \subseteq gX_y$, proving $gX_y = X_{gy}$. Therefore the action of G permutes the pieces of $\mathcal{P}$, so $\mathcal{P}$ is a system of imprimitivity of X . Now define

$$\theta: Y \rightarrow \mathcal{P}, \quad \theta(y) = X_y.$$

This map is surjective by definition of $\mathcal{P}$. It is injective because distinct elements of Y have disjoint fibers: if $X_y = X_{y'}$, then this common fiber is non-empty, so choose x in it. Then $\varphi(x) = y$ and $\varphi(x) = y'$, hence $y = y'$. Finally, θ is a G -set homomorphism. For $g \in G$ and $y \in Y$,

$$\theta(gy) = X_{gy} = gX_y = g\theta(y).$$

Therefore θ is a G -set isomorphism between Y and the system of imprimitivity $\mathcal{P}$. $\square$

If X is a G -set, we use $[X]$ to denote the isomorphism class of X .

Exercise 3.11. Let G be a finite group, and let $S(G)$ be the set of isomorphism classes of finite G -sets. Show that we can define sum and product operations on $S(G)$ by $[X] + [Y] = [X \cup Y]$ and $[X][Y] = [X \times Y]$.

Solution. We must check that these operations do not depend on the chosen representatives of the isomorphism classes. Suppose $X \cong X'$ and $Y \cong Y'$ as G -sets. Choose G -set isomorphisms

$$\alpha: X \rightarrow X' \quad \text{and} \quad \beta: Y \rightarrow Y'.$$

First consider the sum operation. We regard $X \cup Y$ as the disjoint union of X and Y , with G acting on each piece in the usual way. Define

$$\alpha \cup \beta: X \cup Y \rightarrow X' \cup Y'$$

by

$$(\alpha \cup \beta)(z) = \begin{cases} \alpha(z), & z \in X, \\ \beta(z), & z \in Y. \end{cases}$$

Since α and β are bijections, $\alpha \cup \beta$ is a bijection. It also commutes with the G -action, because if $z \in X$, then

$$(\alpha \cup \beta)(gz) = \alpha(gz) = g\alpha(z) = g(\alpha \cup \beta)(z),$$

and the same calculation using β works when $z \in Y$. Thus $X \cup Y \cong X' \cup Y'$ as G -sets, so $[X] + [Y] = [X \cup Y]$ is well-defined. Now consider the product operation. The product $X \times Y$ is a G -set by the diagonal action

$$g(x, y) = (gx, gy).$$

Define

$$\alpha \times \beta: X \times Y \rightarrow X' \times Y', \quad (\alpha \times \beta)(x, y) = (\alpha(x), \beta(y)).$$

Since α and β are bijections, $\alpha \times \beta$ is a bijection. Also,

$$(\alpha \times \beta)(g(x, y)) = (\alpha \times \beta)(gx, gy) = (\alpha(gx), \beta(gy)) = (g\alpha(x), g\beta(y)) = g(\alpha(x), \beta(y)) = g(\alpha \times \beta)(x, y).$$

Therefore $\alpha \times \beta$ is a G -set isomorphism, so $X \times Y \cong X' \times Y'$ as G -sets. Hence $[X][Y] = [X \times Y]$ is well-defined. $\square$

Exercise 3.12. (cont.) Let $B(G)$ be the set obtained from $S(G)$ by adjoining formal additive inverses of isomorphism classes, in the same way that $\mathbb{Z}$ is obtained from $\mathbb{N}$ by adjoining the additive inverses of positive integers. (The additive identity here will be the isomorphism class of the empty set.) Show that the operations defined above on $S(G)$ extend to give a commutative ring structure on $B(G)$. This ring $B(G)$ is called the *Burnside ring* of G .

Solution. By Exercise 3.11, the operations

$$[X] + [Y] = [X \cup Y], \quad [X][Y] = [X \times Y]$$

are well-defined on isomorphism classes of finite G -sets. The set $B(G)$ is obtained by formally adjoining additive inverses, so every element of $B(G)$ may be represented as a formal difference

$$[X] - [Y],$$

where X and Y are finite G -sets. Addition is then defined by collecting the positive and negative parts:

$$([X] - [Y]) + ([U] - [V]) = [X \cup U] - [Y \cup V].$$

This gives an abelian group under addition. The zero element is $[\emptyset]$, and the additive inverse of $[X] - [Y]$ is $[Y] - [X]$, since

$$([X] - [Y]) + ([Y] - [X]) = [X \cup Y] - [Y \cup X] = [\emptyset]$$

in the formal group obtained from $S(G)$. We extend multiplication by the distributive rule:

$$([X] - [Y])([U] - [V]) = [X \times U] - [X \times V] - [Y \times U] + [Y \times V].$$

This agrees with the original product on $S(G)$ when $Y = V = \emptyset$. The usual bijections

$$X \cup Y \cong Y \cup X, \quad (X \cup Y) \cup Z \cong X \cup (Y \cup Z)$$

are G -set isomorphisms, so addition is commutative and associative. Similarly,

$$X \times Y \cong Y \times X, \quad (X \times Y) \times Z \cong X \times (Y \times Z)$$

as G -sets, so multiplication is commutative and associative. The multiplicative identity is the class $[\{*\}]$ of the one-point G -set, where $g* = *$ for every $g \in G$, because

$$X \times \{*\} \cong X$$

as G -sets. Finally, Cartesian product distributes over disjoint union:

$$X \times (Y \cup Z) \cong (X \times Y) \cup (X \times Z)$$

as G -sets. Hence multiplication distributes over addition in $B(G)$. Therefore the operations on $S(G)$ extend to make $B(G)$ a commutative ring. $\square$

Exercise 3.13. (cont.) Show that any element of $B(G)$ can be written uniquely as a $\mathbb{Z}$ -linear combination of isomorphism classes of transitive G -sets.

Solution. Let $\mathcal{T}$ be the set of isomorphism classes of transitive finite G -sets. Let X be a finite G -set. By Proposition 3.3, X has a unique decomposition into orbits:

$$X = O_1 \cup \cdots \cup O_r.$$

Each orbit O_i is a transitive G -set, and hence

$$[X] = [O_1] + \cdots + [O_r]$$

in $S(G)$. If several of the orbits are isomorphic as G -sets, we collect their isomorphism classes together. Thus

$$[X] = n_1[T_1] + \cdots + n_k[T_k],$$

where the T_i are pairwise non-isomorphic transitive G -sets and $n_i \in \mathbb{N}$ records the number of orbits of X isomorphic to T_i . This expression is unique in $S(G)$, because the coefficient n_i is exactly the number

of orbits of X whose isomorphism class is $[T_i]$. Therefore $S(G)$ is the free commutative monoid on the set $\mathcal{T}$. Now let $\alpha \in B(G)$. By construction of $B(G)$, we may write

$$\alpha = [X] - [Y]$$

for finite G -sets X and Y . Decompose both X and Y into transitive orbits:

$$[X] = n_1[T_1] + \cdots + n_k[T_k], \quad [Y] = m_1[T_1] + \cdots + m_k[T_k],$$

after enlarging the list of T_i if necessary. Then

$$\alpha = [X] - [Y] = (n_1 - m_1)[T_1] + \cdots + (n_k - m_k)[T_k].$$

Hence every element of $B(G)$ is a $\mathbb{Z}$ -linear combination of isomorphism classes of transitive G -sets. Since $S(G)$ is the free commutative monoid on $\mathcal{T}$, adjoining formal additive inverses turns it into the free abelian group on $\mathcal{T}$. Thus an element of $B(G)$ is determined by its integer coefficient attached to each transitive isomorphism class. Therefore no non-trivial $\mathbb{Z}$ -linear relation among distinct transitive isomorphism classes is possible, and the expression above is unique. $\square$

Exercise 3.14. (cont.) Let $H \leq G$. Show that there is a unique ring homomorphism from $B(G)$ to $\mathbb{Z}$ which sends an isomorphism class $[X]$ to the number of elements of X fixed by H .

Solution. For a finite G -set X , define

$$X^H = \{x \in X \mid hx = x \text{ for every } h \in H\}.$$

We first check that the rule $[X] \mapsto |X^H|$ is well-defined on isomorphism classes. Suppose that $X \cong Y$ as G -sets, and let

$$f: X \rightarrow Y$$

be a G -set isomorphism. If $x \in X^H$, then for every $h \in H$,

$$hf(x) = f(hx) = f(x),$$

so $f(x) \in Y^H$. Applying the same argument to f^{-1} shows that f restricts to a bijection $X^H \rightarrow Y^H$. Hence $|X^H| = |Y^H|$. Define

$$\varphi_H([X]) = |X^H|$$

on $S(G)$. This map respects addition, because

$$(X \cup Y)^H = X^H \cup Y^H$$

as a disjoint union. Hence

$$\varphi_H([X] + [Y]) = |(X \cup Y)^H| = |X^H| + |Y^H| = \varphi_H([X]) + \varphi_H([Y]).$$

It also respects multiplication, because an element $(x, y) \in X \times Y$ is fixed by every element of H iff both x and y are fixed by every element of H . Thus

$$(X \times Y)^H = X^H \times Y^H,$$

and so

$$\varphi_H([X][Y]) = |(X \times Y)^H| = |X^H| |Y^H| = \varphi_H([X])\varphi_H([Y]).$$

The multiplicative identity $[\{*\}]$ is sent to 1, since the one point of $\{*\}$ is fixed by every element of H . Since $B(G)$ is obtained from $S(G)$ by adjoining formal additive inverses, the additive map φ_H extends uniquely to $B(G)$ by

$$\varphi_H([X] - [Y]) = |X^H| - |Y^H|.$$

The multiplication calculation above shows that this extension is multiplicative, so it is a ring homomorphism $B(G) \rightarrow \mathbb{Z}$. Finally, this homomorphism is unique. If $\psi: B(G) \rightarrow \mathbb{Z}$ is any ring homomorphism satisfying $\psi([X]) = |X^H|$ for every finite G -set X , then for every formal difference $[X] - [Y] \in B(G)$,

$$\psi([X] - [Y]) = \psi([X]) - \psi([Y]) = |X^H| - |Y^H|.$$

Thus ψ must agree with φ_H on every element of $B(G)$, proving uniqueness. $\square$

Exercise 3.15. (cont.) Show that any non-zero homomorphism from $B(G)$ to $\mathbb{Z}$ arises from the construction in Exercise 3.14 and that the intersection of the kernels of all such homomorphisms is zero.

Solution. By Proposition 3.4, every transitive finite G -set is isomorphic to G/K for some subgroup $K \leq G$. Here G/K means the set of left cosets of K in G , with G acting by left multiplication; it is not being used as a quotient group. Thus, by Exercise 3.13, the elements $[G/K]$, with K running over representatives of conjugacy classes of subgroups, generate $B(G)$ additively.

Let $\psi: B(G) \rightarrow \mathbb{Z}$ be a non-zero ring homomorphism. Since $[G/G]$ is the multiplicative identity of $B(G)$, we have

$$\psi([G/G])^2 = \psi([G/G]^2) = \psi([G/G]).$$

The only integers satisfying $n^2 = n$ are 0 and 1. Hence $\psi([G/G])$ is either 0 or 1. If it were 0, then for every $\alpha \in B(G)$,

$$\psi(\alpha) = \psi([G/G]\alpha) = \psi([G/G])\psi(\alpha) = 0,$$

contrary to the assumption that ψ is non-zero. Thus $\psi([G/G]) = 1$. Choose a subgroup $H \leq G$ with $\psi([G/H]) \neq 0$ and with $|H|$ minimal among all such subgroups. We claim that $\psi = \varphi_H$, where φ_H is the fixed-point homomorphism from Exercise 3.14. Let $K \leq G$. We decompose the product $G/H \times G/K$ into G -orbits using Proposition 3.3, where G acts diagonally:

$$u(aH, bK) = (uaH, ubK).$$

Since the first coordinate of an element of $G/H \times G/K$ has the form aH , acting by a^{-1} sends (aH, bK) to

$$(H, a^{-1}bK).$$

Hence every G -orbit has a representative of the form (H, gK) . We now compute the stabilizer of this representative. An element $u \in G$ fixes (H, gK) iff

$$u(H, gK) = (H, gK),$$

which is equivalent to the two conditions

$$uH = H \quad \text{and} \quad ugK = gK.$$

The first condition says $u \in H$. The second condition says $g^{-1}ug \in K$, or equivalently $u \in gKg^{-1}$. Therefore the stabilizer is

$$H \cap gKg^{-1}.$$

By Proposition 3.4, the orbit of (H, gK) is therefore isomorphic to

$$G/(H \cap gKg^{-1}).$$

Combining the orbit decomposition from Proposition 3.3 with the orbit-stabilizer identification from Proposition 3.4, we get

$$[G/H][G/K] = \sum [G/(H \cap gKg^{-1})],$$

where the sum runs over representatives of the H -orbits on G/K . Applying ψ gives

$$\psi([G/H])\psi([G/K]) = \sum \psi([G/(H \cap gKg^{-1})]).$$

If $H \cap gKg^{-1}$ is a proper subgroup of H , then its order is smaller than $|H|$, so its term is zero by the minimal choice of H . The only remaining terms occur when

$$H \cap gKg^{-1} = H,$$

equivalently when $H \leq gKg^{-1}$. This is exactly the condition that the coset gK is fixed by every element of H . Indeed, gK is fixed by H iff for every $h \in H$,

$$hgK = gK.$$

Multiplying on the left by g^{-1} , this is equivalent to

$$g^{-1}hgK = K,$$

which holds iff $g^{-1}hg \in K$. Thus gK is fixed by every element of H iff $g^{-1}Hg \leq K$, equivalently iff $H \leq gKg^{-1}$. Hence the number of remaining terms is $|(G/K)^H|$, and the equality above becomes

$$\psi([G/H])\psi([G/K]) = |(G/K)^H|\psi([G/H]).$$

Since $\psi([G/H]) \neq 0$, we may cancel it in $\mathbb{Z}$, obtaining

$$\psi([G/K]) = |(G/K)^H| = \varphi_H([G/K]).$$

Since the classes $[G/K]$ generate $B(G)$ additively, this proves that $\psi = \varphi_H$. Thus every non-zero homomorphism $B(G) \rightarrow \mathbb{Z}$ arises from the construction in Exercise 3.14.

It remains to prove that the intersection of the kernels is zero. Let

$$\alpha = \sum_K a_K [G/K] \in B(G),$$

where K runs over one representative from each conjugacy class of subgroups of G . Suppose that $\alpha \neq 0$. Choose K with $a_K \neq 0$ and with $|K|$ maximal among all subgroups appearing with non-zero coefficient. We evaluate at the fixed-point homomorphism φ_K . For another subgroup L , the set $(G/L)^K$ is non-empty iff some coset gL is fixed by K , which is equivalent to

$$g^{-1}Kg \leq L.$$

Thus, if $(G/L)^K$ is non-empty, then $|K| \leq |L|$. If also $a_L \neq 0$, maximality of $|K|$ gives $|L| \leq |K|$, and hence $|L| = |K|$. Therefore $g^{-1}Kg = L$, so L is conjugate to K . Since we chose one representative from each conjugacy class, this forces $L = K$. Hence all non-zero coefficient terms except the K -term make no contribution to $\varphi_K(\alpha)$, and

$$\varphi_K(\alpha) = a_K |(G/K)^K|.$$

The coset $K \in G/K$ is fixed by every element of K , so $|(G/K)^K| > 0$. Since $a_K \neq 0$, we have $\varphi_K(\alpha) \neq 0$. Therefore α is not in the kernel of φ_K . Thus no non-zero element of $B(G)$ lies in the kernels of all fixed-point homomorphisms, and so the intersection of all these kernels is zero. $\square$

An element e of a ring R is called an *idempotent* if $e^2 = e$.

Exercise 3.16. (cont.) Show that if G has no self-normalizing proper subgroup, then $B(G)$ has no idempotents other than the additive and multiplicative identities. (A subgroup H of G is called *self-normalizing* if $N_G(H) = H$. We shall see in Section 11 that there is an important class of groups, namely the nilpotent groups, that have no self-normalizing proper subgroups.)

Solution. Let $e \in B(G)$ be an idempotent, so $e^2 = e$. For every subgroup $H \leq G$, let

$$\varphi_H: B(G) \rightarrow \mathbb{Z}$$

be the fixed-point homomorphism defined by

$$\varphi_H([X]) = |X^H|$$

for every finite G -set X . Since φ_H is a ring homomorphism,

$$\varphi_H(e)^2 = \varphi_H(e^2) = \varphi_H(e).$$

Hence $\varphi_H(e)$ is an integer satisfying $n^2 = n$, so

$$\varphi_H(e) \in \{0, 1\}.$$

We next show that these values do not change when we enlarge a subgroup inside its normalizer. Suppose that

$$H < K \leq N_G(H)$$

and that $|K : H| = p$ is prime. Since $K \leq N_G(H)$, the quotient group K/H acts on X^H for every finite G -set X . The fixed points of this action are exactly X^K . Since K/H has prime order p , Corollary 3.5 shows that each orbit of K/H on X^H has size either 1 or p . The orbits of size 1 are precisely the points of X^K , while every other orbit has size p . Hence, for some integer $r \geq 0$,

$$|X^H| = |X^K| + rp.$$

Therefore

$$|X^H| \equiv |X^K| \pmod{p}.$$

By additivity, the same congruence holds for every element of $B(G)$; in particular,

$$\varphi_H(e) \equiv \varphi_K(e) \pmod{p}.$$

Since both $\varphi_H(e)$ and $\varphi_K(e)$ are either 0 or 1, this congruence forces

$$\varphi_H(e) = \varphi_K(e).$$

Now let $H < G$ be a proper subgroup. By hypothesis, H is not self-normalizing, so

$$H < N_G(H).$$

Thus $N_G(H)/H$ is a non-trivial finite group. Choose a subgroup K/H of prime order in $N_G(H)/H$. Then

$$H < K \leq N_G(H),$$

and the previous paragraph gives $\varphi_H(e) = \varphi_K(e)$. If $K < G$, we repeat the same argument with K in place of H . Since G is finite, this process must eventually reach G . Hence

$$\varphi_H(e) = \varphi_G(e)$$

for every subgroup $H \leq G$.

Therefore all fixed-point homomorphisms take the same value on e , and this common value is either 0 or 1. If it is 0, then e lies in the kernel of every φ_H , so Exercise 3.15 gives $e = 0$. If it is 1, then for every $H \leq G$,

$$\varphi_H(e - [G/G]) = \varphi_H(e) - \varphi_H([G/G]) = 1 - 1 = 0,$$

since G/G is the one-point G -set. Again by Exercise 3.15, we get

$$e - [G/G] = 0.$$

Thus $e = [G/G]$, the multiplicative identity of $B(G)$. Therefore the only idempotents in $B(G)$ are 0 and 1. □

Chapter 2

The General Linear Group

4 Basic Structure

Setup and notation

- F a field, $n \in \mathbb{N}$. $\mathcal{M}_n(F)$ denotes the set of all $n \times n$ matrices with entries in F ; we write $M = (m_{ij})$ with $m_{ij} \in F$ the (i, j) -entry (row i , column j).
- The *general linear group* $\mathrm{GL}(n, F)$ is the subset of $\mathcal{M}_n(F)$ of invertible matrices, equivalently those with non-zero determinant. It forms a group under matrix multiplication; I denotes the identity.
- More generally, for a finite-dimensional F -vector space V , $\mathrm{GL}(V)$ is the group of invertible linear transformations of V under composition. Taking $V = F^n$ recovers $\mathrm{GL}(n, F)$ up to obvious isomorphism, and any n -dimensional F -vector space is isomorphic to F^n , so we lose nothing by working with $\mathrm{GL}(n, F)$.
- If F is finite, then $|F| = p^a$ for a prime p and $a \in \mathbb{N}$ (Moore's theorem, 1893). For q a prime power we write $\mathrm{GL}(n, q)$ for $\mathrm{GL}(n, F)$ where F is the unique field of order q .

Proposition 4.1. *Let E be a finite abelian group of exponent p , where p is prime. Then $\mathrm{Aut}(E) \cong \mathrm{GL}(n, p)$, where $n \in \mathbb{N}$ is such that $|E| = p^n$.*

(Recall that the exponent of a group is the least common multiple of the orders of its elements.)

Proof. Let $K = \mathbb{Z}/p\mathbb{Z}$. We give E the structure of a K -vector space. Since E is abelian, we may define vector addition by

$$x + y = xy$$

for $x, y \in E$, with the identity element of E serving as the zero vector. If $a \in K$, choose an integer representative r of a , and define scalar multiplication by

$$ax = x^r.$$

This is well-defined: if $r \equiv s \pmod{p}$, then $r - s = mp$ for some $m \in \mathbb{Z}$, and since E has exponent p we have $x^{r-s} = (x^p)^m = e$, so $x^r = x^s$.

The vector-space axioms follow from the group law in E . For instance, if $a \in K$ is represented by r , then

$$a(x + y) = (xy)^r = x^r y^r = ax + ay,$$

where we use that E is abelian. The remaining compatibility laws are checked similarly from the identities $x^{r+s} = x^r x^s$, $x^{rs} = (x^s)^r$, and $x^1 = x$. Thus E is a vector space over K .

Now every group endomorphism of E is automatically K -linear. Indeed, if $\varphi \in \mathrm{End}(E)$ and $a \in K$ is represented by r , then

$$\varphi(x + y) = \varphi(xy) = \varphi(x)\varphi(y) = \varphi(x) + \varphi(y)$$

and

$$\varphi(ax) = \varphi(x^r) = \varphi(x)^r = a\varphi(x).$$

Conversely, every K -linear transformation of E is a homomorphism of the underlying abelian group, because it preserves vector addition, which is precisely the group operation on E . Hence the automorphisms of E as a group are exactly the invertible K -linear transformations of E . If $\dim_K E = n$, then $|E| = |K|^n = p^n$. Therefore

$$\text{Aut}(E) \cong \text{GL}(E) \cong \text{GL}(n, p),$$

as required. $\square$

Elementary abelian p -groups

- If E is as in Proposition 4.1 with basis $\{x_1, \dots, x_n\}$ over the field of p elements, then as groups $E = \langle x_1 \rangle \times \dots \times \langle x_n \rangle$ with each $\langle x_i \rangle$ cyclic of order p .
- Hence any finite abelian group of prime exponent p is isomorphic with a direct product of copies of $\mathbb{Z}_p$. Such groups are called *elementary abelian p -groups*.
- The *rank* of an elementary abelian p -group E is n , where $|E| = p^n$.
- In particular $\text{Aut}(\mathbb{Z}_p) \cong \text{GL}(1, p) \cong (\mathbb{Z}/p\mathbb{Z})^\times$ by Proposition 4.1; this was previously established in Proposition 2.1.

Proposition 4.2. *Let $n \in \mathbb{N}$, and let q be a prime power. Then*

$$|\text{GL}(n, q)| = \prod_{k=1}^n (q^n - q^{k-1}) = q^{\frac{n(n-1)}{2}} (q^n - 1) \cdots (q - 1).$$

Proof. Let F be the field of order q . An $n \times n$ matrix over F is invertible if and only if its rows form a linearly independent subset of F^n . We therefore count the number of ways to choose the rows of such a matrix.

The first row may be any non-zero vector of F^n , so there are $q^n - 1$ choices. Suppose now that $1 < k \leq n$ and that the first $k - 1$ rows have already been chosen to be linearly independent. Their span is a $(k - 1)$ -dimensional subspace of F^n , and hence contains q^{k-1} vectors. The k th row must avoid this span, and so there are

$$q^n - q^{k-1}$$

choices for it.

Multiplying these independent choices gives

$$|\text{GL}(n, q)| = \prod_{k=1}^n (q^n - q^{k-1}).$$

Finally, since

$$q^n - q^{k-1} = q^{k-1}(q^{n-k+1} - 1),$$

we obtain

$$\prod_{k=1}^n (q^n - q^{k-1}) = q^{0+1+\dots+(n-1)} (q^n - 1)(q^{n-1} - 1) \cdots (q - 1) = q^{\frac{n(n-1)}{2}} (q^n - 1) \cdots (q - 1).$$

$\square$

Fixing notation for the section

- We now fix a field F and some $n \in \mathbb{N}$, and write G instead of $\text{GL}(n, F)$.
- For $M = (m_{ij}) \in \mathcal{M}_n(F)$: the *main diagonal* of M consists of the entries m_{ii} for $1 \leq i \leq n$.
- M is *diagonal* if its only non-zero entries lie on the main diagonal.
- M is *upper triangular* if all entries below the main diagonal are zero, i.e. $m_{ij} = 0$ whenever $i > j$.

Proposition 4.3. *The set B consisting of all invertible upper triangular matrices is a subgroup of G , called the standard Borel subgroup.*

Proof. The identity matrix lies in B . If $M = (m_{ij})$ and $N = (n_{ij})$ are upper triangular, then for $i > j$ the (i, j) -entry of MN is

$$\sum_{k=1}^n m_{ik}n_{kj}.$$

If $k < i$, then $m_{ik} = 0$, while if $k \geq i > j$, then $n_{kj} = 0$. Hence every term in the sum is zero, so MN is again upper triangular. Since M and N are invertible, MN is also invertible; hence $MN \in B$.

It remains to show that inverses of elements of B again lie in B . Let $M = (m_{ij}) \in B$, and let $N = (n_{ij}) = M^{-1}$. Since M is upper triangular, its determinant is the product of its diagonal entries. As M is invertible, this determinant is non-zero, and hence each m_{ii} is non-zero.

We show that $n_{ij} = 0$ whenever $i > j$. Since $MN = I$, for each i and j we have

$$\sum_{k=1}^n m_{ik}n_{kj} = \delta_{ij}.$$

Fix j . Taking $i = n$ and $j < n$ gives $m_{nn}n_{nj} = 0$, and hence $n_{nj} = 0$. Now argue upward from the bottom row. Suppose that for some i we already know $n_{kj} = 0$ whenever $k > i$ and $j < k$. If $j < i$, then the (i, j) -entry of $MN = I$ gives

$$0 = \sum_{k=1}^n m_{ik}n_{kj} = m_{ii}n_{ij} + \sum_{k=i+1}^n m_{ik}n_{kj}.$$

The sum on the right is zero by the induction hypothesis, so $m_{ii}n_{ij} = 0$. Since $m_{ii} \neq 0$, it follows that $n_{ij} = 0$. Thus N is upper triangular.

Therefore B contains the identity, is closed under multiplication, and is closed under inverses. Hence $B \leq G$. $\square$

More generally, a *Borel subgroup* of G is any conjugate of the standard Borel subgroup B .

Permutation matrices

- A *permutation matrix* is a matrix in which every row and column has a unique non-zero entry, and all non-zero entries equal 1.
- The identity matrix is a permutation matrix; every permutation matrix is obtained from I by switching columns (or rows).
- Every permutation matrix is orthogonal, so its inverse (= transpose) is again a permutation matrix. Hence all $n \times n$ permutation matrices lie in G .

Proposition 4.4. *The set W consisting of all permutation matrices is a subgroup of G , called the Weyl subgroup.*

Proof. We already know that the identity matrix lies in W , and that the inverse of a permutation matrix is again a permutation matrix. It remains to check closure under multiplication. Let $M = (m_{ij})$ and $N = (n_{ij})$ be permutation matrices, and set $P = MN = (p_{ij})$. Then

$$p_{ij} = \sum_{k=1}^n m_{ik}n_{kj}.$$

Fix a row i . Since M is a permutation matrix, there is a unique k_i such that $m_{ik_i} = 1$, and all other entries in row i of M are zero. Hence

$$p_{ij} = n_{k_i j}$$

for every j . The k_i th row of N has exactly one non-zero entry, and that entry is 1, so the i th row of P has exactly one non-zero entry, also equal to 1.

Similarly, fix a column j . Since N is a permutation matrix, there is a unique l_j such that $n_{l_j j} = 1$, and all other entries in column j of N are zero. Hence

$$p_{ij} = m_{il_j}$$

for every i . The l_j th column of M has exactly one non-zero entry, and that entry is 1, so the j th column of P has exactly one non-zero entry, also equal to 1. Thus P is a permutation matrix. Therefore W is closed under multiplication, and so $W \leq G$. $\square$

Action on the standard basis

- Let $V_n(F)$ denote the vector space of n -dimensional column vectors with entries in F , and let $\mathbf{v}_1, \dots, \mathbf{v}_n$ be the standard basis.
- Multiplying $\mathbf{v}_i$ on the left by an $n \times n$ matrix M produces the i th column of M ; we say that M sends $\mathbf{v}_i$ to the i th column of M .

Proposition 4.5. $W \cong S_n$.

Proof. Let $X = \{1, \dots, n\}$. For each $w \in W$ and each $i \in X$, the i th column of w is some standard basis vector $\mathbf{v}_k$. Define

$$\varphi(w)(i) = k$$

where $w\mathbf{v}_i = \mathbf{v}_k$. Since w is a permutation matrix, each standard basis vector appears exactly once among the columns of w . Hence $\varphi(w)$ is a bijection $X \rightarrow X$, so $\varphi(w) \in S_n$. This gives a map $\varphi: W \rightarrow S_n$.

If $w, u \in W$, then for each $i \in X$ we have

$$u\mathbf{v}_i = \mathbf{v}_{\varphi(u)(i)}$$

and therefore

$$wu\mathbf{v}_i = w\mathbf{v}_{\varphi(u)(i)} = \mathbf{v}_{\varphi(w)(\varphi(u)(i))}.$$

Thus $\varphi(wu) = \varphi(w) \circ \varphi(u)$, and φ is a homomorphism.

If $\varphi(w) = \varphi(u)$, then $w\mathbf{v}_i = u\mathbf{v}_i$ for every i , so w and u have the same columns. Hence $w = u$, and φ is injective. Conversely, if $\sigma \in S_n$, let w_σ be the matrix whose i th column is $\mathbf{v}_{\sigma(i)}$ for each i . Then w_σ is a permutation matrix and $\varphi(w_\sigma) = \sigma$, so φ is surjective. Therefore $W \cong S_n$. $\square$

Permutation matrices as permutations

- We often implicitly regard a permutation matrix w as the element of S_n sending i to j , where $\mathbf{v}_j$ is the i th column of w .
- For example, the matrix

$$\begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}$$

corresponds to $(1\ 3)(2\ 4) \in S_4$, and so if w is this matrix then we may write $w(1) = 3$, and so forth.

- Useful observation: if $w(i) = j$, then for any $M \in \mathcal{M}_n(F)$, the j th row of wM equals the i th row of M , and the i th column of Mw equals the j th column of M .

Transvections

- For distinct $1 \leq i, j \leq n$ and $\alpha \in F$, define $X_{ij}(\alpha)$ to be the $n \times n$ matrix whose (k, l) -entry equals α if $(k, l) = (i, j)$ and equals δ_{kl} for all other (k, l) .
- For example, $X_{23}(\alpha) \in \mathcal{M}_3(F)$ is the matrix

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & \alpha \\ 0 & 0 & 1 \end{pmatrix}.$$

- These matrices $X_{ij}(\alpha)$, and their conjugates by elements of G , are called *transvections*.

Lemma 4.6. Let $\alpha, \beta \in F$, and let i and j be distinct.

- (i) $X_{ij}(\alpha)$ has determinant 1 and hence lies in G .
- (ii) If $\alpha \neq 0$, then $X_{ij}(\alpha) \in B$ iff $i < j$.
- (iii) $X_{ij}(\alpha)X_{ij}(\beta) = X_{ij}(\alpha + \beta)$, and hence $X_{ij}(\alpha)^{-1} = X_{ij}(-\alpha)$.

(iv) $[X_{ij}(\alpha), X_{jk}(\beta)] = X_{ik}(\alpha\beta)$ whenever i, j, k are distinct.

(v) If $w \in W$, then $wX_{ij}(\alpha)w^{-1} = X_{w(i)w(j)}(\alpha)$.

(vi) $X_{ij}(\alpha)$ sends $\mathbf{v}_j$ to $\mathbf{v}_j + \alpha\mathbf{v}_i$ and fixes $\mathbf{v}_k$ whenever $k \neq j$.

(vii) If $M \in \mathcal{M}_n(F)$, then the i th row of $X_{ij}(\alpha)M$ is equal to the sum of the i th row of M and α times the j th row of M , and for $k \neq i$ the k th row of $X_{ij}(\alpha)M$ is equal to the k th row of M .

Proof. For this proof, write E_{rs} for the matrix with 1 in the (r, s) -entry and 0 everywhere else. Thus

$$X_{ij}(\alpha) = I + \alpha E_{ij}.$$

For (i), if $i < j$, then $X_{ij}(\alpha)$ is upper triangular with every diagonal entry equal to 1; if $i > j$, then it is lower triangular with every diagonal entry equal to 1. In either case its determinant is 1, and hence $X_{ij}(\alpha) \in G$.

For (ii), assume $\alpha \neq 0$. If $i < j$, then the only possible non-zero off-diagonal entry of $X_{ij}(\alpha)$ lies above the main diagonal, so $X_{ij}(\alpha) \in B$. If $i > j$, then the (i, j) -entry lies below the main diagonal and is non-zero, so $X_{ij}(\alpha)$ is not upper triangular. Hence $X_{ij}(\alpha) \in B$ iff $i < j$.

For (iii), since $i \neq j$ we have $E_{ij}^2 = 0$. Therefore

$$X_{ij}(\alpha)X_{ij}(\beta) = (I + \alpha E_{ij})(I + \beta E_{ij}) = I + (\alpha + \beta)E_{ij} = X_{ij}(\alpha + \beta).$$

Taking $\beta = -\alpha$ gives $X_{ij}(\alpha)X_{ij}(-\alpha) = I$, and hence $X_{ij}(\alpha)^{-1} = X_{ij}(-\alpha)$.

For (iv), suppose that i, j, k are distinct. Using the convention $[x, y] = xyx^{-1}y^{-1}$ and part (iii), we compute

$$\begin{aligned} [X_{ij}(\alpha), X_{jk}(\beta)] &= (I + \alpha E_{ij})(I + \beta E_{jk})(I - \alpha E_{ij})(I - \beta E_{jk}) \\ &= (I + \alpha E_{ij} + \beta E_{jk} + \alpha\beta E_{ik})(I - \alpha E_{ij})(I - \beta E_{jk}) \\ &= (I + \beta E_{jk} + \alpha\beta E_{ik})(I - \beta E_{jk}) \\ &= I + \alpha\beta E_{ik} \\ &= X_{ik}(\alpha\beta). \end{aligned}$$

Here we used $E_{ij}E_{jk} = E_{ik}$, together with the fact that the other products of these matrix units which occur in the calculation vanish because i, j, k are distinct.

For (v), let $w \in W$. Since we are regarding w as the corresponding permutation of $\{1, \dots, n\}$, we have $w\mathbf{v}_r = \mathbf{v}_{w(r)}$ for every r . If $r = w(j)$, then

$$wX_{ij}(\alpha)w^{-1}\mathbf{v}_r = wX_{ij}(\alpha)\mathbf{v}_j = w(\mathbf{v}_j + \alpha\mathbf{v}_i) = \mathbf{v}_{w(j)} + \alpha\mathbf{v}_{w(i)}.$$

If $r \neq w(j)$, then $w^{-1}(r) \neq j$, so $X_{ij}(\alpha)$ fixes $\mathbf{v}_{w^{-1}(r)}$, and hence $wX_{ij}(\alpha)w^{-1}$ fixes $\mathbf{v}_r$. Thus $wX_{ij}(\alpha)w^{-1}$ sends $\mathbf{v}_{w(j)}$ to $\mathbf{v}_{w(j)} + \alpha\mathbf{v}_{w(i)}$ and fixes all other standard basis vectors. Therefore

$$wX_{ij}(\alpha)w^{-1} = X_{w(i)w(j)}(\alpha).$$

For (vi), the j th column of $X_{ij}(\alpha)$ is $\mathbf{v}_j + \alpha\mathbf{v}_i$, while its k th column is $\mathbf{v}_k$ whenever $k \neq j$. Since multiplying $\mathbf{v}_k$ on the left by a matrix gives the k th column of that matrix, the asserted action on the standard basis follows.

For (vii), let $M = (m_{rs}) \in \mathcal{M}_n(F)$, and write $X_{ij}(\alpha)M = (p_{rs})$. Then

$$p_{rs} = \sum_{t=1}^n (X_{ij}(\alpha))_{rt} m_{ts}.$$

If $r = i$, this gives $p_{is} = m_{is} + \alpha m_{js}$ for every s , so the i th row of $X_{ij}(\alpha)M$ is the i th row of M plus α times the j th row of M . If $r \neq i$, then the r th row of $X_{ij}(\alpha)$ is the r th row of the identity matrix, so $p_{rs} = m_{rs}$ for every s . Hence every row other than the i th row is unchanged. $\square$

Root subgroups

- For distinct i and j , define $X_{ij} = \{X_{ij}(\alpha) \mid \alpha \in F\}$; by parts (i) and (iii) of Lemma 4.6, this is a subgroup of G .
- The subgroups X_{ij} are called *root subgroups* of G . (Terminology from the theory of Lie algebras.)

- These subgroups feed into the Bruhat Decomposition Theorem, the main result of the section.

Lemma 4.7. *Let $M \in G$. Then there is a product b of upper triangular transvections such that the following property holds: For each $1 \leq i \leq n$, bM has exactly one row, say the k_i th row, whose entries in the first $i - 1$ columns are zero and which has a non-zero entry in the i th column.*

Proof. Let $M = (m_{ij}) \in G$. Since M is invertible, its first column is not the zero column. Choose k_1 maximal such that $m_{k_1,1} \neq 0$; equivalently, $m_{i,1} = 0$ for all $i > k_1$. For example, if $n = 5$ and $k_1 = 3$, then M has the schematic form

$$M = \begin{pmatrix} * & * & * & * & * \\ * & * & * & * & * \\ * & * & * & * & * \\ 0 & * & * & * & * \\ 0 & * & * & * & * \end{pmatrix},$$

where the symbol $*$ denotes an entry whose value is not relevant to the argument.

We now premultiply by transvections of the form $X_{ik_1}(\alpha)$ with $i < k_1$, choosing $\alpha = -m_{i,1}m_{k_1,1}^{-1}$ in order to clear the $(i, 1)$ -entry. By Lemma 4.6(vii), this replaces row i by $(\text{row } i) + \alpha(\text{row } k_1)$; since row k_1 has first entry $m_{k_1,1}$, the new first entry in row i is 0. Each such transvection is upper triangular because $i < k_1$. Thus we obtain a matrix $M' = (m'_{ij})$ whose only non-zero entry in the first column lies in row k_1 . In the example above,

$$M' = \begin{pmatrix} 0 & * & * & * & * \\ 0 & * & * & * & * \\ * & * & * & * & * \\ 0 & * & * & * & * \\ 0 & * & * & * & * \end{pmatrix}.$$

All transvections used so far lie in B , and the resulting matrix is still invertible.

Since M' is invertible, its first two columns are linearly independent. Therefore the second column of M' has a non-zero entry in some row other than the k_1 th row. Choose $k_2 \neq k_1$ maximal such that $m'_{k_2,2} \neq 0$ and $m'_{i,2} = 0$ for all $i > k_2$ with $i \neq k_1$. We premultiply M' by transvections of the form $X_{ik_2}(\alpha)$, where $i < k_2$ and $i \neq k_1$, so as to clear all entries in the second column except those in rows k_1 and k_2 . These transvections are again upper triangular, and they do not disturb the first column because row k_2 has zero first entry.

We continue in this way. Suppose that the rows $k_1, \dots, k_{r-1}$ have already been chosen, and that after the preceding operations the first $r - 1$ columns have the required shape. Since the matrix is still invertible, the first r columns are linearly independent, so the r th column must have a non-zero entry in some row not among $k_1, \dots, k_{r-1}$. Choose k_r maximal among such rows. Then premultiply by transvections $X_{ik_r}(\alpha)$ with $i < k_r$ and $i \notin \{k_1, \dots, k_{r-1}\}$ to clear the other entries in the r th column among the rows whose first $r - 1$ entries are zero. Because row k_r has zero entries in the first $r - 1$ columns, these operations preserve all earlier columns.

After n stages we obtain a matrix bM , where b is a product of upper triangular transvections, with the desired property. For example, if $n = 5$ and

$$(k_1, k_2, k_3, k_4, k_5) = (3, 5, 4, 1, 2),$$

then the final shape is

$$bM = \begin{pmatrix} 0 & 0 & 0 & * & * \\ 0 & 0 & 0 & 0 & * \\ * & * & * & * & * \\ 0 & 0 & * & * & * \\ 0 & * & * & * & * \end{pmatrix}.$$

Of course, the numbers $k_1, \dots, k_n$ are just $1, \dots, n$ in some order. Thus for each $1 \leq i \leq n$, the matrix bM has exactly one row, namely row k_i , whose entries in the first $i - 1$ columns are zero and whose entry in the i th column is non-zero. $\square$

Theorem (Bruhat Decomposition Theorem). $G = BWB$.

Proof. Let $M \in G$. By Lemma 4.7, there is a product b of upper triangular transvections such that, for each $1 \leq i \leq n$, the matrix bM has exactly one row, say the k_i th row, whose entries in the first $i - 1$ columns are zero and whose entry in the i th column is non-zero. Since each upper triangular transvection

lies in B , and since $B \leq G$ by Proposition 4.3, we have $b \in B$.

Let $w \in W$ be the permutation matrix whose k_i th column is $\mathbf{v}_i$ for each i . Equivalently, $w(k_i) = i$. By the row-permutation observation above, the i th row of wbM is equal to the k_i th row of bM . Hence the entries of the i th row of wbM in columns $1, \dots, i-1$ are all zero. This holds for every i , so wbM is upper triangular. Since w , b , and M are invertible, wbM is also invertible; therefore $wbM \in B$.

It follows that

$$M = b^{-1}w^{-1}(wbM) \in BWB.$$

Thus $G \subseteq BWB$. The reverse inclusion is immediate from $B \leq G$ and $W \leq G$, so $G = BWB$. $\square$

We have now expressed G as a union of the double cosets BwB , as w ranges through W . We will now show that this union is disjoint. Again we first need a lemma.

Lemma 4.8. *If $w_1, w_2 \in W$ and $b \in B$ are such that $w_1bw_2 \in B$, then $w_2 = w_1^{-1}$.*

Proof. Fix $j \in \{1, \dots, n\}$. Choose i such that $w_1(i) = j$, and then choose k such that $w_2(k) = i$. With our convention for permutation matrices, $w_1(i) = j$ means that left multiplication by w_1 sends row i to row j . Thus the j th row of w_1b is the i th row of b . Also, since $w_2(k) = i$, the k th column of $(w_1b)w_2$ is the i th column of w_1b . Therefore the (j, k) -entry of w_1bw_2 is the (i, i) -entry of b .

Since $b \in B$, its diagonal entries are non-zero, so this (j, k) -entry of w_1bw_2 is non-zero. But $w_1bw_2 \in B$ is upper triangular, and hence a non-zero (j, k) -entry can occur only when $j \leq k$. By construction,

$$k = w_2^{-1}(i) = w_2^{-1}w_1^{-1}(j).$$

Thus the permutation $\sigma = w_2^{-1}w_1^{-1}$ satisfies $\sigma(j) \geq j$ for every j . Summing these inequalities over all j gives

$$\sum_{j=1}^n \sigma(j) \geq \sum_{j=1}^n j.$$

The two sums are equal because σ is a permutation of $\{1, \dots, n\}$, so every inequality $\sigma(j) \geq j$ must in fact be an equality. Hence $\sigma(j) = j$ for every j , and so $\sigma = 1$. Therefore $w_2^{-1}w_1^{-1} = 1$, which is equivalent to $w_2 = w_1^{-1}$. $\square$

Corollary 4.9. *If w and w' are distinct elements of W , then BwB and $Bw'B$ are disjoint; consequently, G is the disjoint union as w runs through W of the $n!$ double cosets BwB .*

Proof. Suppose first that $BwB \cap Bw'B$ is non-empty. Since double cosets are either equal or disjoint, this gives $BwB = Bw'B$. In particular, $w' \in BwB$, so there exist $b, b' \in B$ such that

$$w' = bw b'.$$

Rearranging, we obtain

$$w^{-1}b^{-1}w' = b' \in B.$$

Applying Lemma 4.8 with $w_1 = w^{-1}$, $b = b^{-1}$, and $w_2 = w'$, we get

$$w' = (w^{-1})^{-1} = w.$$

Thus if $w \neq w'$, then BwB and $Bw'B$ are disjoint.

By Bruhat Decomposition Theorem, every element of G lies in BwB , so

$$G = \bigcup_{w \in W} BwB.$$

The disjointness just proved shows that this is a disjoint union. Finally, by Proposition 4.5, we have $W \cong S_n$, and hence $|W| = n!$. Therefore G is the disjoint union of the $n!$ double cosets BwB . $\square$

The following somewhat technical observation, which we shall need in the next section, is an immediate consequence of what we have done.

Corollary 4.10. *Let $M \in G$, and let w be the unique element of W such that $M \in BwB$. Then w sends $\mathbf{v}_i$ to $\mathbf{v}_{k_i}$ for each i , where the numbers k_i are as defined in the statement of Lemma 4.7. In particular, if M sends $\mathbf{v}_1$ to $\alpha_1\mathbf{v}_1 + \dots + \alpha_k\mathbf{v}_k$ where $\alpha_k \neq 0$, then w sends $\mathbf{v}_1$ to $\mathbf{v}_k$.*

Proof. In the proof of Bruhat Decomposition Theorem, Lemma 4.7 gives row indices $k_1, \dots, k_n$ and an element $b \in B$ such that bM has the prescribed row-echelon form. The permutation matrix used there is the matrix $p \in W$ satisfying

$$p(k_i) = i$$

for every i , or equivalently, the k_i th column of p is $\mathbf{v}_i$. With this choice, left multiplication by p moves row k_i of bM to row i , and hence $pbM \in B$. It follows that

$$M \in Bp^{-1}B.$$

By the uniqueness of the Bruhat double coset from Corollary 4.9, the unique element $w \in W$ such that $M \in BwB$ is $w = p^{-1}$. Therefore

$$w(i) = p^{-1}(i) = k_i,$$

and so $w\mathbf{v}_i = \mathbf{v}_{k_i}$ for each i .

For the final assertion, $M\mathbf{v}_1$ is the first column of M . If

$$M\mathbf{v}_1 = \alpha_1\mathbf{v}_1 + \dots + \alpha_k\mathbf{v}_k$$

with $\alpha_k \neq 0$ and no later terms, then k is precisely the lowest row in which the first column of M has a non-zero entry. Thus the index selected in Lemma 4.7 for the first column is $k_1 = k$, and the result above gives $w\mathbf{v}_1 = \mathbf{v}_k$. $\square$

We close this section with a result giving a smaller generating set for G than that given by the Bruhat decomposition. Once again, we start with a lemma.

Lemma 4.11. *Let $b \in B$. Then there exists a product t of transvections such that tb is a diagonal matrix having the same main diagonal entries as b .*

Proof. Write $b = (b_{ij})$. Since $b \in B$, it is upper triangular and invertible, so each diagonal entry b_{ii} is non-zero. We shall clear the entries above the main diagonal by premultiplying by transvections, beginning at the far right. This direction is important: operations used to clear column j will add multiples of row j to rows above it, and if we have already treated columns to the right of j , then row j has zero entries in those right-hand columns except at its own diagonal position where appropriate, so the lower-right block is not disturbed.

For example, in the 3×3 case, clearing the $(1, 2)$ -entry first would give

$$X_{12}(-ad_2^{-1}) \begin{pmatrix} d_1 & a & c \\ 0 & d_2 & e \\ 0 & 0 & d_3 \end{pmatrix} = \begin{pmatrix} d_1 & 0 & c - ad_2^{-1}e \\ 0 & d_2 & e \\ 0 & 0 & d_3 \end{pmatrix},$$

so an operation aimed at column 2 can change an entry in column 3. This is precisely what the right-to-left procedure avoids. By contrast, within one fixed column the relevant transvections may be applied in any order: they add the same pivot row to different rows, and they leave that pivot row unchanged. Start with column n . For each $i < n$, premultiply by

$$X_{in}(-b_{in}b_{nn}^{-1}).$$

By Lemma 4.6(vii), this adds $-b_{in}b_{nn}^{-1}$ times row n to row i , and therefore clears the (i, n) -entry. These operations do not change any diagonal entry: row n has zeros in columns $1, \dots, n-1$, so adding a multiple of row n to row $i < n$ does not affect the (i, i) -entry, and row n itself is unchanged.

Now move to column $n-1$. Using transvections of the form $X_{i,n-1}(\alpha)$ with $i < n-1$, and choosing α so as to cancel the current $(i, n-1)$ -entry, we clear all entries above the $(n-1, n-1)$ -entry. These operations do not alter the diagonal entries, and they do not reintroduce entries in column n , because after the previous stage row $n-1$ has zero in column n .

Continuing leftward in this way, at the stage for column j we use transvections $X_{ij}(\alpha)$ with $i < j$ to clear the entries above the (j, j) -entry. The lower-right block formed by columns and rows $j+1, \dots, n$ has already been put in diagonal form, and it is not disturbed because row j has zeros in those columns. The diagonal entries are preserved throughout.

Let t be the product of all the transvections used, in the order in which they are applied. Then tb has every off-diagonal entry equal to zero and has the same main diagonal entries as b . Hence tb is the required diagonal matrix. $\square$

Theorem 4.12. *G is generated by the set consisting of all invertible diagonal matrices and all transvections.*

Proof. Let H be the subgroup of G generated by all invertible diagonal matrices and all transvections. By Bruhat Decomposition Theorem, it is enough to prove that $B \leq H$ and $W \leq H$.

First let $b \in B$. By Lemma 4.11, there is a product t of transvections such that $tb = d$, where d is a diagonal matrix with the same main diagonal entries as b . Since b is invertible, d is an invertible diagonal matrix. Thus $t \in H$ and $d \in H$, so

$$b = t^{-1}d \in H.$$

Hence $B \leq H$.

It remains to show that $W \leq H$. By Proposition 4.5, the group W is generated by the permutation matrices corresponding to transpositions. Let $i \neq j$. Consider

$$X_{ji}(1)X_{ij}(-1)X_{ji}(1).$$

This product only acts non-trivially on the two-dimensional subspace spanned by $\mathbf{v}_i$ and $\mathbf{v}_j$. With respect to the ordered basis $(\mathbf{v}_i, \mathbf{v}_j)$, its action is represented by

$$\begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix} = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}.$$

Thus, equivalently by Lemma 4.6(vi), this product sends $\mathbf{v}_i$ to $\mathbf{v}_j$, sends $\mathbf{v}_j$ to $-\mathbf{v}_i$, and fixes every $\mathbf{v}_k$ with $k \notin \{i, j\}$. Now premultiply by the diagonal matrix whose (i, i) -entry is -1 and whose other diagonal entries are 1. The resulting matrix sends $\mathbf{v}_i$ to $\mathbf{v}_j$, sends $\mathbf{v}_j$ to $\mathbf{v}_i$, and fixes all other standard basis vectors, so it is exactly the permutation matrix for the transposition $(i j)$. In characteristic 2 this argument still works, since then $-1 = 1$.

Therefore each transposition matrix lies in H . Since every permutation is a product of transpositions, every permutation matrix is a product of such transposition matrices. Hence $W \leq H$.

Now Bruhat Decomposition Theorem gives

$$G = BWB \leq H.$$

The reverse inclusion $H \leq G$ is immediate from the definition of H , so $G = H$. □

Exercises

Exercise 4.1. Show that $\text{GL}(2, 2) \cong S_3$.

Solution. Let $E = \mathbb{Z}_2 \times \mathbb{Z}_2$. Then E is abelian, has exponent 2, and $|E| = 4 = 2^2$. Hence Proposition 4.1 gives

$$\text{Aut}(E) \cong \text{GL}(2, 2).$$

It remains to show that $\text{Aut}(E) \cong S_3$. Write

$$E = \{e, a, b, c\},$$

where e is the identity and a, b, c are the three non-identity elements. Since E has exponent 2, each of a, b, c has order 2. Also, the product of two distinct elements among a, b, c is the third one.

Every automorphism of E fixes e and permutes the set $X = \{a, b, c\}$. Therefore we get a homomorphism

$$\theta: \text{Aut}(E) \rightarrow S_X \cong S_3, \quad \theta(\varphi) = \varphi|_X.$$

If $\varphi, \psi \in \text{Aut}(E)$ and $x \in X$, then $\psi(x) \in X$, and so

$$\theta(\varphi \circ \psi)(x) = (\varphi \circ \psi)(x) = \varphi(\psi(x)) = \theta(\varphi)(\theta(\psi)(x)).$$

Hence $\theta(\varphi \circ \psi) = \theta(\varphi) \circ \theta(\psi)$, so θ is a homomorphism.

If $\theta(\varphi)$ is the identity permutation of X , then φ fixes a, b, c and also fixes e , so φ is the identity automorphism of E . Hence θ is injective.

By Proposition 4.2, we have

$$|\text{GL}(2, 2)| = (2^2 - 1)(2^2 - 2) = 6.$$

Therefore $|\text{Aut}(E)| = 6$. Also $|S_X| = 3! = 6$. Since θ is an injective homomorphism between finite groups of the same order, θ is an isomorphism. Thus

$$\text{Aut}(E) \cong S_X \cong S_3.$$

Combining this with $\text{Aut}(E) \cong \text{GL}(2, 2)$ gives $\text{GL}(2, 2) \cong S_3$. $\square$

Exercise 4.2. (cont.) Construct a monomorphism $\varphi: \text{GL}(1, 4) \rightarrow \text{GL}(2, 2)$ that corresponds to the inclusion of A_3 in S_3 . (Recall that the field $\mathbb{F}_4$ of 4 elements can be written as $\{0, 1, \alpha, \alpha^2\}$, where $\alpha + 1 = \alpha^2$ and $\lambda + \lambda = 0$ for all $\lambda \in \mathbb{F}_4$.) Show further that the extension of φ to a map from $\mathbb{F}_4$ to $\mathcal{M}_2(\mathbb{F}_2)$, where $\mathbb{F}_2 = \{0, 1\}$ is the field of 2 elements, is a monomorphism of rings.

Solution. We regard $\mathbb{F}_4$ as a 2-dimensional vector space over $\mathbb{F}_2$ with ordered basis

$$\mathcal{B} = (1, \alpha).$$

For each $\lambda \in \mathbb{F}_4$, let

$$m_\lambda: \mathbb{F}_4 \rightarrow \mathbb{F}_4, \quad m_\lambda(x) = \lambda x$$

be multiplication by λ . We define

$$\tilde{\varphi}: \mathbb{F}_4 \rightarrow \mathcal{M}_2(\mathbb{F}_2), \quad \tilde{\varphi}(\lambda) = [m_\lambda]_{\mathcal{B}}^{\mathcal{B}}.$$

Thus m_λ is the linear map, and $\tilde{\varphi}(\lambda)$ is the matrix of that map in the basis $\mathcal{B}$.

We compute this map on the elements of $\mathbb{F}_4$. Multiplication by 0 is the zero map, so

$$\tilde{\varphi}(0) = [m_0]_{\mathcal{B}}^{\mathcal{B}} = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}.$$

Multiplication by 1 is the identity map, so

$$\tilde{\varphi}(1) = [m_1]_{\mathcal{B}}^{\mathcal{B}} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}.$$

Next, since $\alpha \cdot 1 = \alpha$ and $\alpha \cdot \alpha = \alpha^2 = 1 + \alpha$, the first column of $\tilde{\varphi}(\alpha)$ is the coordinate column of α , and the second column is the coordinate column of $1 + \alpha$. Hence

$$\tilde{\varphi}(\alpha) = [m_\alpha]_{\mathcal{B}}^{\mathcal{B}} = \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}.$$

Finally, since $\alpha^2 \cdot 1 = \alpha^2 = 1 + \alpha$ and $\alpha^2 \cdot \alpha = \alpha^3 = 1$, we get

$$\tilde{\varphi}(\alpha^2) = [m_{\alpha^2}]_{\mathcal{B}}^{\mathcal{B}} = \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix}.$$

Restricting $\tilde{\varphi}$ to the non-zero elements $\mathbb{F}_4^\times = \{1, \alpha, \alpha^2\} = \text{GL}(1, 4)$ gives

$$\varphi: \text{GL}(1, 4) \rightarrow \text{GL}(2, 2), \quad \varphi(\lambda) = [m_\lambda]_{\mathcal{B}}^{\mathcal{B}}.$$

This is a group homomorphism because, for $\lambda, \mu \in \mathbb{F}_4^\times$,

$$m_{\lambda\mu} = m_\lambda \circ m_\mu,$$

and taking matrices in the basis $\mathcal{B}$ gives

$$\varphi(\lambda\mu) = [m_{\lambda\mu}]_{\mathcal{B}}^{\mathcal{B}} = [m_\lambda]_{\mathcal{B}}^{\mathcal{B}}[m_\mu]_{\mathcal{B}}^{\mathcal{B}} = \varphi(\lambda)\varphi(\mu).$$

It is injective because if $\varphi(\lambda) = \varphi(\mu)$, then $[m_\lambda]_{\mathcal{B}}^{\mathcal{B}} = [m_\mu]_{\mathcal{B}}^{\mathcal{B}}$, so $m_\lambda = m_\mu$. Applying both maps to 1 gives $\lambda = \mu$. Therefore φ is a monomorphism.

It remains to connect this with the inclusion $A_3 \leq S_3$. Since φ is injective and $\text{GL}(1, 4) = \mathbb{F}_4^\times$ has order 3, the subgroup $\text{im } \varphi$ has order 3. Under the isomorphism $\text{GL}(2, 2) \cong S_3$ from Exercise 4.1, this image corresponds to a subgroup of S_3 of order 3. The only such subgroup is

$$\{e, (1\ 2\ 3), (1\ 3\ 2)\} = A_3.$$

Thus φ corresponds to the usual inclusion $A_3 \leq S_3$. Finally, the map $\tilde{\varphi}: \mathbb{F}_4 \rightarrow \mathcal{M}_2(\mathbb{F}_2)$ is a ring homomorphism. Indeed, for $\lambda, \mu, x \in \mathbb{F}_4$,

$$m_{\lambda+\mu}(x) = (\lambda + \mu)x = \lambda x + \mu x = (m_\lambda + m_\mu)(x),$$

so $m_{\lambda+\mu} = m_\lambda + m_\mu$. Taking matrices in the basis $\mathcal{B}$ gives

$$\tilde{\varphi}(\lambda + \mu) = [m_{\lambda+\mu}]_{\mathcal{B}}^{\mathcal{B}} = [m_\lambda]_{\mathcal{B}}^{\mathcal{B}} + [m_\mu]_{\mathcal{B}}^{\mathcal{B}} = \tilde{\varphi}(\lambda) + \tilde{\varphi}(\mu).$$

Similarly,

$$m_{\lambda\mu} = m_\lambda \circ m_\mu,$$

so

$$\tilde{\varphi}(\lambda\mu) = [m_{\lambda\mu}]_{\mathcal{B}}^{\mathcal{B}} = [m_\lambda]_{\mathcal{B}}^{\mathcal{B}}[m_\mu]_{\mathcal{B}}^{\mathcal{B}} = \tilde{\varphi}(\lambda)\tilde{\varphi}(\mu),$$

and $\tilde{\varphi}(1) = I$. If $\tilde{\varphi}(\lambda) = \tilde{\varphi}(\mu)$, then $[m_\lambda]_{\mathcal{B}}^{\mathcal{B}} = [m_\mu]_{\mathcal{B}}^{\mathcal{B}}$, so $m_\lambda = m_\mu$. Applying both maps to 1 gives $\lambda = \mu$. Hence $\tilde{\varphi}$ is injective, and so it is a monomorphism of rings. $\square$

Exercise 4.3. (cont.) Construct an explicit monomorphism from $\text{GL}(n, 4)$ to $\text{GL}(2n, 2)$ for any $n \in \mathbb{N}$.

Solution. Let $\mathcal{B} = (1, \alpha)$ be the ordered $\mathbb{F}_2$ -basis of $\mathbb{F}_4$ used in Exercise 4.2. Write

$$\rho(\lambda) = [m_\lambda]_{\mathcal{B}}^{\mathcal{B}} \in \mathcal{M}_2(\mathbb{F}_2)$$

for the matrix of multiplication by λ on $\mathbb{F}_4$. By Exercise 4.2, the map $\rho: \mathbb{F}_4 \rightarrow \mathcal{M}_2(\mathbb{F}_2)$ is an injective ring homomorphism.

Now let $M = (\lambda_{ij}) \in \text{GL}(n, 4)$. Define $\Phi(M)$ to be the $2n \times 2n$ block matrix over $\mathbb{F}_2$ whose (i, j) -block is $\rho(\lambda_{ij})$:

$$\Phi(M) = \begin{pmatrix} \rho(\lambda_{11}) & \rho(\lambda_{12}) & \cdots & \rho(\lambda_{1n}) \\ \rho(\lambda_{21}) & \rho(\lambda_{22}) & \cdots & \rho(\lambda_{2n}) \\ \vdots & \vdots & \ddots & \vdots \\ \rho(\lambda_{n1}) & \rho(\lambda_{n2}) & \cdots & \rho(\lambda_{nn}) \end{pmatrix}.$$

This is the same as replacing each entry of M by its 2×2 multiplication matrix $[m_{\lambda_{ij}}]_{\mathcal{B}}^{\mathcal{B}}$.

We claim that

$$\Phi: \text{GL}(n, 4) \rightarrow \text{GL}(2n, 2)$$

is a monomorphism. First, if $M = (\lambda_{ij})$ and $N = (\mu_{ij})$ are in $\text{GL}(n, 4)$, then the (i, j) -block of $\Phi(M)\Phi(N)$ is

$$\sum_{k=1}^n \rho(\lambda_{ik})\rho(\mu_{kj}).$$

Since ρ is a ring homomorphism, this equals

$$\sum_{k=1}^n \rho(\lambda_{ik}\mu_{kj}) = \rho\left(\sum_{k=1}^n \lambda_{ik}\mu_{kj}\right),$$

which is exactly the (i, j) -block of $\Phi(MN)$. Hence $\Phi(MN) = \Phi(M)\Phi(N)$.

If $M \in \text{GL}(n, 4)$, let M^{-1} be its inverse over $\mathbb{F}_4$. Then

$$\Phi(M)\Phi(M^{-1}) = \Phi(I_n) = I_{2n},$$

so $\Phi(M)$ is invertible. Hence $\Phi(M) \in \text{GL}(2n, 2)$.

Finally, suppose $\Phi(M) = \Phi(N)$. Then every corresponding 2×2 block is equal, so $\rho(\lambda_{ij}) = \rho(\mu_{ij})$ for all i, j . Since ρ is injective, $\lambda_{ij} = \mu_{ij}$ for all i, j , and therefore $M = N$. Thus Φ is injective.

Therefore Φ is an explicit monomorphism from $\text{GL}(n, 4)$ to $\text{GL}(2n, 2)$. $\square$

Exercise 4.4. (cont.) More generally, show for any $n \in \mathbb{N}$ and any prime power q that $\text{GL}(2n, q)$ has a subgroup that is isomorphic with $\text{GL}(n, q^2)$. Will $\text{GL}(n, q^m)$ always have a subgroup isomorphic with $\text{GL}(mn, q)$ for any $m, n \in \mathbb{N}$ and any prime power q ?

Solution. Let $\mathbb{F}_{q^2}$ be the field with q^2 elements. Since $\mathbb{F}_{q^2}$ is a 2-dimensional vector space over $\mathbb{F}_q$, choose an ordered $\mathbb{F}_q$ -basis

$$\mathcal{B} = (\beta_1, \beta_2)$$

of $\mathbb{F}_{q^2}$. For each $\lambda \in \mathbb{F}_{q^2}$, let

$$m_\lambda: \mathbb{F}_{q^2} \rightarrow \mathbb{F}_{q^2}, \quad m_\lambda(x) = \lambda x,$$

and define

$$\rho(\lambda) = [m_\lambda]_{\mathcal{B}}^{\mathcal{B}} \in \mathcal{M}_2(\mathbb{F}_q).$$

This is an injective ring homomorphism. Indeed, for $\lambda, \mu \in \mathbb{F}_{q^2}$ and $x \in \mathbb{F}_{q^2}$, we have

$$m_{\lambda+\mu}(x) = (\lambda + \mu)x = \lambda x + \mu x = (m_\lambda + m_\mu)(x),$$

so $\rho(\lambda + \mu) = \rho(\lambda) + \rho(\mu)$. Also

$$m_{\lambda\mu}(x) = (\lambda\mu)x = \lambda(\mu x) = (m_\lambda \circ m_\mu)(x),$$

so $\rho(\lambda\mu) = \rho(\lambda)\rho(\mu)$. Finally, if $\rho(\lambda) = \rho(\mu)$, then $m_\lambda = m_\mu$, so evaluating both maps at 1 gives $\lambda = \mu$. Thus ρ is injective.

Now let $M = (\lambda_{ij}) \in \text{GL}(n, q^2)$. Define $\Phi(M)$ to be the $2n \times 2n$ block matrix over $\mathbb{F}_q$ whose (i, j) -block is $\rho(\lambda_{ij})$:

$$\Phi(M) = \begin{pmatrix} \rho(\lambda_{11}) & \rho(\lambda_{12}) & \cdots & \rho(\lambda_{1n}) \\ \rho(\lambda_{21}) & \rho(\lambda_{22}) & \cdots & \rho(\lambda_{2n}) \\ \vdots & \vdots & \ddots & \vdots \\ \rho(\lambda_{n1}) & \rho(\lambda_{n2}) & \cdots & \rho(\lambda_{nn}) \end{pmatrix}.$$

Let $N = (\mu_{ij}) \in \text{GL}(n, q^2)$. The (i, j) -block of $\Phi(M)\Phi(N)$ is

$$\sum_{k=1}^n \rho(\lambda_{ik})\rho(\mu_{kj}) = \sum_{k=1}^n \rho(\lambda_{ik}\mu_{kj}) = \rho\left(\sum_{k=1}^n \lambda_{ik}\mu_{kj}\right),$$

which is exactly the (i, j) -block of $\Phi(MN)$. Hence

$$\Phi(MN) = \Phi(M)\Phi(N)$$

for $M, N \in \text{GL}(n, q^2)$. If $M \in \text{GL}(n, q^2)$, then

$$\Phi(M)\Phi(M^{-1}) = \Phi(I_n) = I_{2n},$$

so $\Phi(M) \in \text{GL}(2n, q)$. Also, if $\Phi(M) = \Phi(N)$, then $\rho(\lambda_{ij}) = \rho(\mu_{ij})$ for all i, j , and since ρ is injective, $\lambda_{ij} = \mu_{ij}$ for all i, j . Hence $M = N$. Therefore Φ is an injective homomorphism

$$\Phi: \text{GL}(n, q^2) \rightarrow \text{GL}(2n, q).$$

Thus $\text{GL}(2n, q)$ has a subgroup isomorphic with $\text{GL}(n, q^2)$.

The proposed reverse-looking statement is not always true. Take $n = 1$, $m = 2$, and $q = 2$. Then it would say that $\text{GL}(1, 4)$ has a subgroup isomorphic with $\text{GL}(2, 2)$. But

$$|\text{GL}(1, 4)| = |\mathbb{F}_4^\times| = 3,$$

whereas by Proposition 4.2,

$$|\text{GL}(2, 2)| = (2^2 - 1)(2^2 - 2) = 6.$$

A group of order 3 cannot have a subgroup of order 6. Therefore $\text{GL}(n, q^m)$ does not always have a subgroup isomorphic with $\text{GL}(mn, q)$. $\square$

Exercise 4.5. Show that $\text{GL}(4, 2) \cong A_8$.

Solution. Let $G = \text{GL}(4, 2)$. We first record the plan. By Proposition 4.2,

$$|G| = (2^4 - 1)(2^4 - 2)(2^4 - 2^2)(2^4 - 2^3) = 15 \cdot 14 \cdot 12 \cdot 8 = 20160.$$

Also

$$|A_8| = \frac{8!}{2} = 20160.$$

Thus it is enough to construct an embedding

$$G \hookrightarrow S_8,$$

because the image will then have order $8!/2$, hence will be an index 2 subgroup of S_8 . The unique index 2 subgroup of S_8 is A_8 .

The hard part is finding the eight objects on which G should act. We shall find them by first constructing a subgroup $H \leq G$ with $H \cong A_7$; then the eight objects will be the left cosets of H in G .

Step 1: Constructing fifteen Fano planes.

Let $\Omega = \{1, \dots, 7\}$. A *Fano plane structure* on Ω is a set of seven 3-element subsets of Ω , called lines, such that any two points of Ω lie in exactly one line. For example, take

$$\mathcal{P}_0 = \{\{1, 2, 3\}, \{1, 4, 5\}, \{1, 6, 7\}, \{2, 4, 6\}, \{2, 5, 7\}, \{3, 4, 7\}, \{3, 5, 6\}\}.$$

We may identify the points $1, \dots, 7$ with the seven non-zero vectors of $\mathbb{F}_2^3$ by

point	1	2	3	4	5	6	7
vector	e_1	e_2	$e_1 + e_2$	e_3	$e_1 + e_3$	$e_2 + e_3$	$e_1 + e_2 + e_3$.

With this identification, the lines of $\mathcal{P}_0$ are exactly the triples

$$\{u, v, u + v\}$$

where u and v are distinct non-zero vectors. Therefore every invertible linear map of $\mathbb{F}_2^3$ permutes the seven points and preserves the lines, giving

$$\mathrm{GL}(3, 2) \leq \mathrm{Aut}(\mathcal{P}_0).$$

Conversely, every automorphism of $\mathcal{P}_0$ is determined by the images of three non-collinear points, for instance $1, 2, 4$. These images are again non-collinear, hence form a basis of $\mathbb{F}_2^3$, and so determine a unique invertible linear map. Thus

$$\mathrm{Aut}(\mathcal{P}_0) \cong \mathrm{GL}(3, 2).$$

By Proposition 4.2,

$$|\mathrm{Aut}(\mathcal{P}_0)| = |\mathrm{GL}(3, 2)| = (2^3 - 1)(2^3 - 2)(2^3 - 2^2) = 168.$$

Moreover $\mathrm{Aut}(\mathcal{P}_0) \leq A_7$. To see this, note that by Theorem 4.12, $\mathrm{GL}(3, 2)$ is generated by transvections, since the only invertible diagonal matrix over $\mathbb{F}_2$ is the identity. A transvection of $\mathbb{F}_2^3$ fixes three non-zero vectors and interchanges the remaining four non-zero vectors in two pairs. Hence it is an even permutation of the seven non-zero vectors. So every element of $\mathrm{Aut}(\mathcal{P}_0)$ is an even permutation.

Let

$$\mathcal{P} = A_7 \mathcal{P}_0$$

be the orbit of $\mathcal{P}_0$ under the natural action of A_7 on Ω . Since the stabilizer of $\mathcal{P}_0$ in A_7 is $\mathrm{Aut}(\mathcal{P}_0)$, we have

$$|\mathcal{P}| = \frac{|A_7|}{|\mathrm{Aut}(\mathcal{P}_0)|} = \frac{7!/2}{168} = 15.$$

So $\mathcal{P}$ is a set of fifteen Fano plane structures on the same seven-point set Ω .

We shall need one elementary fact about these fifteen Fano planes: any two distinct elements of $\mathcal{P}$ have exactly one common line. It is enough to check this with one of the two planes equal to $\mathcal{P}_0$. Let $\ell = \{1, 2, 3\}$ and let $\tau = (1\ 2\ 3)$. Then $\tau\mathcal{P}_0$ and $\tau^2\mathcal{P}_0$ are the two Fano planes

$$\begin{aligned} \tau\mathcal{P}_0 &= \{\{1, 2, 3\}, \{2, 4, 5\}, \{2, 6, 7\}, \{3, 4, 6\}, \{3, 5, 7\}, \{1, 4, 7\}, \{1, 5, 6\}\}, \\ \tau^2\mathcal{P}_0 &= \{\{1, 2, 3\}, \{3, 4, 5\}, \{3, 6, 7\}, \{1, 4, 6\}, \{1, 5, 7\}, \{2, 4, 7\}, \{2, 5, 6\}\}. \end{aligned}$$

Each has exactly the line ℓ in common with $\mathcal{P}_0$. Since $\mathrm{Aut}(\mathcal{P}_0)$ is transitive on the seven lines of $\mathcal{P}_0$, the same statement holds for every line of $\mathcal{P}_0$. Thus the two rotations of each of the seven lines give fourteen distinct elements of $\mathcal{P}$ different from $\mathcal{P}_0$. Since $|\mathcal{P}| = 15$, these are all the other elements of $\mathcal{P}$. Therefore

every element of $\mathcal{P} - \{\mathcal{P}_0\}$ has exactly one line in common with $\mathcal{P}_0$. Applying an element of A_7 gives the same conclusion for every pair of distinct elements of $\mathcal{P}$.

It follows also that if $\mathcal{Q} \in \mathcal{P}$ and ℓ is a line of $\mathcal{Q}$, then exactly three elements of $\mathcal{P}$ contain ℓ : namely $\mathcal{Q}$ and the two rotations of $\mathcal{Q}$ around ℓ .

Step 2: Turning the fifteen planes into a vector space.

We now put a vector space structure over $\mathbb{F}_2$ on

$$V = \mathcal{P} \cup \{0\}.$$

The zero element is the new symbol 0. For $\mathcal{Q} \in \mathcal{P}$, define

$$\mathcal{Q} + 0 = 0 + \mathcal{Q} = \mathcal{Q}, \quad \mathcal{Q} + \mathcal{Q} = 0.$$

If $\mathcal{Q}_1, \mathcal{Q}_2 \in \mathcal{P}$ are distinct, let ℓ be their unique common line. There is a unique third element $\mathcal{Q}_3 \in \mathcal{P}$ which contains ℓ . Define

$$\mathcal{Q}_1 + \mathcal{Q}_2 = \mathcal{Q}_3.$$

This addition is commutative, has identity 0, and every element has order 2. We check associativity. If one of the three entries is 0, or if two of the three entries are equal, associativity follows directly from the definition. If $\mathcal{Q}_1, \mathcal{Q}_2, \mathcal{Q}_3$ are distinct and have one common line, then

$$\mathcal{Q}_1 + \mathcal{Q}_2 = \mathcal{Q}_3, \quad \mathcal{Q}_2 + \mathcal{Q}_3 = \mathcal{Q}_1,$$

and hence

$$(\mathcal{Q}_1 + \mathcal{Q}_2) + \mathcal{Q}_3 = \mathcal{Q}_3 + \mathcal{Q}_3 = 0 = \mathcal{Q}_1 + \mathcal{Q}_1 = \mathcal{Q}_1 + (\mathcal{Q}_2 + \mathcal{Q}_3).$$

It remains to check the case where $\mathcal{Q}_1, \mathcal{Q}_2, \mathcal{Q}_3$ are distinct and do not all share one common line. There are $15 \cdot 14 \cdot 12 = 2520$ ordered triples of this type: after choosing $\mathcal{Q}_1$ and $\mathcal{Q}_2$, the third entry can be any element of $\mathcal{P}$ except the three planes containing the common line of $\mathcal{Q}_1$ and $\mathcal{Q}_2$. We show that A_7 is transitive on these triples by looking at one example. Take $\mathcal{P}_0$ as above and set

$$\begin{aligned} \mathcal{Q}_1 &= \{\{1, 2, 3\}, \{1, 4, 6\}, \{1, 5, 7\}, \{2, 4, 7\}, \{2, 5, 6\}, \{3, 4, 5\}, \{3, 6, 7\}\}, \\ \mathcal{Q}_2 &= \{\{1, 2, 4\}, \{1, 3, 5\}, \{1, 6, 7\}, \{2, 3, 7\}, \{2, 5, 6\}, \{3, 4, 6\}, \{4, 5, 7\}\}. \end{aligned}$$

The three pairwise common lines are

$$\mathcal{P}_0 \cap \mathcal{Q}_1 = \{\{1, 2, 3\}\}, \quad \mathcal{P}_0 \cap \mathcal{Q}_2 = \{\{1, 6, 7\}\}, \quad \mathcal{Q}_1 \cap \mathcal{Q}_2 = \{\{2, 5, 6\}\}.$$

Any element of A_7 which fixes the ordered triple $(\mathcal{P}_0, \mathcal{Q}_1, \mathcal{Q}_2)$ must fix these three lines setwise. It therefore fixes their pairwise intersections 1, 2, and 6; then it fixes the remaining point on each of the three lines, namely 3, 7, and 5; and finally it fixes the last point 4. Thus the stabilizer of this ordered triple is trivial. Its orbit has size $|A_7| = 2520$, which is the total number of triples in this case. Hence every such triple is in this one orbit, so it is enough to check associativity for the displayed example. Direct use of the addition rule gives

$$\begin{aligned} \mathcal{P}_0 + \mathcal{Q}_1 &= \{\{1, 2, 3\}, \{1, 4, 7\}, \{1, 5, 6\}, \{2, 4, 5\}, \{2, 6, 7\}, \{3, 4, 6\}, \{3, 5, 7\}\}, \\ \mathcal{Q}_1 + \mathcal{Q}_2 &= \{\{1, 2, 7\}, \{1, 3, 6\}, \{1, 4, 5\}, \{2, 3, 4\}, \{2, 5, 6\}, \{3, 5, 7\}, \{4, 6, 7\}\}, \end{aligned}$$

and both $(\mathcal{P}_0 + \mathcal{Q}_1) + \mathcal{Q}_2$ and $\mathcal{P}_0 + (\mathcal{Q}_1 + \mathcal{Q}_2)$ are equal to

$$\{\{1, 2, 6\}, \{1, 3, 7\}, \{1, 4, 5\}, \{2, 3, 5\}, \{2, 4, 7\}, \{3, 4, 6\}, \{5, 6, 7\}\}.$$

Thus V is an elementary abelian group of order 16, hence a 4-dimensional vector space over $\mathbb{F}_2$.

Step 3: Getting A_7 inside $\text{GL}(4, 2)$.

The action of A_7 on $\mathcal{P}$ preserves the addition on V , since the definition of addition only uses common lines. Therefore we obtain a homomorphism

$$A_7 \rightarrow \text{GL}(V).$$

We now show this homomorphism is injective. We use the elementary fact that A_7 is simple. Here is a proof. Let $1 \neq N \trianglelefteq A_7$, and choose a non-identity element of N moving as few letters as possible. The possible non-identity cycle types in A_7 are represented by

$$(1\ 2\ 3), \quad (1\ 2)(3\ 4), \quad (1\ 2\ 3\ 4\ 5), \quad (1\ 2\ 3\ 4\ 5\ 6\ 7), \quad (1\ 2\ 3)(4\ 5\ 6), \quad (1\ 2\ 3\ 4)(5\ 6), \quad (1\ 2\ 3)(4\ 5)(6\ 7).$$

If the chosen element is not a 3-cycle, then after conjugating in A_7 and replacing by its inverse if necessary, one of the following commutator calculations applies:

$$\begin{aligned} [(1\ 2)(3\ 4), (1\ 2\ 5)] &= (1\ 2\ 5), \\ [(1\ 2\ 3\ 4\ 5), (1\ 2\ 3)] &= (1\ 4\ 2), \\ [(1\ 2\ 3\ 4\ 5\ 6\ 7), (1\ 2\ 3)] &= (1\ 4\ 2), \\ [(1\ 2\ 3)(4\ 5\ 6), (1\ 2\ 4)] &= (1\ 4\ 3\ 5\ 2), \\ [(1\ 2\ 3\ 4)(5\ 6), (1\ 2\ 3)] &= (1\ 4\ 2), \\ [(1\ 2\ 3)(4\ 5)(6\ 7), (1\ 2\ 4)] &= (1\ 4\ 3\ 5\ 2). \end{aligned}$$

Each commutator lies in N because N is normal in A_7 , and each displayed commutator is non-trivial and moves fewer letters than the original element. This contradicts the minimal choice unless N contains a 3-cycle. Since all 3-cycles are conjugate in A_7 , N contains every 3-cycle. Finally, 3-cycles generate A_7 : every even permutation is a product of pairs of transpositions, and a product of two transpositions is either a 3-cycle or, when the transpositions are disjoint,

$$(a\ b)(c\ d) = (a\ c\ b)(a\ c\ d).$$

It follows that $N = A_7$. Thus A_7 is simple.

The kernel of the action of A_7 on V is a normal subgroup of A_7 . The action is not trivial because the orbit $\mathcal{P} = A_7\mathcal{P}_0$ has 15 elements. Hence the kernel is not all of A_7 , and by simplicity it must be trivial. Thus

$$A_7 \hookrightarrow \mathrm{GL}(V).$$

Since $\dim_{\mathbb{F}_2} V = 4$, we have $\mathrm{GL}(V) \cong \mathrm{GL}(4, 2)$. Therefore $G = \mathrm{GL}(4, 2)$ contains a subgroup

$$H \cong A_7.$$

Step 4: Acting on the eight cosets of H .

Since

$$|H| = |A_7| = \frac{7!}{2} = 2520,$$

we have

$$|G : H| = \frac{20160}{2520} = 8.$$

Thus G acts by left multiplication on the set G/H of eight left cosets of H . This gives a homomorphism

$$\varphi : G \rightarrow S_8.$$

We claim that $\ker \varphi = 1$. The kernel of the action is contained in H , because the identity coset H is fixed precisely by elements of H . Also $\ker \varphi \trianglelefteq G$, so $\ker \varphi \trianglelefteq H$. Since $H \cong A_7$ is simple, $\ker \varphi$ is either 1 or H .

It remains to rule out $\ker \varphi = H$. If $\ker \varphi = H$, then $H \trianglelefteq G$ and G/H has order 8. But G is generated by transvections: over $\mathbb{F}_2$ every invertible diagonal matrix is the identity, so Theorem 4.12 says that G is generated by transvections. Also, since $4 \geq 3$, every transvection is a commutator. Indeed, for distinct i, j , choose k distinct from both i and j , and Lemma 4.6(iv) gives

$$[X_{ik}(1), X_{kj}(1)] = X_{ij}(1).$$

Hence $G = G'$, so every quotient of G is perfect. This is impossible for a quotient of order 8. Indeed, if a group Q has order 8, then either Q has an element of order 4 or 8, in which case it has a cyclic subgroup of index 2 and therefore a non-trivial abelian quotient, or every non-identity element of Q has order 2, in which case Q is abelian because $(ab)^2 = 1$ implies $ab = ba$. Thus no group of order 8 is perfect. Therefore $\ker \varphi \neq H$, and so $\ker \varphi = 1$.

Therefore φ embeds G into S_8 . Its image has order $20160 = 8!/2$, so it is an index 2 subgroup of S_8 . Finally, the unique index 2 subgroup of S_8 is A_8 . Indeed, an index 2 subgroup is normal, so it is the kernel of a non-trivial homomorphism $S_8 \rightarrow C_2$; since transpositions generate S_8 and are all conjugate, this kernel consists exactly of the even permutations. Hence

$$\varphi(G) = A_8.$$

Thus

$$\mathrm{GL}(4, 2) \cong A_8.$$

□

Exercise 4.6. Let β be a non-trivial automorphism of the field F . Use β to construct an outer automorphism of $\text{GL}(n, F)$. Do all outer automorphisms of $\text{GL}(n, F)$ arise in this way?

Solution. For $A = (a_{ij}) \in \text{GL}(n, F)$, define

$$\Phi_\beta(A) = (\beta(a_{ij})).$$

That is, Φ_β applies β to every entry of the matrix. Since β preserves addition and multiplication, the (i, j) -entry of $\Phi_\beta(AB)$ is

$$\beta \left(\sum_{k=1}^n a_{ik} b_{kj} \right) = \sum_{k=1}^n \beta(a_{ik}) \beta(b_{kj}),$$

which is exactly the (i, j) -entry of $\Phi_\beta(A)\Phi_\beta(B)$. Hence

$$\Phi_\beta(AB) = \Phi_\beta(A)\Phi_\beta(B).$$

If $A \in \text{GL}(n, F)$, then A^{-1} exists and

$$\Phi_\beta(A)\Phi_\beta(A^{-1}) = \Phi_\beta(AA^{-1}) = I,$$

so $\Phi_\beta(A)$ is again invertible. Thus Φ_β is a homomorphism $\text{GL}(n, F) \rightarrow \text{GL}(n, F)$. Its inverse is obtained by applying β^{-1} entry by entry, and hence Φ_β is an automorphism of $\text{GL}(n, F)$.

We now show that it is outer. Suppose, for contradiction, that Φ_β is inner. Then there exists $P \in \text{GL}(n, F)$ such that

$$\Phi_\beta(A) = PAP^{-1}$$

for every $A \in \text{GL}(n, F)$. Since β is non-trivial and fixes 0 and 1, choose $\lambda \in F^\times$ such that $\beta(\lambda) \neq \lambda$. Taking $A = \lambda I$, we get on one hand

$$\Phi_\beta(\lambda I) = \beta(\lambda)I.$$

On the other hand, if Φ_β were conjugation by P , then

$$\Phi_\beta(\lambda I) = P(\lambda I)P^{-1} = \lambda PP^{-1} = \lambda I.$$

Thus $\beta(\lambda)I = \lambda I$, forcing $\beta(\lambda) = \lambda$, a contradiction. Therefore Φ_β is not inner, and hence it is outer. Not all outer automorphisms arise in this way. For $n \geq 2$, define

$$\tau(A) = (A^{-1})^T.$$

Then

$$\tau(AB) = ((AB)^{-1})^T = (B^{-1}A^{-1})^T = (A^{-1})^T(B^{-1})^T = \tau(A)\tau(B),$$

so τ is an automorphism of $\text{GL}(n, F)$. It is outer by the same scalar-matrix test: choose $\lambda \in F^\times$ with $\lambda^{-1} \neq \lambda$; such a λ exists under the present hypothesis, since otherwise every non-zero element of F would satisfy $x^2 = 1$, forcing $|F^\times| \leq 2$ and leaving no non-trivial field automorphism. Then $\tau(\lambda I) = \lambda^{-1}I$, while every inner automorphism fixes λI . Finally, τ is not obtained from a field automorphism, because a field automorphism γ sends

$$X_{ij}(\alpha) \mapsto X_{ij}(\gamma(\alpha)),$$

keeping the same matrix position (i, j) , whereas

$$\tau(X_{ij}(\alpha)) = (X_{ij}(\alpha)^{-1})^T = X_{ji}(-\alpha),$$

which swaps the position from (i, j) to (j, i) . Therefore field automorphisms give outer automorphisms of $\text{GL}(n, F)$, but they do not account for all of them. $\square$

Further Exercises

Exercise 4.7. A matrix is said to be *monomial* if each row and column has exactly one non-zero entry. Let N be the subset of $G = \text{GL}(n, F)$ consisting of all monomial matrices. Show that $N \leq G$, that $T = B \cap N$ is the subgroup of G consisting of all diagonal matrices, that $N = N_G(T)$, and that $N = T \rtimes W$.

Solution. Let $\mathcal{D}$ denote the subgroup of G consisting of all invertible diagonal matrices. We first show that

$$N = \mathcal{D}W.$$

Let $m = (m_{ij}) \in N$. In each row i of m , there is exactly one non-zero entry. Let that entry occur in column $\sigma(i)$. Since each column also has exactly one non-zero entry, σ is a permutation of $\{1, \dots, n\}$. Let w be the permutation matrix whose non-zero entries occur in the same positions as the non-zero entries of m , and let

$$D = \text{diag}(m_{1,\sigma(1)}, \dots, m_{n,\sigma(n)}).$$

Then $D \in \mathcal{D}$, because all these entries are non-zero, and left multiplication by D scales row i of w by $m_{i,\sigma(i)}$. Hence $m = Dw$, so $m \in \mathcal{D}W$. Thus $N \subseteq \mathcal{D}W$. Conversely, if $D \in \mathcal{D}$ and $w \in W$, then Dw is obtained from the permutation matrix w by scaling each row by a non-zero scalar. Therefore every row and column still has exactly one non-zero entry, so $Dw \in N$. Hence $\mathcal{D}W \subseteq N$, and therefore $N = \mathcal{D}W$. We now prove that $N \leq G$. The identity matrix is monomial, so N is non-empty. Let $m, l \in N$. Write

$$m = Dw, \quad l = Eu,$$

where $D, E \in \mathcal{D}$ and $w, u \in W$. Then

$$ml = DwEu = D(wEw^{-1})(wu).$$

The matrix wEw^{-1} is diagonal, because conjugating a diagonal matrix by a permutation matrix only permutes the diagonal entries. Also $wu \in W$. Hence ml is diagonal times permutation, so $ml \in N$. Similarly,

$$m^{-1} = (Dw)^{-1} = w^{-1}D^{-1} = (w^{-1}D^{-1}w)w^{-1}.$$

Again $w^{-1}D^{-1}w$ is diagonal and $w^{-1} \in W$, so $m^{-1} \in N$. Therefore $N \leq G$.

Next we identify $B \cap N$. Let $x \in B \cap N$. Since $x \in N$, the non-zero entries of x have the form of a permutation: in row i , the unique non-zero entry occurs in column $\sigma(i)$, where σ is a permutation of $\{1, \dots, n\}$. Since $x \in B$, all entries below the diagonal are zero, so this non-zero entry cannot lie below the diagonal. Hence

$$\sigma(i) \geq i$$

for every i . But σ is a permutation, so

$$\sum_{i=1}^n \sigma(i) = \sum_{i=1}^n i.$$

Since each $\sigma(i) \geq i$, equality of the sums forces $\sigma(i) = i$ for every i . Thus the only non-zero entries of x are diagonal entries, so x is diagonal. Therefore $B \cap N \subseteq \mathcal{D}$. Conversely, every invertible diagonal matrix is both upper triangular and monomial, so $\mathcal{D} \subseteq B \cap N$. Hence

$$T = B \cap N = \mathcal{D}.$$

In particular, T is the subgroup of G consisting of all invertible diagonal matrices.

We now prove that $N = N_G(T)$. For this assertion we assume $F \neq \mathbb{F}_2$; otherwise, if $n > 1$, then $T = \{I\}$ and $N_G(T) = G$, so the displayed equality is false. First let $m \in N$. Write $m = Dw$ with $D \in T$ and $w \in W$. If $t \in T$, then

$$mtm^{-1} = Dwtw^{-1}D^{-1}.$$

The matrix wtw^{-1} is diagonal, hence lies in T . Since diagonal matrices commute, we get

$$Dwtw^{-1}D^{-1} = wtw^{-1} \in T.$$

Thus $mTm^{-1} \subseteq T$. Applying the same argument to $m^{-1} \in N$ gives $T \subseteq mTm^{-1}$, and hence $mTm^{-1} = T$. Therefore $m \in N_G(T)$, so $N \subseteq N_G(T)$.

Conversely, let $g \in N_G(T)$. We prove that g is monomial. Let

$$V = F^n$$

with standard basis $\mathbf{v}_1, \dots, \mathbf{v}_n$. Every matrix $A \in G$ acts on V by left multiplication:

$$L_A: V \rightarrow V, \quad L_A(x) = Ax.$$

In particular, $g\mathbf{v}_i = L_g(\mathbf{v}_i)$ is exactly the i th column of the matrix g . Since $F \neq \mathbb{F}_2$, choose $\lambda \in F^\times$ with $\lambda \neq 1$. For each i , define

$$t_i = \text{diag}(1, \dots, 1, \lambda, 1, \dots, 1),$$

where λ is in the i th diagonal position. Then $t_i \in T$. As a linear transformation, L_{t_i} satisfies

$$L_{t_i}(\mathbf{v}_i) = \lambda \mathbf{v}_i,$$

and, for $k \neq i$,

$$L_{t_i}(\mathbf{v}_k) = \mathbf{v}_k.$$

Thus the λ -eigenspace of L_{t_i} is exactly

$$E_\lambda(L_{t_i}) = F\mathbf{v}_i = \{\alpha \mathbf{v}_i \mid \alpha \in F\}.$$

Indeed, if $x = \sum_{k=1}^n a_k \mathbf{v}_k$ and $t_i x = \lambda x$, then for every $k \neq i$ we have $a_k = \lambda a_k$, so $(\lambda - 1)a_k = 0$, and hence $a_k = 0$. Thus only the $\mathbf{v}_i$ -coordinate can remain.

Because $g \in N_G(T)$ and $t_i \in T$, the conjugate

$$s_i = gt_i g^{-1}$$

also lies in T . Hence s_i is diagonal. Now compute the action of s_i on $g\mathbf{v}_i$:

$$s_i(g\mathbf{v}_i) = (gt_i g^{-1})(g\mathbf{v}_i) = gt_i \mathbf{v}_i = g(\lambda \mathbf{v}_i) = \lambda g\mathbf{v}_i.$$

So $g\mathbf{v}_i$ is a λ -eigenvector of the diagonal matrix s_i . More precisely, conjugation transports eigenspaces:

$$E_\lambda(L_{s_i}) = g(E_\lambda(L_{t_i})) = g(F\mathbf{v}_i).$$

Since s_i is diagonal, its λ -eigenspace is spanned by those standard basis vectors whose diagonal entries in s_i are equal to λ . Since s_i is conjugate to t_i , this eigenspace has the same dimension as $E_\lambda(L_{t_i})$, namely dimension 1. Therefore there is some index j such that

$$E_\lambda(L_{s_i}) = F\mathbf{v}_j.$$

In particular,

$$g\mathbf{v}_i \in F\mathbf{v}_j.$$

Thus there is some $\alpha_i \in F^\times$ such that

$$g\mathbf{v}_i = \alpha_i \mathbf{v}_j.$$

But $g\mathbf{v}_i$ is the i th column of g , so the i th column of g has exactly one non-zero entry. Since this holds for every i , every column of g has exactly one non-zero entry. Finally, g is invertible, so its columns are linearly independent. Hence two different columns cannot be non-zero scalar multiples of the same standard basis vector. Therefore the non-zero entries occur in different rows, and every row also has exactly one non-zero entry. Thus g is monomial, so $g \in N$. Hence $N_G(T) \subseteq N$. Combining both inclusions gives

$$N = N_G(T).$$

It remains to prove the semidirect product statement. We already proved $N = TW$, since $T = \mathcal{D}$. Also $T \trianglelefteq N$: if $m = Dw \in N$ and $t \in T$, then the same computation as above gives

$$mtm^{-1} = twt^{-1} \in T.$$

Finally,

$$T \cap W = \{I\}.$$

Indeed, if $x \in T \cap W$, then x is diagonal, say

$$x = \text{diag}(\mu_1, \dots, \mu_n),$$

and x is also a permutation matrix. Since x is diagonal, the only possible non-zero entry in row i is the diagonal entry μ_i . Since x is a permutation matrix, the non-zero entry in each row is equal to 1. Hence $\mu_i = 1$ for every i , and so $x = I$. The reverse inclusion $I \in T \cap W$ is immediate. Therefore $T \cap W = \{I\}$. Since $N = TW$, $T \trianglelefteq N$, and $T \cap W = \{I\}$, we conclude that

$$N = T \rtimes W.$$

□

Let G be a group. Suppose that G has subgroups B and N of G satisfying the following conditions:

- G is generated by B and N .
- $T = B \cap N$ is a normal subgroup of N .
- $W = N/T$ is generated by a finite set S of involutions (elements of order 2). In addition, if for each $w \in W$ we choose some $\dot{w} \in N$ such that $\dot{w}T = w$, then we must have
 - $\dot{s}B\dot{w} \subseteq B\dot{w}B \cup B\dot{s}wB$ for any $s \in S$ and $w \in W$,
 - $\dot{s}B\dot{s} \not\subseteq B$ for any $s \in S$.

In this case we say that B and N form a *BN-pair* of G , or that (G, B, N, S) is a *Tits system* (after Jacques Tits). We call B the *Borel subgroup* of G , and $W = N/B \cap N$ the *Weyl group* associated with the Tits system. The *rank* of the Tits system is defined to be $|S|$.

Exercise 4.8. (cont.) Let $G = \text{GL}(n, F)$, let B be the standard Borel subgroup of G , let N be the subgroup of G consisting of all monomial matrices, and let $T = B \cap N$. By Exercise 4.7 above, we know that we can regard $W = N/T$ as being imbedded in G as the group of permutation matrices. (By making this identification, we can replace $\dot{w}$ by w in the above formulas.) Let S be the subset of W consisting of those permutation matrices that are obtained from the identity matrix by switching two adjacent columns. (In other words, if we identify W with S_n as in Proposition 4.5, then S corresponds to the set $\{(1\ 2), (2\ 3), \dots, (n-1\ n)\}$.) Show that (G, B, N, S) is a Tits system of rank $n-1$.

Solution. We check the defining conditions one at a time. To keep the quotient notation separate from the matrix notation, write $s_i \in S \leq N/T$ for the generator corresponding to the adjacent transposition $(i\ i+1)$, and write $w_i \in W \leq G$ for its permutation-matrix representative. Thus, in the notation of the definition above, we choose

$$\dot{s}_i = w_i.$$

Similarly, for an element $w \in W$, we use the same symbol w for its permutation-matrix representative in G ; in the dotted notation this means $\dot{w} = w$. With these choices, the condition

$$\dot{s}B\dot{w} \subseteq B\dot{w}B \cup B\dot{s}wB$$

becomes

$$w_iBw \subseteq BwB \cup Bw_iwB.$$

We now verify the defining conditions using these representatives.

First, by the Bruhat Decomposition Theorem,

$$G = BWB.$$

Since every permutation matrix is monomial, we have $W \leq N$. Hence

$$G = BWB \subseteq \langle B, N \rangle.$$

The reverse inclusion is automatic because $B \leq G$ and $N \leq G$. Therefore

$$G = \langle B, N \rangle.$$

Second, Exercise 4.7 gives $T \trianglelefteq N$.

Third, Exercise 4.7 gives $N = T \rtimes W$. Thus the quotient N/T is naturally isomorphic to W : the map

$$W \rightarrow N/T, \quad w \mapsto wT$$

is surjective because every element of N has the form tw with $t \in T$ and $w \in W$, and it is injective because its kernel is $T \cap W = \{I\}$. By Proposition 4.5, $W \cong S_n$. Write

$$S = \{s_1, \dots, s_{n-1}\},$$

where s_i is the generator of N/T corresponding to the adjacent transposition

$$(i\ i+1).$$

The representative $w_i \in W$ switches the adjacent basis vectors $\mathbf{v}_i$ and $\mathbf{v}_{i+1}$. Then $w_i^2 = I$, so s_i has order 2 in N/T . The elements s_i generate N/T , because their representatives w_i generate the permutation-matrix subgroup W : by repeatedly swapping neighbouring entries, we can move any chosen entry to any chosen position, and hence build any permutation matrix. Therefore N/T is generated by the finite set S of involutions. Since $|S| = n - 1$, the rank will be $n - 1$ once the remaining two conditions are checked. It remains to verify the two double-coset conditions. If $n = 1$, then $S = \emptyset$ and these conditions are vacuous, so assume $n \geq 2$. Fix $s_i \in S$, and let $w_i \in W$ be its adjacent permutation-matrix representative. Write

$$X_i(\alpha) = X_{i,i+1}(\alpha).$$

We first prove that

$$w_i B w \subseteq B w B \cup B w_i w B$$

for all $w \in W$. Let $b = (b_{rs}) \in B$. Since b is invertible upper triangular, $b_{ii} \neq 0$. Set

$$\alpha = b_{i,i+1} b_{ii}^{-1}, \quad b_0 = b X_i(-\alpha).$$

Right multiplication by $X_i(-\alpha)$ subtracts α times column i from column $i + 1$. Hence the $(i, i + 1)$ -entry of b_0 is

$$b_{i,i+1} - \alpha b_{ii} = 0.$$

Thus $b = b_0 X_i(\alpha)$, with $b_0 \in B$. Moreover $w_i b_0 w_i \in B$. Indeed, conjugating by w_i only swaps the adjacent rows and columns i and $i + 1$; the only possible below-diagonal entry which could appear is the old $(i, i + 1)$ -entry of b_0 , and this entry is zero. Hence $b_0 \in B \cap w_i B w_i$.

Now

$$w_i b w = w_i b_0 X_i(\alpha) w = (w_i b_0 w_i)(w_i X_i(\alpha) w).$$

Since $w_i b_0 w_i \in B$, it is enough to show that $w_i X_i(\alpha) w$ lies in $B w B \cup B w_i w B$. Put

$$p = w^{-1}(i), \quad q = w^{-1}(i + 1).$$

By Lemma 4.6(v),

$$X_i(\alpha) w = w X_{pq}(\alpha).$$

If $p < q$, then $X_{pq}(\alpha) \in B$, and therefore

$$w_i X_i(\alpha) w = w_i w X_{pq}(\alpha) \in B w_i w B.$$

Now suppose that $p > q$. If $\alpha = 0$, then $w_i X_i(0) w = w_i w \in B w_i w B$. If $\alpha \neq 0$, then in the rows and columns $i, i + 1$ we have the 2×2 calculation

$$w_i X_i(\alpha) = \begin{pmatrix} 0 & 1 \\ 1 & \alpha \end{pmatrix} = \begin{pmatrix} -\alpha^{-1} & 1 \\ 0 & \alpha \end{pmatrix} \begin{pmatrix} 1 & 0 \\ \alpha^{-1} & 1 \end{pmatrix}.$$

Thus $w_i X_i(\alpha) = b_1 X_{i+1,i}(\alpha^{-1})$ for some $b_1 \in B$. Therefore

$$w_i X_i(\alpha) w = b_1 X_{i+1,i}(\alpha^{-1}) w = b_1 w X_{qp}(\alpha^{-1}).$$

Since $p > q$, we have $q < p$, so $X_{qp}(\alpha^{-1}) \in B$. Hence

$$w_i X_i(\alpha) w \in B w B.$$

In all cases, $w_i X_i(\alpha) w \in B w B \cup B w_i w B$, and so

$$w_i B w \subseteq B w B \cup B w_i w B.$$

Finally, we verify that the representative w_i satisfies $w_i B w_i \not\subseteq B$. Take $X_i(1) = X_{i,i+1}(1) \in B$. Since $w_i = w_i^{-1}$, Lemma 4.6(v) gives

$$w_i X_i(1) w_i = X_{i+1,i}(1).$$

This matrix has a non-zero entry below the diagonal, so it is not in B . Hence $w_i B w_i \not\subseteq B$.

All the defining conditions of a Tits system are now verified. Therefore (G, B, N, S) is a Tits system. Since S consists of the $n - 1$ adjacent transposition generators, its rank is $n - 1$. $\square$

Exercise 4.9. (cont.) Let G be a group with a BN -pair. Verify that, for $w \in W$, the set $B\dot{w}B$ is independent of the choice of $\dot{w} \in N$ such that $\dot{w}T = w$. Show that we have a Bruhat decomposition

$$G = \bigcup_{w \in W} BwB$$

in which the union is disjoint, where BwB is taken to mean $B\dot{w}B$ for any $\dot{w} \in N$ with $\dot{w}T = w$.

Solution. Let

$$T = B \cap N, \quad W = N/T.$$

For each $w \in W$, choose a representative $\dot{w} \in N$ with $\dot{w}T = w$, and write

$$D_w = B\dot{w}B.$$

We will prove that D_w is well-defined and that

$$G = \coprod_{w \in W} D_w,$$

where $\coprod$ means that the union is disjoint.

Step 1: D_w does not depend on the representative. Suppose that $n, n' \in N$ both represent the same element $w \in W$. This means

$$nT = n'T.$$

Hence $n' = nt$ for some $t \in T$. Since $T = B \cap N$, we have $t \in B$. Therefore

$$Bn'B = BntB = BnB,$$

because the final B already contains t . Thus $B\dot{w}B$ depends only on w , not on the chosen representative $\dot{w}$.

Step 2: multiplying by a simple double coset has only two possible outcomes. Let $s \in S$ and $w \in W$. We claim that

$$D_s D_w \subseteq D_w \cup D_{sw}.$$

Indeed,

$$D_s D_w = B\dot{s}B\dot{w}B = B(\dot{s}B\dot{w})B.$$

By the BN -pair axiom,

$$\dot{s}B\dot{w} \subseteq B\dot{w}B \cup B\dot{s}B\dot{w}.$$

The element $\dot{s}B\dot{w}$ represents $sw \in W$, so $B\dot{s}B\dot{w} = D_{sw}$. Multiplying the inclusion above on the left and right by B gives

$$D_s D_w \subseteq D_w \cup D_{sw}.$$

Also,

$$D_{sw} = B\dot{s}B\dot{w}B \subseteq B\dot{s}B\dot{w}B = D_s D_w.$$

So the useful statement is

$$D_{sw} \subseteq D_s D_w \subseteq D_w \cup D_{sw}.$$

Step 3: the union covers G . Let

$$X = \bigcup_{w \in W} D_w.$$

We show that $X = G$. First,

$$B = D_1 \subseteq X.$$

Also, if $b \in B$, then

$$bD_w \subseteq D_w,$$

because $D_w = B\dot{w}B$ already has B on the left. Hence X is closed under left multiplication by elements of B .

Now take $s \in S$. If $x \in D_w$, then

$$\dot{s}x \in \dot{s}D_w = \dot{s}B\dot{w}B.$$

By Step 2,

$$\dot{s}B\dot{w}B \subseteq D_w \cup D_{sw} \subseteq X.$$

Thus X is closed under left multiplication by each chosen representative $\dot{s}$.

We now explain why these elements generate G . Since $G = \langle B, N \rangle$, it is enough to build every element of N using elements of B and the chosen representatives $\dot{s}$. Let $n \in N$. Since $W = N/T$ is generated by S , we can write

$$nT = s_1 s_2 \cdots s_m$$

with each $s_i \in S$. Then $\dot{s}_1 \dot{s}_2 \cdots \dot{s}_m$ represents the same coset of T as n , so

$$n = \dot{s}_1 \dot{s}_2 \cdots \dot{s}_m t$$

for some $t \in T$. But $T \leq B$. Therefore every element of N is built from elements of B and the $\dot{s}$'s. Hence G is generated by B and the chosen $\dot{s}$'s. Since $1 \in X$ and X is closed under left multiplication by those generators, every element of G lies in X . Therefore

$$G = \bigcup_{w \in W} D_w.$$

Step 4: the union is disjoint. We use the following elementary fact: two double cosets for the same subgroup B are either equal or disjoint. So, to prove disjointness, it is enough to show that

$$D_w = D_v \implies w = v.$$

Assume $D_w = D_v$. We first show that left multiplication by a generator $s \in S$ preserves this equality, in the sense that

$$D_{sw} = D_{sv}.$$

Since $D_w = D_v$, we have

$$D_s D_w = D_s D_v.$$

By Step 2,

$$D_{sw} \subseteq D_s D_w.$$

Thus

$$D_{sw} \subseteq D_s D_v \subseteq D_v \cup D_{sv} = D_w \cup D_{sv}.$$

Since D_{sw} is itself one double coset, it cannot be partly in two different double cosets. Therefore

$$D_{sw} = D_w \quad \text{or} \quad D_{sw} = D_{sv}.$$

Doing the same argument with w and v switched gives

$$D_{sv} = D_w \quad \text{or} \quad D_{sv} = D_{sw}.$$

In every case, $D_{sw} = D_{sv}$: either this is stated directly, or both double cosets are equal to D_w . Now repeat this argument. If $x = s_1 \cdots s_m$ is any product of elements of S , then

$$D_{xw} = D_{xv}.$$

Since S generates W , this holds for every $x \in W$. Choose $x = v^{-1}$. Then

$$D_{v^{-1}w} = D_{v^{-1}v} = D_1 = B.$$

Let $y = v^{-1}w$. Choose a representative $\dot{y} \in N$. Since

$$D_y = B\dot{y}B = B,$$

we have $\dot{y} \in B$. But $\dot{y} \in N$ as well, so $\dot{y} \in B \cap N = T$. Hence

$$y = \dot{y}T = T,$$

which is the identity element of W . Therefore $v^{-1}w = 1$, so $w = v$.

Thus different elements of W give different double cosets, and since double cosets are either equal or disjoint, the union is disjoint. We have proved

$$G = \coprod_{w \in W} B\dot{w}B.$$

Since $B\dot{w}B$ is independent of the chosen representative, we may write this as

$$G = \coprod_{w \in W} BwB,$$

where BwB means $B\dot{w}B$ for any $\dot{w} \in N$ representing w . □

5 Parabolic Subgroups

Setup and Complete Flags

- Throughout, $G = \mathrm{GL}(n, F)$ for some field F and some $n \in \mathbb{N}$, with $V_n(F)$ the space of column vectors of length n over F .
- Elements of G are viewed as matrices, with respect to the standard basis $\mathbf{v}_1, \dots, \mathbf{v}_n$, of invertible linear transformations of $V_n(F)$.
- A *complete flag* on $V_n(F)$ is a sequence of subspaces

$$0 \subset V_1 \subset V_2 \subset \dots \subset V_{n-1} \subset V_n = V_n(F),$$

denoted $(V_1, \dots, V_n)$. Since $\subset$ is proper containment, $\dim_F V_i = i$.

- The *standard flag* is given by $V_i = F\mathbf{v}_1 \oplus \dots \oplus F\mathbf{v}_i = V_{i-1} \oplus F\mathbf{v}_i$ (with $V_0 = 0$).

The G -Action on Complete Flags

- G acts on complete flags by $g(V_1, \dots, V_n) = (gV_1, \dots, gV_n)$.
- This action is transitive: any complete flag $(W_1, \dots, W_n)$ is obtained from the standard flag by the matrix whose i th column is $\mathbf{w}_i \in W_i - W_{i-1}$.
- The stabilizer of the standard flag is precisely the standard Borel subgroup B , so by Lemma 3.2 the Borel subgroups of G are exactly the stabilizers of complete flags on $V_n(F)$.

Upper Unitriangular Matrices and the Decomposition of B

- An upper triangular matrix is *upper unitriangular* if all its main-diagonal entries equal 1, equivalently if it is upper triangular with only eigenvalue 1.
- The set U of all invertible upper unitriangular matrices is a subgroup of B (imitate the proof of Proposition 4.3).
- Let T be the set of diagonal matrices in G . Then $T \leq B$ and $U \cap T = 1$.

Proposition 5.1. $B = U \rtimes T$.

Proof. Let $\varphi: B \rightarrow T$ be the map which sends an upper triangular matrix to the diagonal matrix with the same main diagonal. If $b = (b_{ij})$ and $c = (c_{ij})$ lie in B , then the (i, i) -entry of bc is $b_{ii}c_{ii}$, since all entries below the main diagonal are zero. Hence

$$\varphi(bc) = \varphi(b)\varphi(c),$$

so φ is a homomorphism. Its kernel consists exactly of those upper triangular matrices whose diagonal entries are all 1, so $\ker \varphi = U$. Therefore $U \trianglelefteq B$.

Now let $b \in B$. Since $\varphi(b) \in T$ and φ restricts to the identity map on T , we have

$$\varphi(b\varphi(b)^{-1}) = \varphi(b)\varphi(b)^{-1} = I.$$

Thus $b\varphi(b)^{-1} \in U$, and so

$$b = (b\varphi(b)^{-1})\varphi(b) \in UT.$$

Therefore $B = UT$. Finally, $U \cap T = 1$, because a diagonal matrix whose diagonal entries are all 1 is the identity matrix. Hence $B = U \rtimes T$. $\square$

Action of U and T on the Standard Flag

- An element of U sends each $\mathbf{v}_i$ to $\mathbf{v}_i$ plus some element of V_{i-1} ; conversely, any matrix with this property lies in U .
- Hence U consists of the matrices stabilizing each V_i and inducing the identity on each quotient V_i/V_{i-1} .
- T consists exactly of the matrices stabilizing each of $F\mathbf{v}_1, \dots, F\mathbf{v}_n$; we say T is the *common stabilizer* of the $F\mathbf{v}_i$.

Unipotent Elements and Subgroups

- An element of G is *unipotent* if its characteristic polynomial is $(X - 1)^n$.
- By Jordan form, any unipotent element of G is conjugate to an element of U .
- A subgroup of G is *unipotent* if all its elements are unipotent; U is unipotent and comprises all unipotent elements of B .

Theorem (Kolchin's Theorem). *Any unipotent subgroup of G is conjugate with a subgroup of U .*

Proof. Let H be a unipotent subgroup of G . We first observe that it is enough to show that H stabilizes some complete flag on $V_n(F)$. Indeed, if H stabilizes a complete flag $\mathcal{F}$, then $\mathcal{F} = g\mathcal{F}_0$ for some $g \in G$, where $\mathcal{F}_0$ is the standard flag. By Lemma 3.2, the stabilizer of $\mathcal{F}$ is gBg^{-1} , and hence $g^{-1}Hg \leq B$. Since conjugation preserves characteristic polynomials, every element of $g^{-1}Hg$ is still unipotent. The unipotent elements of B all lie in U , so $g^{-1}Hg \leq U$. Thus H is conjugate with a subgroup of U .

It remains to show that H stabilizes some complete flag. We argue by induction on n . The case $n = 1$ is immediate. Assume $n > 1$. By Proposition 13.28, there exists some $0 \neq \mathbf{v} \in V_n(F)$ such that $x\mathbf{v} = \mathbf{v}$ for every $x \in H$. Let $W = F\mathbf{v}$. Then H stabilizes W , and hence acts on the quotient $V_n(F)/W$. For each $x \in H$, the induced transformation on $V_n(F)/W$ is again unipotent: with respect to a basis beginning with $\mathbf{v}$, the matrix of x has the block form

$$\begin{pmatrix} 1 & * \\ 0 & \bar{x} \end{pmatrix},$$

so the characteristic polynomial of $\bar{x}$ is $(X - 1)^{n-1}$. Therefore the induced subgroup on $V_n(F)/W$ is unipotent. By the induction hypothesis, it stabilizes a complete flag on $V_n(F)/W$. Pulling this flag back to $V_n(F)$ and placing W at the beginning gives a complete flag on $V_n(F)$ stabilized by H . This completes the induction, and hence proves the theorem. $\square$

General Flags and Parabolic Subgroups

- A *flag* on $V_n(F)$ is a nested sequence of non-zero subspaces of $V_n(F)$ which terminates in $V_n(F)$ and has no repeated terms; that is, a sequence $(W_1, \dots, W_r)$ with

$$0 \subset W_1 \subset W_2 \subset \dots \subset W_{r-1} \subset W_r = V_n(F).$$

Since $\subset$ is proper, $r \leq n$; a complete flag is the case $r = n$.

- The G -set of all flags is not transitive: two flags $(W_1, \dots, W_r)$ and $(W'_1, \dots, W'_s)$ lie in the same orbit iff $r = s$ and $\dim_F W_i = \dim_F W'_i$ for all i (same *dimension sequence*).
- A *parabolic subgroup* of G is the stabilizer of some flag on $V_n(F)$.
- Given a flag $(W_1, \dots, W_r)$ with parabolic stabilizer P , choose subspaces Y_i with $W_i = W_{i-1} \oplus Y_i$. The *unipotent radical* of P is the subgroup $U_P \leq P$ of matrices inducing the identity on each W_i/W_{i-1} ; e.g. $U_B = U$.
- A *Levi complement* of U_P is the subgroup $L_P \leq P$ which is the common stabilizer of the Y_i .
- A Levi complement is isomorphic with $\mathrm{GL}(y_1, F) \times \dots \times \mathrm{GL}(y_r, F)$, where $y_i = \dim_F Y_i = \dim_F W_i - \dim_F W_{i-1}$. (In particular, any two Levi complements of U_P are isomorphic.)
- Taking $Y_i = F\mathbf{v}_i$ gives $L_B = T$, isomorphic with a direct product of n copies of $\mathrm{GL}(1, F) \cong F^\times$.

Proposition 5.2. *If P is a parabolic subgroup of G , then we have $P = U_P \rtimes L_P$, where U_P is the unipotent radical of P and L_P is a Levi complement of U_P . Furthermore, $P = N_G(U_P)$.*

Proof. Let P be the stabilizer of the flag $(W_1, \dots, W_r)$, and put $W_0 = 0$. Choose subspaces Y_i such that $W_i = W_{i-1} \oplus Y_i$ for each i . Then

$$V_n(F) = Y_1 \oplus \dots \oplus Y_r.$$

With respect to this direct-sum decomposition, elements of P are block upper triangular. This means that if $p \in P$ and $y_i \in Y_i$, then $p(y_i)$ may have a part in the current block Y_i and earlier blocks

$Y_1, \dots, Y_{i-1}$, but it cannot have a part in any later block. Equivalently, there are maps $a_i: Y_i \rightarrow Y_i$ such that

$$p(y_i) = w_{i-1} + a_i(y_i)$$

with $w_{i-1} \in W_{i-1}$. The map a_i is the action induced by p on the quotient W_i/W_{i-1} , so it is invertible. We now split p into its diagonal-block part and its upper-block-unipotent part. Define $l \in G$ by letting l act as a_i on each Y_i and by making l preserve every Y_i . Then $l \in L_P$, because l is exactly the block diagonal part of p . Set

$$u = pl^{-1}.$$

We claim that $u \in U_P$. Let $y_i \in Y_i$, and write $z_i = l^{-1}(y_i) \in Y_i$. Since l acts as a_i on Y_i , we have $a_i(z_i) = y_i$. Hence

$$u(y_i) = p(z_i) = y_i + w_{i-1}$$

for some $w_{i-1} \in W_{i-1}$. Therefore, modulo W_{i-1} , the vector $u(y_i)$ is the same as y_i . Since the Y_i build the flag one block at a time, this says exactly that u stabilizes every W_i and induces the identity transformation on each quotient W_i/W_{i-1} . Thus $u \in U_P$, and

$$p = ul \in U_P L_P.$$

Since $p \in P$ was arbitrary, $P = U_P L_P$.

This product is a semidirect product. First, U_P is normal in P , since it is the kernel of the homomorphism

$$P \rightarrow \mathrm{GL}(W_1/W_0) \times \cdots \times \mathrm{GL}(W_r/W_{r-1})$$

which records the induced actions on the successive quotients. Second, $U_P \cap L_P = 1$. Indeed, if $x \in U_P \cap L_P$, then x preserves each Y_i , but it also acts as the identity on W_i/W_{i-1} . Therefore x is the identity on every block Y_i , and hence $x = 1$. We have proved

$$P = U_P \rtimes L_P.$$

It remains to prove that $P = N_G(U_P)$. Since $U_P \trianglelefteq P$, every element of P normalizes U_P , so

$$P \leq N_G(U_P).$$

Conversely, suppose that $g \in N_G(U_P)$. We shall prove that g stabilizes every space in the flag, and hence that $g \in P$.

First we recover W_1 from U_P alone. We claim that W_1 is exactly the common fixed subspace of U_P , meaning

$$W_1 = \{v \in V_n(F) \mid uv = v \text{ for every } u \in U_P\}.$$

Every vector in W_1 is fixed because elements of U_P act as the identity on $W_1/W_0 = W_1$. Conversely, take a vector

$$v = y_1 + \cdots + y_r$$

with $y_i \in Y_i$. If some $y_j \neq 0$ for $j > 1$, choose a non-zero vector $x \in Y_1$. Extend y_j to a basis of Y_j , and define a block-unitriangular element u by

$$u(y_j) = y_j + x,$$

while fixing the remaining basis vectors of Y_j and fixing all other blocks. This element lies in U_P : it only adds an earlier-block vector to a vector in the j th block, so it acts as the identity on every quotient W_i/W_{i-1} . But then

$$uv = v + x \neq v.$$

Thus any vector with a non-zero component outside W_1 is moved by some element of U_P . Hence the common fixed subspace of U_P is exactly W_1 .

Now use the assumption that g normalizes U_P . We show that $gW_1 = W_1$. Let $w \in W_1$. To prove $gw \in W_1$, it is enough to show that gw is fixed by every element of U_P . Let $u \in U_P$. Since $g \in N_G(U_P)$, we have $g^{-1}ug \in U_P$. Therefore

$$u(gw) = g(g^{-1}ug)w.$$

The element $g^{-1}ug$ lies in U_P , and $w \in W_1$ is fixed by every element of U_P . Hence $(g^{-1}ug)w = w$, so

$$u(gw) = gw.$$

This holds for every $u \in U_P$, so gw belongs to the common fixed subspace of U_P , which is W_1 . Thus $gW_1 \leq W_1$, and equality follows because g is invertible. Therefore $gW_1 = W_1$. Now pass to the quotient $V_n(F)/W_1$. On this quotient, the group U_P induces the unipotent radical corresponding to the shorter flag

$$W_2/W_1 \subset \cdots \subset W_r/W_1.$$

Because g normalizes U_P and stabilizes W_1 , the induced map of g on $V_n(F)/W_1$ normalizes this induced unipotent radical. The same common-fixed-subspace argument on the quotient shows that the first fixed layer of the quotient, namely W_2/W_1 , is stabilized by g . Hence $gW_2 = W_2$. Repeating this argument one layer at a time gives

$$gW_i = W_i$$

for every i . Thus g stabilizes the whole flag, so $g \in P$. Therefore $N_G(U_P) \leq P$. Combining both inclusions gives $P = N_G(U_P)$. $\square$

Subflags and Staircase Groups

- A *subflag* of the standard flag $(V_1, \dots, V_n)$ is a flag $(W_1, \dots, W_r)$ in which each $W_i = V_j$ for some j ; there are 2^{n-1} such subflags.
- A *staircase group* is a parabolic subgroup of G that stabilizes some subflag of the standard flag.
- Example ($n = 6$, subflag $0 \subset V_2 \subset V_3 \subset V_6$): the staircase group consists of all matrices in $G = \text{GL}(6, F)$ of the form

$$\begin{pmatrix} * & * & * & * & * & * \\ * & * & * & * & * & * \\ 0 & 0 & * & * & * & * \\ 0 & 0 & 0 & * & * & * \\ 0 & 0 & 0 & * & * & * \\ 0 & 0 & 0 & * & * & * \end{pmatrix}.$$

Imagine a “staircase” separating the zero entries from the arbitrary entries.

- Its unipotent radical consists of all matrices in G of the form

$$\begin{pmatrix} 1 & 0 & * & * & * & * \\ 0 & 1 & * & * & * & * \\ 0 & 0 & 1 & * & * & * \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix},$$

and the Levi complement (with canonical choices $Y_2 = F\mathbf{v}_3$ and $Y_3 = F\mathbf{v}_4 \oplus F\mathbf{v}_5 \oplus F\mathbf{v}_6$) consists of all matrices in G of the form

$$\begin{pmatrix} * & * & 0 & 0 & 0 & 0 \\ * & * & 0 & 0 & 0 & 0 \\ 0 & 0 & * & 0 & 0 & 0 \\ 0 & 0 & 0 & * & * & * \\ 0 & 0 & 0 & * & * & * \\ 0 & 0 & 0 & * & * & * \end{pmatrix},$$

isomorphic with $\text{GL}(2, F) \times \text{GL}(1, F) \times \text{GL}(3, F)$.

- In general, the unipotent radical of any staircase group is contained in B .
- The G -set of all flags is partitioned into the orbits of the subflags of the standard flag, since any flag has the same dimension sequence as some subflag of the standard flag. By Lemma 3.2, the parabolic subgroups of G are exactly the conjugates of the staircase groups.

Subgroups Containing B

Lemma 5.3. *The only non-zero subspaces of $V_n(F)$ left invariant by B are the subspaces $V_1, \dots, V_n$ that appear in the standard flag.*

Proof. First observe that each V_i is stabilized by B : if $j \leq i$ and $b \in B$, then $b\mathbf{v}_j$ is a linear combination of $\mathbf{v}_1, \dots, \mathbf{v}_j$, hence lies in V_i . Therefore $bV_i \subseteq V_i$, and equality follows since b is invertible.

Conversely, let Y be a non-zero subspace of $V_n(F)$ stabilized by B . Choose k minimal such that $Y \subseteq V_k$. Then $Y \not\subseteq V_{k-1}$, so there is some

$$y = \alpha_1 \mathbf{v}_1 + \dots + \alpha_k \mathbf{v}_k \in Y$$

with $\alpha_k \neq 0$. Let $b \in B$ be the matrix which is the identity except that its k th column is y :

$$b = \begin{pmatrix} & \alpha_1 & & \\ I_{k-1} & \vdots & & 0 \\ & \alpha_{k-1} & & \\ 0 & \alpha_k & & 0 \\ 0 & 0 & & I_{n-k} \end{pmatrix}.$$

This matrix is upper triangular and invertible, since its k th diagonal entry is $\alpha_k \neq 0$. Also $b\mathbf{v}_k = y$. Since Y is stabilized by B and $b^{-1} \in B$, we have

$$\mathbf{v}_k = b^{-1}y \in Y.$$

Now let

$$z = \beta_1 \mathbf{v}_1 + \dots + \beta_k \mathbf{v}_k \in V_k - V_{k-1}.$$

Then $\beta_k \neq 0$, and the same matrix construction gives some $c \in B$ with $c\mathbf{v}_k = z$. Since $\mathbf{v}_k \in Y$ and Y is stabilized by B , it follows that $z \in Y$. Thus every vector in $V_k - V_{k-1}$ lies in Y . In particular, if $w \in V_{k-1}$, then $w + \mathbf{v}_k \in Y$, and since $\mathbf{v}_k \in Y$, we get $w \in Y$. Hence $V_{k-1} \subseteq Y$, and together with $\mathbf{v}_k \in Y$ this gives $V_k \subseteq Y$. Since $Y \subseteq V_k$ by choice of k , we conclude that $Y = V_k$. $\square$

Theorem 5.4. *The only subgroups of G that contain B are the staircase groups.*

Proof. First observe that every staircase group contains B , because B stabilizes every subspace in the standard flag and therefore stabilizes every subflag of the standard flag. We prove the converse. Let H be a subgroup with

$$B \leq H \leq G.$$

If a subspace of $V_n(F)$ is stabilized by H , then it is also stabilized by B . Hence Lemma 5.3 shows that the only possible non-zero H -invariant subspaces are among

$$V_1, \dots, V_n.$$

List the ones which are actually stabilized by H as

$$V_{a_1}, \dots, V_{a_r}, \quad 1 \leq a_1 < \dots < a_r = n,$$

and put $a_0 = 0$. Let P be the staircase group stabilizing the subflag

$$0 \subset V_{a_1} \subset \dots \subset V_{a_r} = V_n(F).$$

Then $H \leq P$. We shall prove that $H = P$.

First we prove the basic case in which the only H -invariant subspaces are 0 and $V_n(F)$. In this case we claim that $H = G$. If $n = 1$ this is immediate, so assume $n > 1$. The subspace

$$Y = \langle h\mathbf{v}_1 \mid h \in H \rangle$$

is non-zero and H -invariant, so by assumption $Y = V_n(F)$. Thus some $h \in H$ sends $\mathbf{v}_1$ to a vector

$$h\mathbf{v}_1 = \alpha_1 \mathbf{v}_1 + \dots + \alpha_n \mathbf{v}_n$$

with $\alpha_n \neq 0$. By Bruhat Decomposition Theorem, write $h = b_1 w b_2$ with $b_1, b_2 \in B$ and $w \in W$. Since $B \leq H$ and $h \in H$, it follows that $w \in H$. By Corollary 4.10, the fact that the last non-zero term in

$h\mathbf{v}_1$ is the $\mathbf{v}_n$ -term implies that $w(1) = n$. Let $j = w^{-1}(1)$. Then $j > 1$, so $X_{1j} \leq B \leq H$. Using Lemma 4.6(v), we get

$$wX_{1j}(\alpha)w^{-1} = X_{n1}(\alpha)$$

for every $\alpha \in F$. Hence $X_{n1} \leq H$.

We now get all the remaining root subgroups from this one lower root subgroup. The upper root subgroups X_{ij} with $i < j$ already lie in B , and hence in H . If $n = 2$, this already gives every root subgroup. If $n > 2$, then for $1 < i < n$ and $\alpha \in F$, Lemma 4.6(iv) gives

$$X_{ni}(\alpha) = [X_{n1}(\alpha), X_{1i}(1)] \in H$$

and

$$X_{i1}(\alpha) = [X_{in}(\alpha), X_{n1}(1)] \in H.$$

For distinct $1 < i, j < n$, the same commutator formula gives

$$X_{ij}(\alpha) = [X_{i1}(\alpha), X_{1j}(1)] \in H.$$

Thus H contains every root subgroup X_{ij} , and it also contains every invertible diagonal matrix because all diagonal matrices lie in B . By Theorem 4.12, $H = G$. This proves the basic case.

We return to the general subgroup H . By Proposition 5.2, the staircase group P decomposes as

$$P = U_P \rtimes L_P.$$

Since P is a staircase group, its unipotent radical U_P is contained in B , so $U_P \leq H$. Therefore it is enough to show that a Levi complement L_P is contained in H . We choose the standard Levi complement whose blocks are

$$F\mathbf{v}_{a_{k-1}+1} \oplus \cdots \oplus F\mathbf{v}_{a_k}, \quad 1 \leq k \leq r.$$

Fix one block, and write

$$s = a_{k-1} + 1, \quad t = a_k.$$

If $s = t$, there is nothing to prove for this block. Assume $s < t$. Because there is no H -invariant standard flag subspace strictly between $V_{a_{k-1}}$ and V_{a_k} , the subspace

$$V_{a_{k-1}} + \langle h\mathbf{v}_s \mid h \in H \rangle$$

must be all of V_t . Hence there is some $h \in H$ such that the coefficient of $\mathbf{v}_t$ in $h\mathbf{v}_s$ is non-zero. Write $h = b_1 w b_2$ by Bruhat Decomposition Theorem. Again $w \in H$. Also $w \in P$, because $b_1, b_2 \in B \leq P$ and $h \in P$. Thus w preserves each set of indices

$$\{1, \dots, a_\ell\}$$

which comes from the subflag stabilized by P . In particular, w sends the block $\{s, \dots, t\}$ to itself. The row-pivot description in Corollary 4.10 applied to the s th column says that $w(s) = t$, because the first $s - 1$ columns use only rows $1, \dots, s - 1$, while the s th column has no entries below row t and has a non-zero entry in row t . Let $q = w^{-1}(s)$. Since w preserves the block and $w(s) = t$, we have $s < q \leq t$. Therefore $X_{sq} \leq B \leq H$, and Lemma 4.6(v) gives

$$wX_{sq}(\alpha)w^{-1} = X_{ts}(\alpha)$$

for every $\alpha \in F$. Hence $X_{ts} \leq H$.

Now the same commutator ladder used in the basic case works inside this block. All upper root subgroups X_{ij} with $s \leq i < j \leq t$ lie in B , hence in H . Together with $X_{ts} \leq H$, this gives every root subgroup inside the block: for $s < i < t$,

$$X_{ti}(\alpha) = [X_{ts}(\alpha), X_{si}(1)] \in H$$

and

$$X_{is}(\alpha) = [X_{it}(\alpha), X_{ts}(1)] \in H,$$

while for distinct $s < i, j < t$,

$$X_{ij}(\alpha) = [X_{is}(\alpha), X_{sj}(1)] \in H.$$

Thus, for each block of L_P , the subgroup H contains all root subgroups supported inside that block. Since H also contains all diagonal matrices, Theorem 4.12 applied block-by-block shows that $L_P \leq H$. Hence

$$P = U_P \rtimes L_P \leq H.$$

We already had $H \leq P$, so $H = P$. Therefore every subgroup of G containing B is a staircase group. $\square$

Exercises

Let $G = \text{GL}(n, F)$, where $n \geq 2$.

Exercise 5.1. Suppose that $g \in G$ fixes some $0 \neq \mathbf{v} \in V_n(F)$ and induces the identity transformation on $V_n(F)/F\mathbf{v}$. Show that g is conjugate with an element of the root subgroup X_{12} .

Solution. Since g induces the identity transformation on $V_n(F)/F\mathbf{v}$, for every $\mathbf{x} \in V_n(F)$ we have

$$g\mathbf{x} + F\mathbf{v} = \mathbf{x} + F\mathbf{v}.$$

Equivalently,

$$g\mathbf{x} - \mathbf{x} \in F\mathbf{v}.$$

Thus there is a scalar $\lambda(\mathbf{x}) \in F$ such that

$$g\mathbf{x} = \mathbf{x} + \lambda(\mathbf{x})\mathbf{v}.$$

The scalar $\lambda(\mathbf{x})$ is unique because $\mathbf{v} \neq 0$. The assignment $\lambda: V_n(F) \rightarrow F$ is linear: for example, comparing

$$g(\mathbf{x} + \mathbf{y}) = (\mathbf{x} + \mathbf{y}) + \lambda(\mathbf{x} + \mathbf{y})\mathbf{v}$$

with

$$g\mathbf{x} + g\mathbf{y} = \mathbf{x} + \lambda(\mathbf{x})\mathbf{v} + \mathbf{y} + \lambda(\mathbf{y})\mathbf{v}$$

gives $\lambda(\mathbf{x} + \mathbf{y}) = \lambda(\mathbf{x}) + \lambda(\mathbf{y})$, and scalar multiplication is similar. Also $g\mathbf{v} = \mathbf{v}$, so $\lambda(\mathbf{v}) = 0$.

If $\lambda = 0$, then $g\mathbf{x} = \mathbf{x}$ for every $\mathbf{x}$, so $g = I = X_{12}(0)$, and there is nothing to prove. Assume now that $\lambda \neq 0$. Choose $\mathbf{w}_1 = \mathbf{v}$. Since $\lambda \neq 0$, there exists some vector $\mathbf{w}_2$ with

$$\lambda(\mathbf{w}_2) = \alpha \neq 0.$$

Then

$$g\mathbf{w}_2 = \mathbf{w}_2 + \alpha\mathbf{w}_1.$$

The kernel of λ has dimension $n - 1$. Since $\lambda(\mathbf{w}_1) = 0$, we have $\mathbf{w}_1 \in \ker \lambda$. Choose vectors $\mathbf{w}_3, \dots, \mathbf{w}_n$ such that

$$\mathbf{w}_1, \mathbf{w}_3, \dots, \mathbf{w}_n$$

is a basis of $\ker \lambda$. Since $\mathbf{w}_2 \notin \ker \lambda$, the list $\mathbf{w}_1, \mathbf{w}_2, \dots, \mathbf{w}_n$ is a basis of $V_n(F)$. In this basis, g sends

$$\mathbf{w}_1 \mapsto \mathbf{w}_1, \quad \mathbf{w}_2 \mapsto \mathbf{w}_2 + \alpha\mathbf{w}_1, \quad \mathbf{w}_i \mapsto \mathbf{w}_i \quad (i \geq 3).$$

Therefore the matrix of g in the basis $\mathbf{w}_1, \dots, \mathbf{w}_n$ is $X_{12}(\alpha)$. If $p \in G$ is the change-of-basis matrix whose columns are $\mathbf{w}_1, \dots, \mathbf{w}_n$, then

$$p^{-1}gp = X_{12}(\alpha).$$

Hence g is conjugate with an element of the root subgroup X_{12} . □

Exercise 5.2. We say a basis $\mathcal{B}$ of $V_n(F)$ belongs to a complete flag $(V_1, \dots, V_n)$ if each V_i contains exactly one element of $\mathcal{B}$ that is not in V_{i-1} .

(a) Show that if $(V_1, \dots, V_n)$ and $(W_1, \dots, W_n)$ are complete flags on $V_n(F)$, then there is a basis of $V_n(F)$ that belongs to both flags.

(b) Use part (a) to give a different proof of the Bruhat Decomposition Theorem.

Solution.

(a) Put $V_0 = W_0 = 0$. We build the required basis one vector at a time from intersections of the two flags. Fix $i \in \{1, \dots, n\}$. Since $W_n = V_n(F)$, we have

$$V_i \cap W_n = V_i,$$

and this is not contained in V_{i-1} . Therefore there is a smallest integer j_i such that

$$V_i \cap W_{j_i} \not\subseteq V_{i-1}.$$

Choose

$$\mathbf{b}_i \in (V_i \cap W_{j_i}) - V_{i-1}.$$

This means that $\mathbf{b}_i$ enters the V -flag at the i th step, and it enters the W -flag no later than the j_i th step.

First we check that these vectors form a basis belonging to the V -flag. Since $\mathbf{b}_1 \in V_1$ and $\mathbf{b}_1 \neq 0$, the first vector works. Suppose that $\mathbf{b}_1, \dots, \mathbf{b}_{i-1}$ are independent and span V_{i-1} . Since $\mathbf{b}_i \notin V_{i-1}$, the vector $\mathbf{b}_i$ is not in the span of the previous vectors. Hence $\mathbf{b}_1, \dots, \mathbf{b}_i$ are independent. They are i independent vectors inside the i -dimensional space V_i , so they span V_i . By induction,

$$V_i = \langle \mathbf{b}_1, \dots, \mathbf{b}_i \rangle$$

for every i . Thus the basis belongs to the V -flag.

It remains to show that the same basis also belongs to the W -flag. Fix $m \in \{1, \dots, n\}$. For each i , the space

$$V_{i-1} \cap W_m$$

is contained in $V_i \cap W_m$, and the dimension can increase by at most 1 because V_i/V_{i-1} is one-dimensional. By the definition of j_i , this increase happens exactly when $j_i \leq m$. Therefore

$$\#\{i \mid j_i \leq m\} = \sum_{i=1}^n (\dim(V_i \cap W_m) - \dim(V_{i-1} \cap W_m)).$$

The sum telescopes:

$$\sum_{i=1}^n (\dim(V_i \cap W_m) - \dim(V_{i-1} \cap W_m)) = \dim(V_n \cap W_m) - \dim(V_0 \cap W_m) = \dim W_m = m.$$

Also, by the minimality of j_i , the vector $\mathbf{b}_i$ lies in W_m exactly when $j_i \leq m$. Thus exactly m of the vectors $\mathbf{b}_1, \dots, \mathbf{b}_n$ lie in W_m . These m vectors are independent and lie inside the m -dimensional space W_m , so they form a basis of W_m . Hence, when we pass from W_{m-1} to W_m , exactly one new basis vector appears. Therefore the same basis belongs to the W -flag as well.

(b) Let

$$\mathcal{F}_0 = (V_1, \dots, V_n)$$

be the standard complete flag, and let $g \in G$. Then

$$g\mathcal{F}_0 = (gV_1, \dots, gV_n)$$

is also a complete flag. By part (a), there is a basis

$$\mathcal{B} = \{\mathbf{b}_1, \dots, \mathbf{b}_n\}$$

which belongs to both $\mathcal{F}_0$ and $g\mathcal{F}_0$. Order the basis so that

$$\mathbf{b}_i \in V_i - V_{i-1}$$

for every i . Let a be the matrix whose i th column is $\mathbf{b}_i$. Since the i th column of a lies in V_i , the matrix a is upper triangular. Since its columns form a basis, a is invertible. Hence

$$a \in B.$$

Now use the fact that the same basis also belongs to $g\mathcal{F}_0$. For each r , exactly one basis vector lies in

$$gV_r - gV_{r-1}.$$

Thus there is a permutation σ of $\{1, \dots, n\}$ such that

$$\mathbf{b}_{\sigma(r)} \in gV_r - gV_{r-1}$$

for every r . Let $w \in W$ be the permutation matrix which reorders the columns of a so that the r th column of aw is $\mathbf{b}_{\sigma(r)}$. Then

$$g^{-1}\mathbf{b}_{\sigma(r)} \in V_r - V_{r-1}$$

for every r . Therefore the r th column of $g^{-1}aw$ lies in V_r . It follows that $g^{-1}aw$ is upper triangular. It is also invertible, so

$$g^{-1}aw \in B.$$

Write

$$c = g^{-1}aw \in B.$$

Then

$$aw = gc,$$

and therefore

$$g = awc^{-1}.$$

Since $a \in B$, $w \in W$, and $c^{-1} \in B$, we have $g \in BWB$. As $g \in G$ was arbitrary, this gives

$$G \subseteq BWB.$$

The reverse inclusion $BWB \subseteq G$ is immediate from $B \leq G$ and $W \leq G$. Hence

$$G = BWB,$$

which is the Bruhat Decomposition Theorem. □

Exercise 5.3. Let P and Q be distinct parabolic subgroups of G containing the standard Borel subgroup B of G (so that, by Theorem 5.4, P and Q are staircase groups).

- (a) Show that P and Q are not conjugate in G .
- (b) If L_P and L_Q are corresponding Levi complements, find conditions on P and Q that determine when L_P and L_Q will be conjugate in G .

Solution.

- (a) By Theorem 5.4, both P and Q are staircase groups. Thus there are subflags of the standard flag

$$0 \subset V_{a_1} \subset \cdots \subset V_{a_r} = V_n(F)$$

and

$$0 \subset V_{b_1} \subset \cdots \subset V_{b_s} = V_n(F)$$

such that P is the stabilizer of the first subflag and Q is the stabilizer of the second. If the two lists of indices were the same, then the two stabilized standard subflags would be the same, and hence their stabilizers would be the same. Since $P \neq Q$, we know that

$$\{a_1, \dots, a_r\} \neq \{b_1, \dots, b_s\}.$$

We first record the invariant-subspace fact we need. Let R be the staircase group stabilizing

$$0 \subset V_{c_1} \subset \cdots \subset V_{c_t} = V_n(F),$$

and put $c_0 = 0$. We claim that the only non-zero subspaces of $V_n(F)$ stabilized by R are

$$V_{c_1}, \dots, V_{c_t}.$$

This is the block version of Lemma 5.3. Let Y be a non-zero R -invariant subspace. Choose k minimal such that

$$Y \leq V_{c_k}.$$

Then $Y \not\leq V_{c_{k-1}}$, so the image of Y in the quotient

$$V_{c_k}/V_{c_{k-1}}$$

is non-zero. By Proposition 5.2 and the block description of a staircase group, the standard Levi complement of R acts as the full general linear group on the block

$$F\mathbf{v}_{c_{k-1}+1} \oplus \cdots \oplus F\mathbf{v}_{c_k}.$$

Hence it acts on $V_{c_k}/V_{c_{k-1}}$ as the full group $\text{GL}(c_k - c_{k-1}, F)$. Since Y is R -invariant, the image of Y in this quotient is invariant under this full general linear group. But the only non-zero subspace invariant under the full general linear group is the whole quotient, because an invertible linear map can send any non-zero vector to any other non-zero vector. Therefore

$$Y + V_{c_{k-1}} = V_{c_k}.$$

We next show that $V_{c_{k-1}} \leq Y$. If $k = 1$, this is immediate because $V_{c_0} = 0$. Assume $k > 1$. Since $Y + V_{c_{k-1}} = V_{c_k}$, there is some vector $y \in Y$ whose component in the k th block is non-zero. Choose an index m in the k th block such that the coefficient of $\mathbf{v}_m$ in y is a non-zero scalar β . Now let $x \in V_{c_{k-1}}$. Write

$$x = \gamma_1 \mathbf{v}_1 + \cdots + \gamma_{c_{k-1}} \mathbf{v}_{c_{k-1}}.$$

For each $\ell \leq c_{k-1}$, the root subgroup element $X_{\ell m}(\alpha)$ sends $\mathbf{v}_m$ to $\mathbf{v}_m + \alpha \mathbf{v}_\ell$. Therefore, if we choose

$$u = \prod_{\ell=1}^{c_{k-1}} X_{\ell m}(\gamma_\ell / \beta),$$

then u sends y to $y + x$. These are upper root subgroup elements because $\ell < m$, so they lie in B . Since $B \leq R$, we have $u \in R$. Since Y is R -invariant, both y and $uy = y + x$ lie in Y , and hence

$$x = (y + x) - y \in Y.$$

Since $x \in V_{c_{k-1}}$ was arbitrary, $V_{c_{k-1}} \leq Y$. Together with $Y + V_{c_{k-1}} = V_{c_k}$, this gives $V_{c_k} \leq Y$. Since $Y \leq V_{c_k}$ by the choice of k , we get $Y = V_{c_k}$. This proves the claim.

Now suppose, for a contradiction, that P and Q are conjugate in G . Then there exists some $g \in G$ such that

$$Q = gPg^{-1}.$$

For each i , the subspace V_{a_i} is stabilized by P . We claim that gV_{a_i} is stabilized by Q . Indeed, if $q \in Q$, then $q = gpg^{-1}$ for some $p \in P$. If $gv \in gV_{a_i}$ with $v \in V_{a_i}$, then

$$q(gv) = gpg^{-1}gv = gpv.$$

Since p stabilizes V_{a_i} , we have $pv \in V_{a_i}$, and hence $gpv \in gV_{a_i}$. Thus gV_{a_i} is Q -invariant. By the claim applied to Q , the subspace gV_{a_i} must be one of the subspaces

$$V_{b_1}, \dots, V_{b_s}.$$

Taking dimensions gives

$$a_i = \dim gV_{a_i} = b_j$$

for some j . Therefore

$$\{a_1, \dots, a_r\} \subseteq \{b_1, \dots, b_s\}.$$

Applying the same argument to $P = g^{-1}Qg$ gives the reverse inclusion. Hence

$$\{a_1, \dots, a_r\} = \{b_1, \dots, b_s\},$$

contradicting the fact that these two sets are different. Therefore no such g exists, and P and Q are not conjugate in G .

- (b) Keep the notation from part (a), and put $a_0 = b_0 = 0$. The standard Levi complement of P is the subgroup which preserves each block

$$Y_i = F\mathbf{v}_{a_{i-1}+1} \oplus \cdots \oplus F\mathbf{v}_{a_i},$$

and acts on that block as the full group $\text{GL}(Y_i)$. Thus its block sizes are

$$d_i = \dim Y_i = a_i - a_{i-1} \quad (1 \leq i \leq r).$$

Similarly, the standard Levi complement of Q has blocks

$$Z_j = F\mathbf{v}_{b_{j-1}+1} \oplus \cdots \oplus F\mathbf{v}_{b_j}$$

of sizes

$$e_j = \dim Z_j = b_j - b_{j-1} \quad (1 \leq j \leq s).$$

We claim that L_P and L_Q are conjugate in G exactly when the two lists

$$d_1, \dots, d_r \quad \text{and} \quad e_1, \dots, e_s$$

agree up to permutation.

First suppose that the block sizes agree up to permutation. Then $r = s$, and after reordering the Q -blocks there is a bijection σ such that

$$d_i = e_{\sigma(i)}$$

for every i . Choose a permutation matrix w which sends the standard basis vectors in Y_i onto the standard basis vectors in $Z_{\sigma(i)}$. Conjugating by w sends an arbitrary invertible linear transformation on Y_i to an arbitrary invertible linear transformation on $Z_{\sigma(i)}$. Therefore

$$wL_Pw^{-1} = L_Q.$$

Hence L_P and L_Q are conjugate.

Conversely, suppose that $L_Q = gL_Pg^{-1}$ for some $g \in G$. Conjugation by g sends the block decomposition

$$V_n(F) = Y_1 \oplus \dots \oplus Y_r$$

to

$$V_n(F) = gY_1 \oplus \dots \oplus gY_r.$$

On each gY_i , the conjugate subgroup $gL_Pg^{-1} = L_Q$ contains a factor acting as the full group $\text{GL}(gY_i)$. Thus the spaces gY_i are the same kind of full general-linear blocks for L_Q as the standard blocks Z_j . The standard block decomposition of L_Q has block dimensions $e_1, \dots, e_s$, so the dimensions

$$\dim gY_i = \dim Y_i = d_i$$

must be exactly the same dimensions as the e_j , possibly in a different order. Hence the block sizes of P and Q agree up to permutation.

Therefore the condition is:

$$L_P \text{ and } L_Q \text{ are conjugate in } G \iff \{a_i - a_{i-1} \mid 1 \leq i \leq r\} = \{b_j - b_{j-1} \mid 1 \leq j \leq s\}$$

as multisets.

□

Exercise 5.4. Prove Proposition 5.2.

Solution. This is exactly Proposition 5.2, proved above.

□

Exercise 5.5. Prove the following generalization of Lemma 5.3: If P is a parabolic subgroup of G which is the stabilizer of the flag $(W_1, \dots, W_r)$, then the W_i are the only subspaces of $V_n(F)$ left invariant by P .

Solution. Let

$$0 \subset W_1 \subset \dots \subset W_r = V_n(F)$$

be the flag stabilized by P , and put $c_i = \dim_F W_i$. By the transitivity of the G -action on flags with a fixed dimension sequence, choose $g \in G$ such that

$$gW_i = V_{c_i}$$

for every i . Hence gPg^{-1} is the staircase group stabilizing the standard subflag

$$0 \subset V_{c_1} \subset \dots \subset V_{c_r} = V_n(F).$$

Now let Y be a non-zero subspace of $V_n(F)$ stabilized by P . Then gY is stabilized by gPg^{-1} : if $q = gpg^{-1} \in gPg^{-1}$ and $gy \in gY$, with $p \in P$ and $y \in Y$, then

$$q(gy) = gpy \in gY.$$

By the staircase-group claim proved in Exercise 5.3(a), the only non-zero subspaces stabilized by gPg^{-1} are

$$V_{c_1}, \dots, V_{c_r}.$$

Therefore $gY = V_{c_i}$ for some i . Applying g^{-1} gives

$$Y = g^{-1}V_{c_i} = W_i.$$

Thus the only non-zero subspaces stabilized by P are $W_1, \dots, W_r$. Each W_i is stabilized by P by the definition of P as the stabilizer of the flag, so these are exactly the non-zero P -invariant subspaces. $\square$

Exercise 5.6. (cont.) Using Exercise 5.5, show that any parabolic subgroup of G is self-normalizing.

Solution. Let P be the stabilizer of the flag

$$0 \subset W_1 \subset \dots \subset W_r = V_n(F).$$

We prove that $N_G(P) = P$. The inclusion

$$P \leq N_G(P)$$

is automatic, because every element of a subgroup normalizes that subgroup. It remains to prove the reverse inclusion.

Let $g \in N_G(P)$. We want to show that $g \in P$, i.e. that g stabilizes every W_i . Fix i . We first show that gW_i is stabilized by P . Let $p \in P$ and let $gw \in gW_i$, where $w \in W_i$. Since $g \in N_G(P)$, we have

$$g^{-1}pg \in P.$$

Since P stabilizes W_i , this gives

$$(g^{-1}pg)w \in W_i.$$

Applying g to both sides, we get

$$p(gw) = g(g^{-1}pg)w \in gW_i.$$

Thus gW_i is a non-zero subspace stabilized by P . By Exercise 5.5, the only non-zero subspaces stabilized by P are exactly the spaces in its flag, so

$$gW_i = W_j$$

for some j . But g preserves dimension, so

$$\dim_F W_j = \dim_F(gW_i) = \dim_F W_i.$$

The dimensions in a flag are strictly increasing, so $j = i$. Hence

$$gW_i = W_i$$

for every i . Therefore g stabilizes the whole flag, and so $g \in P$. This proves

$$N_G(P) \leq P.$$

Combining the two inclusions gives $N_G(P) = P$, so every parabolic subgroup of G is self-normalizing. $\square$

Exercise 5.7. Complete the outlined proof of Theorem 5.4.

Solution. Let H be a subgroup of G with $B \leq H \leq G$. If a subspace is stabilized by H , then it is also stabilized by B . Hence Lemma 5.3 says that the only possible non-zero subspaces stabilized by H are among the standard flag subspaces. List the ones actually stabilized by H as

$$V_{a_1}, \dots, V_{a_r}, \quad 1 \leq a_1 < \dots < a_r = n,$$

and put $a_0 = 0$. Let P be the staircase group stabilizing the subflag

$$0 \subset V_{a_1} \subset \dots \subset V_{a_r} = V_n(F).$$

Then $H \leq P$, because every element of H stabilizes each of the spaces in this subflag. We prove the reverse inclusion.

Step 1: reduce to the Levi complement. By Proposition 5.2, we may write

$$P = U_P \rtimes L_P.$$

Since P is a staircase group, its unipotent radical U_P is contained in B . Hence

$$U_P \leq B \leq H.$$

Therefore, to prove $P \leq H$, it remains only to prove

$$L_P \leq H.$$

Choose the standard Levi complement whose blocks are

$$F\mathbf{v}_{a_{k-1}+1} \oplus \dots \oplus F\mathbf{v}_{a_k}, \quad 1 \leq k \leq r.$$

We will show that H contains the full general linear group on each one of these blocks.

Step 2: find one lower root subgroup in a fixed block. Fix a block, and write

$$s = a_{k-1} + 1, \quad t = a_k.$$

If $s = t$, then the block is one-dimensional, and there are no root subgroups to find in it. Assume $s < t$. Since there is no H -invariant standard flag subspace strictly between $V_{a_{k-1}}$ and V_{a_k} , the H -invariant subspace

$$V_{a_{k-1}} + \langle h\mathbf{v}_s \mid h \in H \rangle$$

is stabilized by H . It contains $\mathbf{v}_s$, so it is larger than $V_{a_{k-1}}$; it is contained in V_t , because $H \leq P$ and P stabilizes V_t . Therefore it must equal V_t . Thus there is some $h \in H$ such that the coefficient of $\mathbf{v}_t$ in $h\mathbf{v}_s$ is non-zero.

Write $h = b_1 w b_2$ by the Bruhat Decomposition Theorem, with $b_1, b_2 \in B$ and $w \in W$. Since $B \leq H$ and $h \in H$, we also have $w \in H$. Since $H \leq P$, the element w lies in P , so it permutes the indices inside each block of the staircase. The row-pivot description in Corollary 4.10, applied to the s th column, gives

$$w(s) = t.$$

Indeed, the first $s - 1$ columns of h lie in V_{s-1} , while the s th column lies in V_t and has a non-zero $\mathbf{v}_t$ -coefficient. Thus the row chosen by the Bruhat pivot construction for column s is row t .

Let $q = w^{-1}(s)$. Because w preserves the block $\{s, \dots, t\}$ and sends s to t , we have

$$s < q \leq t.$$

Hence $X_{sq} \leq B \leq H$. By Lemma 4.6(v),

$$wX_{sq}(\alpha)w^{-1} = X_{ts}(\alpha)$$

for every $\alpha \in F$. Therefore

$$X_{ts} \leq H.$$

Step 3: use commutators to get all root subgroups in the block. The upper root subgroups X_{ij} with $s \leq i < j \leq t$ already lie in B , and hence in H . Together with $X_{ts} \leq H$, the commutator formula in Lemma 4.6(iv) gives the remaining root subgroups inside the block. For $s < i < t$ and $\alpha \in F$,

$$X_{ti}(\alpha) = [X_{ts}(\alpha), X_{si}(1)] \in H$$

and

$$X_{is}(\alpha) = [X_{it}(\alpha), X_{ts}(1)] \in H.$$

For distinct $s < i, j < t$,

$$X_{ij}(\alpha) = [X_{is}(\alpha), X_{sj}(1)] \in H.$$

Thus H contains every root subgroup X_{ij} supported inside this block. Since H also contains all diagonal matrices, Theorem 4.12 applied to this block shows that H contains the full general linear group on the block.

Step 4: finish block by block. Repeating Steps 2 and 3 for every block of the staircase shows that every block factor of L_P is contained in H . Hence

$$L_P \leq H.$$

Since $U_P \leq H$ as well, we have

$$P = U_P \rtimes L_P \leq H.$$

We already had $H \leq P$, so $H = P$. Therefore every subgroup of G containing B is a staircase group, which completes the general case of Theorem 5.4. $\square$

Exercise 5.8. Assume that the case $r = 1$ of Theorem 5.4 holds; we sketch an alternate proof of the general case of Theorem 5.4. We use induction on n , where $G = \mathrm{GL}(n, F)$. Let H be a subgroup of G which contains B , and suppose that H stabilizes V_m , where $1 \leq m < n$. Let S be the stabilizer of V_m ; observe that $H \leq S$. We can write $S = U \rtimes (K_1 \times K_2)$, where the elements of $K_1 \cong \mathrm{GL}(m, F)$ have their only non-zero entries in the first m rows and columns, and where K_2 is a direct product of general linear groups whose elements have their non-zero entries concentrated in the last $n - m$ rows and columns. Since $U \leq B \leq H$, by Exercise 2.10 we have $H = U \rtimes Q$ for some $Q \leq K_1 \times K_2$. Use Goursat's theorem (Exercise 2.8), Exercise 5.6, and the induction hypothesis to analyze Q and thereby conclude that H is a staircase group.

Solution. We prove the general case of Theorem 5.4 by strong induction on n , assuming that the case $r = 1$ is already known. Here the case $r = 1$ means the following: if $B \leq H \leq \mathrm{GL}(n, F)$ and H stabilizes no non-zero proper standard flag subspace, then $H = \mathrm{GL}(n, F)$. The case $n = 1$ is immediate, so assume $n > 1$ and that the result is known for all smaller dimensions.

Let H be a subgroup with $B \leq H \leq G$. If H stabilizes no proper standard flag subspace V_m with $1 \leq m < n$, then the assumed $r = 1$ case gives $H = G$, which is a staircase group. Thus we may assume that H stabilizes some V_m , where $1 \leq m < n$. Let S be the stabilizer of V_m . Then $H \leq S$. With respect to the decomposition

$$V_n(F) = V_m \oplus \langle \mathbf{v}_{m+1}, \dots, \mathbf{v}_n \rangle,$$

the elements of S have block form

$$\begin{pmatrix} * & * \\ 0 & * \end{pmatrix}.$$

Let U be the subgroup with identity diagonal blocks and arbitrary upper-right block, let $K_1 \cong \mathrm{GL}(m, F)$ be the upper-left diagonal block group, and let $K_2 \cong \mathrm{GL}(n - m, F)$ be the lower-right diagonal block group. Then

$$S = U \rtimes (K_1 \times K_2).$$

Also $U \leq B \leq H$. By Exercise 2.10, if a subgroup of a semidirect product contains the normal factor, then it is the semidirect product of that normal factor with its intersection with the complement. Applying this to $U \leq H \leq S$, we get

$$H = U \rtimes Q, \quad Q = H \cap (K_1 \times K_2).$$

Thus the problem has been reduced to understanding the subgroup Q of the block diagonal group $K_1 \times K_2$.

Step 1: Q contains the two block Borels. Let

$$B_1 = B \cap K_1, \quad B_2 = B \cap K_2.$$

These are the standard Borel subgroups inside K_1 and K_2 . Since every block diagonal element of $B_1 \times B_2$ is also an element of B , we have

$$B_1 \times B_2 \leq B \leq H.$$

But $B_1 \times B_2 \leq K_1 \times K_2$, so

$$B_1 \times B_2 \leq H \cap (K_1 \times K_2) = Q.$$

Step 2: apply Goursat's theorem to Q . By Exercise 2.8, Goursat's theorem says that a subgroup of a direct product is controlled by its two projections, its two coordinate kernels, and an isomorphism between the corresponding quotient groups. In our case, define

$$P_1 = \{x \in K_1 \mid (x, y) \in Q \text{ for some } y \in K_2\}, \quad I_1 = \{x \in K_1 \mid (x, 1) \in Q\},$$

and define P_2 and I_2 similarly. Then Exercise 2.8 gives

$$I_i \trianglelefteq P_i \leq K_i \quad (i = 1, 2),$$

together with an isomorphism

$$P_1/I_1 \cong P_2/I_2.$$

Moreover, Q is recovered from these data by

$$Q = \{(x, y) \in P_1 \times P_2 \mid \varphi(xI_1) = yI_2\}.$$

The inclusion $B_1 \times B_2 \leq Q$ gives more than just $B_i \leq P_i$. Indeed, if $b_1 \in B_1$, then $(b_1, 1) \in Q$, so $b_1 \in I_1$. Similarly, if $b_2 \in B_2$, then $(1, b_2) \in Q$, so $b_2 \in I_2$. Hence

$$B_i \leq I_i \leq P_i \leq K_i \quad (i = 1, 2).$$

Step 3: use induction and self-normalizing parabolics. Since $K_1 \cong \mathrm{GL}(m, F)$ and $K_2 \cong \mathrm{GL}(n - m, F)$ have smaller dimension than G , the induction hypothesis applies inside both blocks. Because P_i and I_i contain the standard Borel subgroup B_i , the induction hypothesis tells us that P_i and I_i are staircase groups inside K_i . In particular, I_i is a parabolic subgroup of K_i .

Now $I_i \trianglelefteq P_i$. Therefore every element of P_i normalizes I_i , so

$$P_i \leq N_{K_i}(I_i).$$

By Exercise 5.6, parabolic subgroups are self-normalizing. Applying this inside K_i , we get

$$N_{K_i}(I_i) = I_i.$$

Therefore

$$P_i \leq I_i.$$

Since $I_i \leq P_i$ already, we conclude that

$$P_i = I_i \quad (i = 1, 2).$$

Step 4: the Goursat coupling disappears. Since $P_i = I_i$, both quotient groups in Goursat's theorem are trivial:

$$P_1/I_1 = 1, \quad P_2/I_2 = 1.$$

Thus the condition $\varphi(xI_1) = yI_2$ in the description of Q is automatic. Hence

$$Q = P_1 \times P_2.$$

The induction hypothesis also tells us that P_1 and P_2 are staircase groups inside their respective blocks.

Step 5: combine the two block staircases. Suppose P_1 stabilizes a standard subflag inside V_m , say

$$0 \subset V_{a_1} \subset \cdots \subset V_{a_r} = V_m.$$

For the lower block, write

$$W_j = \langle \mathbf{v}_{m+1}, \dots, \mathbf{v}_{m+j} \rangle \quad (1 \leq j \leq n - m).$$

Thus $W_1 \subset \cdots \subset W_{n-m}$ is the standard flag inside the lower block. Suppose P_2 stabilizes the lower-block subflag

$$0 \subset W_{b_1} \subset \cdots \subset W_{b_s} = W_{n-m}.$$

When these lower-block spaces are placed back inside $V_n(F)$, we must add the first block V_m . Thus

$$V_m \oplus W_{b_j} = V_{m+b_j} \quad (1 \leq j \leq s).$$

Thus the two block flags combine to the standard subflag

$$0 \subset V_{a_1} \subset \cdots \subset V_m \subset V_{m+b_1} \subset \cdots \subset V_n(F).$$

The stabilizer of this combined subflag is exactly the group of block upper triangular matrices whose upper-left diagonal block lies in P_1 and whose lower-right diagonal block lies in P_2 :

$$U \rtimes (P_1 \times P_2).$$

But this group is H , since

$$H = U \rtimes Q = U \rtimes (P_1 \times P_2).$$

Therefore H is a staircase group. This completes the induction step, and hence proves the general case of Theorem 5.4 from the assumed $r = 1$ case. $\square$

6 The Special Linear Group

Definition

- The *special linear group* is the subgroup $\mathrm{SL}(n, F)$ of $\mathrm{GL}(n, F)$ consisting of all matrices having determinant 1.
- In other words, $\mathrm{SL}(n, F)$ is the kernel of the homomorphism $\det: \mathrm{GL}(n, F) \rightarrow F^\times$, and hence $\mathrm{SL}(n, F) \trianglelefteq \mathrm{GL}(n, F)$.

Proposition 6.1. *Let $n \in \mathbb{N}$, and let q be a prime power. Then*

$$|\mathrm{SL}(n, q)| = \prod_{k=1}^{n-1} (q^{n+1} - q^k) = q^{\frac{n(n-1)}{2}} (q^n - 1) \cdots (q^2 - 1).$$

Proof. Let F be the field with q elements. The determinant map

$$\det: \mathrm{GL}(n, F) \rightarrow F^\times$$

is surjective: if $\lambda \in F^\times$, then the diagonal matrix with first diagonal entry λ and all other diagonal entries equal to 1 has determinant λ . Its kernel is precisely $\mathrm{SL}(n, F)$. Hence, by the fundamental theorem on homomorphisms,

$$\mathrm{GL}(n, F) / \mathrm{SL}(n, F) \cong F^\times.$$

In particular,

$$|\mathrm{GL}(n, q) : \mathrm{SL}(n, q)| = |F^\times| = q - 1.$$

Proposition 4.2 gives

$$|\mathrm{GL}(n, q)| = q^{\frac{n(n-1)}{2}} (q^n - 1)(q^{n-1} - 1) \cdots (q - 1).$$

Dividing by the index $q - 1$ cancels the final factor, so

$$|\mathrm{SL}(n, q)| = \frac{|\mathrm{GL}(n, q)|}{q - 1} = q^{\frac{n(n-1)}{2}} (q^n - 1) \cdots (q^2 - 1).$$

Finally,

$$\prod_{k=1}^{n-1} (q^{n+1} - q^k) = \prod_{k=1}^{n-1} q^k (q^{n+1-k} - 1) = q^{1+\cdots+(n-1)} (q^n - 1) \cdots (q^2 - 1),$$

which is the same expression. $\square$

Generation by root subgroups

- We showed in Theorem 4.12 that $\mathrm{GL}(n, F)$ is generated by the root subgroups and the diagonal subgroup. We now establish the analogous result for $\mathrm{SL}(n, F)$.

Proposition 6.2. *$\mathrm{SL}(n, F)$ is generated by the root subgroups X_{ij} .*

Proof. If $n = 1$, then $\mathrm{SL}(1, F) = 1$, so there is nothing to prove. We assume now that $n \geq 2$. Let H be the subgroup of $\mathrm{GL}(n, F)$ generated by all root subgroups X_{ij} . By Lemma 4.6(i), every transvection has determinant 1, so $H \leq \mathrm{SL}(n, F)$. We prove the reverse containment.

Let $A \in \mathrm{SL}(n, F)$. We shall premultiply A by products of transvections until it becomes the identity. Since the inverse of a transvection is again a transvection by Lemma 4.6(iii), this will imply that $A \in H$. First we make A upper triangular. By Lemma 4.7, there is a product b of upper triangular transvections such that, for each $1 \leq i \leq n$, the matrix bA has exactly one row whose entries in the first $i - 1$ columns are zero and whose entry in the i th column is non-zero. In other words, the rows of bA already have the correct pivot pattern for an upper triangular matrix, but the pivot rows may be in the wrong order.

For distinct i and j , put

$$s_{ij} = X_{ji}(1)X_{ij}(-1)X_{ji}(1).$$

On the two rows indexed by i and j , premultiplication by s_{ij} has the same effect as the matrix

$$\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix};$$

that is, it sends row i to the negative of row j and sends row j to row i . Thus s_{ij} swaps the two rows up to a harmless sign. Using such signed row swaps, we can move the row with the first pivot into the first row, the row with the second pivot into the second row, and so on. Therefore there is a product c of transvections such that

$$cbA$$

is upper triangular.

Next we make the diagonal entries equal to 1. Let M be an upper triangular element of $\mathrm{SL}(n, F)$, with diagonal entries $\lambda_1, \dots, \lambda_n$. We use the following two-by-two calculation. If $a, c \in F^\times$ and $b' \in F$, then

$$X_{21}(-1)X_{12}(1 - c^{-1})X_{21}(c) = \begin{pmatrix} c & 1 - c^{-1} \\ 0 & c^{-1} \end{pmatrix},$$

and hence

$$X_{21}(-1)X_{12}(1 - c^{-1})X_{21}(c) \begin{pmatrix} a & b' \\ 0 & c \end{pmatrix} = \begin{pmatrix} ac & cb' + c - 1 \\ 0 & 1 \end{pmatrix},$$

so the only point is that the two diagonal entries a, c have become $ac, 1$. Applying this calculation to the final two diagonal positions of M changes the final two diagonal entries from λ_{n-1}, λ_n to $\lambda_{n-1}\lambda_n, 1$, while keeping the matrix upper triangular. Applying the same calculation next to positions $n-2, n-1$, then to positions $n-3, n-2$, and continuing leftward, we obtain an upper triangular matrix whose diagonal entries are

$$\lambda_1\lambda_2 \cdots \lambda_n, 1, \dots, 1.$$

Since $M \in \mathrm{SL}(n, F)$, the product of the diagonal entries of M is its determinant, and hence $\lambda_1 \cdots \lambda_n = 1$. Thus a product of transvections premultiplies M to an upper unitriangular matrix. Applying this to $M = cbA$, we find a product d of transvections such that $dc bA$ is upper unitriangular.

Finally, by Lemma 4.11, an upper triangular matrix can be premultiplied by a product of transvections to become the diagonal matrix with the same main diagonal. Since $dc bA$ is upper unitriangular, this diagonal matrix is the identity. Hence there is a product e of transvections such that

$$edcbA = I.$$

It follows that

$$A = b^{-1}c^{-1}d^{-1}e^{-1}.$$

Each factor on the right lies in H , so $A \in H$. Therefore $\mathrm{SL}(n, F) \leq H$, and hence $\mathrm{SL}(n, F) = H$. $\square$

Proposition 6.3. *The subgroups X_{ij} are conjugate in $\mathrm{SL}(n, F)$.*

Proof. If $n = 1$, there are no root subgroups, so there is nothing to prove. Let X_{ij} and $X_{i'j'}$ be two root subgroups. Choose a permutation matrix w such that $w(i) = i'$ and $w(j) = j'$. By Lemma 4.6(v), for every $\alpha \in F$ we have

$$wX_{ij}(\alpha)w^{-1} = X_{i'j'}(\alpha).$$

If $\det w = 1$, then $w \in \mathrm{SL}(n, F)$, and hence $wX_{ij}w^{-1} = X_{i'j'}$.

It remains to handle the case $\det w = -1$. Let d be the diagonal matrix whose first diagonal entry is -1 and whose remaining diagonal entries are 1. Then $\det d = -1$, so $\det(dw) = 1$, and hence $dw \in \mathrm{SL}(n, F)$. For each $\alpha \in F$,

$$(dw)X_{ij}(\alpha)(dw)^{-1} = dX_{i'j'}(\alpha)d^{-1}.$$

Conjugating by d can only change the parameter α by a sign, depending on whether one of i' or j' is equal to 1. Thus

$$dX_{i'j'}(\alpha)d^{-1} = X_{i'j'}(\pm\alpha).$$

As α ranges over F , so does $\pm\alpha$. Therefore

$$(dw)X_{ij}(dw)^{-1} = X_{i'j'}.$$

Thus any two root subgroups are conjugate in $\mathrm{SL}(n, F)$. $\square$

Scalar matrices and the center

- Let Z be the subgroup of $\mathrm{GL}(n, F)$ consisting of the multiples of the identity matrix by elements of $F^\times$. The elements of Z are called the (non-zero) *scalar matrices*. We clearly have $Z \cong F^\times$.

Proposition 6.4. *The center of $\mathrm{GL}(n, F)$ is Z , and the center of $\mathrm{SL}(n, F)$ is $Z \cap \mathrm{SL}(n, F)$.*

Proof. If $n = 1$, then $\mathrm{GL}(1, F) = F^\times = Z$ and $\mathrm{SL}(1, F) = 1 = Z \cap \mathrm{SL}(1, F)$, so the result is immediate. We assume now that $n > 1$. Let G denote either $\mathrm{GL}(n, F)$ or $\mathrm{SL}(n, F)$. Since scalar matrices commute with every matrix, we have

$$Z \cap G \leq Z(G).$$

It remains to prove the reverse containment. It is enough to show that if an element of G commutes with every element of $\mathrm{SL}(n, F)$, then it is scalar.

Let $M = (m_{rs})$ be such an element of G . Fix distinct indices i and j . By Lemma 4.6(i), the transvection $X_{ij}(1)$ lies in $\mathrm{SL}(n, F)$, so

$$MX_{ij}(1) = X_{ij}(1)M.$$

Writing $X_{ij}(1) = I + E_{ij}$, this equality is equivalent to

$$ME_{ij} = E_{ij}M.$$

Compare entries. The (i, i) -entry of ME_{ij} is 0, while the (i, i) -entry of $E_{ij}M$ is m_{ji} , so $m_{ji} = 0$. The (i, j) -entry of ME_{ij} is m_{ii} , while the (i, j) -entry of $E_{ij}M$ is m_{jj} , so $m_{ii} = m_{jj}$.

Since i and j were arbitrary distinct indices, every off-diagonal entry of M is zero, and all diagonal entries are equal. Hence $M = m_{11}I$ is a scalar matrix. Thus $M \in Z \cap G$. Therefore $Z(G) = Z \cap G$. Taking $G = \mathrm{GL}(n, F)$ gives $Z(\mathrm{GL}(n, F)) = Z$, and taking $G = \mathrm{SL}(n, F)$ gives $Z(\mathrm{SL}(n, F)) = Z \cap \mathrm{SL}(n, F)$. $\square$

Projective groups

- The group $\mathrm{GL}(n, F)/Z$ is called the *projective general linear group*, and the group $\mathrm{SL}(n, F)/Z \cap \mathrm{SL}(n, F)$ is called the *projective special linear group*.
- We denote these groups by $\mathrm{PGL}(n, F)$ and $\mathrm{PSL}(n, F)$, respectively. (Observe that the first isomorphism theorem implies that $\mathrm{PSL}(n, F) \cong Z \mathrm{SL}(n, F)/Z \leq \mathrm{GL}(n, F)/Z = \mathrm{PGL}(n, F)$.)
- As $Z \cong F^\times$, we have $|\mathrm{PGL}(n, q)| = |\mathrm{SL}(n, q)|$ for any prime power q ; however, to determine $|\mathrm{PSL}(n, q)|$ we must be able to count the number of n th roots of unity in the field of q elements.

Proposition 6.5. *Let q be a prime power. Then*

$$|\mathrm{PSL}(2, q)| = \begin{cases} q^3 - q & \text{if } 2 \mid q, \\ \frac{1}{2}(q^3 - q) & \text{if } 2 \nmid q. \end{cases}$$

Proof. Let F be the field with q elements. By Proposition 6.4,

$$Z(\mathrm{SL}(2, F)) = Z \cap \mathrm{SL}(2, F),$$

and hence

$$|\mathrm{PSL}(2, q)| = \frac{|\mathrm{SL}(2, q)|}{|Z \cap \mathrm{SL}(2, q)|}.$$

By Proposition 6.1, we have

$$|\mathrm{SL}(2, q)| = q(q^2 - 1) = q^3 - q.$$

It remains to count the scalar matrices in $\mathrm{SL}(2, q)$. Such a matrix has the form αI , with $\alpha \in F^\times$, and it lies in $\mathrm{SL}(2, F)$ precisely when

$$\det(\alpha I) = \alpha^2 = 1.$$

Thus α is a root of $X^2 - 1$. If $2 \nmid q$, then F has characteristic not equal to 2, so

$$X^2 - 1 = (X - 1)(X + 1)$$

has the two distinct roots 1 and -1 . Hence $|Z \cap \mathrm{SL}(2, q)| = 2$ in this case. If $2 \mid q$, then F has characteristic 2, so $-1 = 1$ and

$$X^2 - 1 = (X - 1)^2$$

has only the single root 1. Hence $|Z \cap \mathrm{SL}(2, q)| = 1$ in this case. Dividing the order of $\mathrm{SL}(2, q)$ by these two possible central orders gives the stated formula. $\square$

Simplicity of $\text{PSL}(n, F)$

- Our main objective in this section is to show that $\text{PSL}(n, F)$ is simple whenever $n \geq 2$, except when $n = 2$ and $|F| = 2$ or 3 .
- This result dates back to L. E. Dickson, another of the early architects of the mathematical tradition of the University of Chicago, who established it in his 1896 Ph.D. thesis for F a finite field; in 1893 E. H. Moore had established the result for F a finite field and $n = 2$, while in 1870 Camille Jordan had established the result for F a finite field of prime order and n arbitrary.
- We first need two lemmas.

Lemma 6.6. *If $n \geq 2$, then every transvection $X_{ij}(\alpha)$ is a commutator of elements of $\text{SL}(n, F)$, except when $n = 2$ and $|F| = 2$ or 3 .*

Proof. First suppose that $n > 2$. Given distinct i and j , choose k distinct from both of them. By Lemma 4.6(iv),

$$[X_{ik}(\alpha), X_{kj}(1)] = X_{ij}(\alpha).$$

Both $X_{ik}(\alpha)$ and $X_{kj}(1)$ lie in $\text{SL}(n, F)$ by Lemma 4.6(i), so $X_{ij}(\alpha)$ is a commutator of elements of $\text{SL}(n, F)$.

It remains to consider the case $n = 2$. Assume that F is not one of the fields with 2 or 3 elements. Then there is some $\beta \in F^\times$ such that $\beta^2 \neq 1$; otherwise every element of $F^\times$ would be a root of $X^2 - 1$, forcing $|F^\times| \leq 2$. Put

$$d = \begin{pmatrix} \beta & 0 \\ 0 & \beta^{-1} \end{pmatrix}.$$

Then $d \in \text{SL}(2, F)$. For any $\gamma \in F$, a direct calculation gives

$$dX_{12}(\gamma)d^{-1} = X_{12}(\beta^2\gamma),$$

and hence

$$[d, X_{12}(\gamma)] = X_{12}(\beta^2\gamma)X_{12}(-\gamma) = X_{12}((\beta^2 - 1)\gamma).$$

Since $\beta^2 - 1 \neq 0$, choosing $\gamma = \alpha(\beta^2 - 1)^{-1}$ gives

$$X_{12}(\alpha) = [d, X_{12}(\gamma)].$$

Similarly,

$$dX_{21}(\gamma)d^{-1} = X_{21}(\beta^{-2}\gamma),$$

so

$$[d, X_{21}(\gamma)] = X_{21}((\beta^{-2} - 1)\gamma).$$

Since $\beta^{-2} - 1 \neq 0$, choosing $\gamma = \alpha(\beta^{-2} - 1)^{-1}$ gives $X_{21}(\alpha)$ as a commutator of elements of $\text{SL}(2, F)$. This proves the assertion in all non-exceptional cases. $\square$

Consequence

- Observe that Lemma 6.6 and Proposition 6.2 together imply that, except when $n = 2$ and $|F| = 2$ or 3 , $\text{SL}(n, F)$ is its own derived group and is consequently also the derived group of $\text{GL}(n, F)$.

Lemma 6.7. *If $n \geq 2$, then the action of $\text{SL}(n, F)$ on the one-dimensional subspaces of $V_n(F)$ is doubly transitive.*

Proof. Let U_1, U_2, W_1, W_2 be one-dimensional subspaces of $V_n(F)$, with $U_1 \neq U_2$ and $W_1 \neq W_2$. Choose non-zero vectors

$$\mathbf{c}_1 \in U_1, \quad \mathbf{c}_2 \in U_2, \quad \mathbf{d}_1 \in W_1, \quad \mathbf{d}_2 \in W_2.$$

Since U_1 and U_2 are distinct one-dimensional subspaces, the vectors $\mathbf{c}_1$ and $\mathbf{c}_2$ are linearly independent. Similarly, $\mathbf{d}_1$ and $\mathbf{d}_2$ are linearly independent. Extend these pairs to bases

$$\mathbf{c}_1, \dots, \mathbf{c}_n \quad \text{and} \quad \mathbf{d}_1, \dots, \mathbf{d}_n$$

of $V_n(F)$. Let C and D be the matrices whose i th columns are $\mathbf{c}_i$ and $\mathbf{d}_i$, respectively. Then $C, D \in \text{GL}(n, F)$.

Put

$$\varepsilon = \frac{\det D}{\det C},$$

and let E be the diagonal matrix with first diagonal entry ε and all other diagonal entries equal to 1. Then

$$\det(DE^{-1}C^{-1}) = (\det D)\varepsilon^{-1}(\det C)^{-1} = 1,$$

so

$$s = DE^{-1}C^{-1} \in \text{SL}(n, F).$$

Now $C^{-1}\mathbf{c}_1 = \mathbf{v}_1$ and $C^{-1}\mathbf{c}_2 = \mathbf{v}_2$, while $E^{-1}\mathbf{v}_1 = \varepsilon^{-1}\mathbf{v}_1$ and $E^{-1}\mathbf{v}_2 = \mathbf{v}_2$. Hence

$$s\mathbf{c}_1 = \varepsilon^{-1}\mathbf{d}_1, \quad s\mathbf{c}_2 = \mathbf{d}_2.$$

Thus $sU_1 = W_1$ and $sU_2 = W_2$. Therefore $\text{SL}(n, F)$ sends any ordered pair of distinct one-dimensional subspaces to any other such ordered pair, so the action is doubly transitive. $\square$

Theorem 6.8. *If $n \geq 2$, then $\text{PSL}(n, F)$ is simple, except when $n = 2$ and $|F| = 2$ or 3.*

Proof. Assume that we are not in the exceptional cases. Put

$$S = \text{SL}(n, F), \quad L = F\mathbf{v}_1, \quad Z_S = Z \cap S.$$

It is enough to prove that if $N \trianglelefteq S$, then either $N \leq Z_S$ or $N = S$. Indeed, the normal subgroups of

$$\text{PSL}(n, F) = S/Z_S$$

correspond by Correspondence Theorem to the normal subgroups $N \trianglelefteq S$ with $Z_S \leq N \leq S$, via

$$N \mapsto N/Z_S.$$

If every normal subgroup of S is either contained in Z_S or equal to S , then any normal subgroup N/Z_S of S/Z_S has only two possibilities. In the first case, $Z_S \leq N \leq Z_S$, so $N = Z_S$, and hence

$$N/Z_S = Z_S/Z_S,$$

which is the trivial subgroup of S/Z_S . In the second case, $N = S$, and hence

$$N/Z_S = S/Z_S,$$

which is the whole quotient group. Thus proving the stated assertion about normal subgroups of S proves that $\text{PSL}(n, F)$ is simple.

Let P be the stabilizer of the line L under the action of S on the one-dimensional subspaces of $V_n(F)$. By Lemma 6.7 and Proposition 3.8, the subgroup P is maximal in S . Now let

$$K = \left\{ I + \sum_{j=2}^n \alpha_j E_{1j} \mid \alpha_j \in F \right\}.$$

Here E_{1j} denotes the matrix with 1 in the $(1, j)$ -entry and 0 everywhere else. Equivalently, K is the subgroup of upper unitriangular matrices whose only possible non-zero entries away from the main diagonal lie in the first row. Since $E_{1j}E_{1k} = 0$ for all $j, k \geq 2$, multiplication in K just adds the parameters α_j , and hence K is abelian. Also, K is exactly the subgroup of P acting trivially on L and on the quotient space $V_n(F)/L$; this description is invariant under conjugation by elements of P , because P stabilizes L . Hence $K \trianglelefteq P$.

Let $N \trianglelefteq S$. First suppose that $N \leq P$. Then N stabilizes L . If $s \in S$, then $sNs^{-1} = N$, and hence N stabilizes sL . Since the action of S on one-dimensional subspaces is transitive by Lemma 6.7, it follows that N stabilizes every one-dimensional subspace of $V_n(F)$. In particular, every element of N stabilizes each standard line $F\mathbf{v}_i$, so every element of N is diagonal. If such an element has diagonal entries $\lambda_1, \dots, \lambda_n$, then stabilizing each line $F(\mathbf{v}_i + \mathbf{v}_j)$ forces $\lambda_i = \lambda_j$ for all $i \neq j$. Thus every element of N is scalar, and hence $N \leq Z_S$.

Now suppose that $N \not\leq P$. Since $N \trianglelefteq S$, the product PN is a subgroup of S . It properly contains P , so the maximality of P gives

$$PN = S.$$

Let $\eta: S \rightarrow S/N$ be the natural map. Since $K \trianglelefteq P$, we have $\eta(K) \trianglelefteq \eta(P)$. But

$$\eta(P) = PN/N = S/N,$$

so $\eta(K) \trianglelefteq S/N$. By Correspondence Theorem, its inverse image

$$\eta^{-1}(\eta(K)) = KN$$

is normal in S . Since K is generated by the root subgroups $X_{12}, \dots, X_{1n}$, and since KN is normal in S , the subgroup KN contains all S -conjugates of these root subgroups. By Proposition 6.3, all root subgroups are conjugate in S , so KN contains every X_{ij} . Proposition 6.2 then gives

$$KN = S.$$

By First Isomorphism Theorem,

$$S/N = KN/N \cong K/(K \cap N).$$

Since K is abelian, the quotient S/N is abelian. Therefore Proposition 2.6 gives $S' \leq N$. By Lemma 6.6, every transvection $X_{ij}(\alpha)$ is a commutator of elements of S , and hence every transvection lies in N . Since the transvections generate S by Proposition 6.2, we conclude that $N = S$.

Thus every normal subgroup of S is either contained in Z_S or equal to S . Therefore $S/Z_S = \text{PSL}(n, F)$ is simple. $\square$

Exercises

Exercise 6.1. Let B be the standard Borel subgroup of $\text{GL}(n, F)$. Determine all subgroups of $\text{SL}(n, F)$ which contain $B \cap \text{SL}(n, F)$.

Solution. Let $S = \text{SL}(n, F)$ and put $B_S = B \cap S$. We claim that the required subgroups are precisely

$$P \cap S,$$

where P ranges over the staircase groups of $\text{GL}(n, F)$. Indeed, if P is a staircase group, then $B \leq P$, and hence $B_S \leq P \cap S \leq S$.

Conversely, suppose that

$$B_S \leq H \leq S.$$

If $n = 1$, then $S = 1$, so the assertion is immediate. Assume now that $n \geq 2$. We first note that the only non-zero subspaces of $V_n(F)$ stabilized by B_S are the standard flag subspaces $V_1, \dots, V_n$. The proof is the same as Lemma 5.3, with one determinant correction. If

$$y = \alpha_1 \mathbf{v}_1 + \dots + \alpha_k \mathbf{v}_k \in V_k - V_{k-1},$$

then $\alpha_k \neq 0$, and because $n \geq 2$ we can choose an upper triangular matrix of determinant 1 whose k th column is y by adjusting one diagonal entry different from the k th one. Thus the argument of Lemma 5.3 applies with B_S in place of B .

List the standard flag subspaces stabilized by H as

$$V_{a_1}, \dots, V_{a_r}, \quad 1 \leq a_1 < \dots < a_r = n.$$

Let P be the staircase group stabilizing the subflag

$$0 \subset V_{a_1} \subset \dots \subset V_{a_r} = V_n(F).$$

Then $H \leq P \cap S$. It remains to prove the reverse containment.

We use the same block argument as in Theorem 5.4. Fix a block

$$F\mathbf{v}_s \oplus \dots \oplus F\mathbf{v}_t, \quad s = a_{k-1} + 1, \quad t = a_k.$$

If $s = t$, there is nothing to prove for this block. Assume $s < t$. Since there is no H -invariant standard flag subspace strictly between V_{s-1} and V_t , there exists $h \in H$ such that the coefficient of $\mathbf{v}_t$ in $h\mathbf{v}_s$ is non-zero. Write $h = b_1 w b_2$ by Bruhat Decomposition Theorem, with $b_1, b_2 \in B$ and w a permutation matrix. Since $h \in S$, this Bruhat decomposition may be rewritten as

$$h = b'_1 \dot{w} b'_2,$$

where $b'_1, b'_2 \in B_S$ and $\dot{w} \in S$ is a diagonal multiple of w . Indeed, choose a diagonal matrix d with $\det(dw) = 1$ and put $\dot{w} = dw$. If $\delta = \det(b_2)$, choose $t_0 \in T$ with $\det(t_0) = \delta^{-1}$. Then

$$h = (b_1 d^{-1} \dot{w} t_0^{-1} \dot{w}^{-1}) \dot{w} (t_0 b_2),$$

and both outside factors lie in B_S . Since $h, b'_1, b'_2 \in H$, we get $\dot{w} \in H$.

The diagonal factor in $\dot{w}$ does not change the permutation pattern, so the same argument used in Theorem 5.4 gives $w(s) = t$. Put $q = w^{-1}(s)$. Then $s < q \leq t$, so the upper root subgroup X_{sq} lies in $B_S \leq H$. Conjugating by $\dot{w}$ sends this root subgroup onto X_{ts} , up to a non-zero scalar change in the parameter. Hence $X_{ts} \leq H$. Since all upper root subgroups inside the block already lie in B_S , the commutator identities of Lemma 4.6(iv) give every root subgroup X_{ij} with $s \leq i \neq j \leq t$.

Therefore, in each block, H contains all root subgroups supported inside that block. By Proposition 6.2, it contains the corresponding embedded special linear group on each block. It also contains every determinant-one diagonal matrix, since $T \cap S \leq B_S$. Hence H contains the full group $L_P \cap S$, where L_P is the standard Levi complement of P : any element of $L_P \cap S$ is block diagonal of total determinant 1, and after multiplying by a suitable element of $T \cap S$ its blocks all have determinant 1. Finally, the unipotent radical U_P of P lies in B_S , so Proposition 5.2 gives

$$P \cap S = U_P(L_P \cap S) \leq H.$$

Together with $H \leq P \cap S$, this proves $H = P \cap S$. □

Exercise 6.2. Show that $\text{PSL}(2, 2) \cong S_3$ and $\text{PSL}(2, 3) \cong A_4$, and that these groups are not simple.

Solution. By Proposition 6.5,

$$|\text{PSL}(2, 2)| = 2^3 - 2 = 6$$

and

$$|\text{PSL}(2, 3)| = \frac{3^3 - 3}{2} = 12.$$

First let F be the field with two elements. The group $\text{SL}(2, F)$ acts on the three non-zero vectors of $V_2(F)$. If a matrix fixes all three non-zero vectors, then in particular it fixes the basis vectors $\mathbf{v}_1$ and $\mathbf{v}_2$, so it is the identity. Hence this action gives an embedding

$$\text{PSL}(2, 2) = \text{SL}(2, F) \hookrightarrow S_3.$$

Since both groups have order 6, we get $\text{PSL}(2, 2) \cong S_3$.

Now let F be the field with three elements. The group $\text{SL}(2, F)$ acts on the four one-dimensional subspaces of $V_2(F)$, namely

$$F \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad F \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \quad F \begin{pmatrix} 1 \\ 1 \end{pmatrix}, \quad F \begin{pmatrix} 1 \\ -1 \end{pmatrix}.$$

The kernel of this action consists of the scalar matrices in $\text{SL}(2, F)$: indeed, if a matrix fixes the first two lines, it is diagonal, and if it also fixes the line $F(\mathbf{v}_1 + \mathbf{v}_2)$, then its two diagonal entries are equal. Thus the kernel is $\{I, -I\}$, and the action induces an embedding

$$\text{PSL}(2, 3) \hookrightarrow S_4.$$

We show that the image lies in A_4 . By Proposition 6.2, $\text{SL}(2, F)$ is generated by X_{12} and X_{21} . Write the four lines as

$$L_0 = F \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad L_1 = F \begin{pmatrix} 1 \\ 1 \end{pmatrix}, \quad L_{-1} = F \begin{pmatrix} 1 \\ -1 \end{pmatrix}, \quad M = F \begin{pmatrix} 0 \\ 1 \end{pmatrix}.$$

Then $X_{12}(1)$ acts as the cycle

$$(L_1 \ L_{-1} \ M)$$

and $X_{21}(1)$ acts as the cycle

$$(L_0 L_1 L_{-1}).$$

These are even permutations, and their powers give the images of the whole root subgroups X_{12} and X_{21} . Therefore the image of $\text{PSL}(2, 3)$ in S_4 is contained in A_4 . Since this image has order 12 and $|A_4| = 12$, it follows that $\text{PSL}(2, 3) \cong A_4$.

Finally, these groups are not simple: $A_3 \trianglelefteq S_3$ is a non-trivial proper normal subgroup, and the Klein four-subgroup is a non-trivial proper normal subgroup of A_4 . $\square$

Exercise 6.3. Show that $\text{PSL}(2, 4)$ and $\text{PSL}(2, 5)$ are both isomorphic with A_5 .

Solution. By Proposition 6.5,

$$|\text{PSL}(2, 4)| = 4^3 - 4 = 60$$

and

$$|\text{PSL}(2, 5)| = \frac{5^3 - 5}{2} = 60.$$

First let F be the field with four elements. The group $\text{SL}(2, F)$ acts on the five one-dimensional subspaces of $V_2(F)$. The kernel of this action consists of the scalar matrices in $\text{SL}(2, F)$, and since F has characteristic 2, the only scalar matrix in $\text{SL}(2, F)$ is I . Thus this action gives an embedding

$$\text{PSL}(2, 4) = \text{SL}(2, F) \hookrightarrow S_5.$$

We show that the image lies in A_5 . Write the five lines as

$$L_a = F \begin{pmatrix} 1 \\ a \end{pmatrix} \quad (a \in F), \quad M = F \begin{pmatrix} 0 \\ 1 \end{pmatrix}.$$

For $\alpha \neq 0$, the transvection $X_{21}(\alpha)$ fixes M and sends

$$L_a \mapsto L_{a+\alpha}.$$

Since the additive group of F has characteristic 2, the translation $a \mapsto a + \alpha$ on the four elements of F is a product of two disjoint transpositions. Hence $X_{21}(\alpha)$ acts as an even permutation on the five lines. Similarly, $X_{12}(\alpha)$ fixes L_0 and, for $\alpha \neq 0$, fixes no other line: if

$$X_{12}(\alpha) \begin{pmatrix} 1 \\ a \end{pmatrix} = \begin{pmatrix} 1 + \alpha a \\ a \end{pmatrix}$$

spans the same line as $\begin{pmatrix} 1 \\ a \end{pmatrix}$ and $a \neq 0$, then the second coordinate forces the scalar to be 1, and hence $\alpha a = 0$, a contradiction. Also M is sent to $L_{\alpha^{-1}}$. Thus $X_{12}(\alpha)$ is an involution fixing exactly one of the five lines, so it is again a product of two disjoint transpositions. By Proposition 6.2, the image of $\text{SL}(2, F)$ in S_5 is generated by these even permutations, and hence lies in A_5 . Since the image has order $60 = |A_5|$, we conclude that

$$\text{PSL}(2, 4) \cong A_5.$$

Now let F be the field with five elements, and put $G = \text{PSL}(2, 5)$. We shall find a subgroup of G of index 5. In $\text{SL}(2, F)$, let

$$C = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad D = \begin{pmatrix} 2 & 0 \\ 0 & -2 \end{pmatrix}, \quad R = \begin{pmatrix} 1 & 1 \\ 2 & 3 \end{pmatrix}.$$

Let c, d, r be their images in G . Since

$$C^2 = D^2 = (CD)^2 = -I$$

and $CD = -DC$, the subgroup

$$V = \{1, c, d, cd\}$$

is a Klein four-subgroup of G . Also $R^3 = I$, so r has order 3. A direct calculation, modulo the scalar subgroup $\{I, -I\}$, gives

$$rcr^{-1} = cd, \quad r(cd)r^{-1} = d, \quad rdr^{-1} = c.$$

Thus r normalizes V and cyclically permutes its three non-identity elements. Therefore

$$H = V\langle r \rangle$$

is a subgroup of G of order 12. Since $|G| = 60$, the index of H in G is 5. Let G act on the five left cosets of H . This gives a homomorphism

$$G \rightarrow S_5.$$

The kernel is normal in G . By Theorem 6.8, the group $G = \text{PSL}(2, 5)$ is simple, since this is not one of the exceptional cases. The action is not trivial because H is a proper subgroup, so the kernel cannot be all of G . Hence the kernel is trivial, and G embeds in S_5 . Finally, composing this embedding with the sign homomorphism $S_5 \rightarrow \{\pm 1\}$ gives a homomorphism from the simple group G to a group of order 2. This homomorphism must be trivial, since a non-trivial one would have a non-trivial proper kernel. Therefore the image of G lies in A_5 . Since both G and A_5 have order 60, the image is all of A_5 , and hence

$$\text{PSL}(2, 5) \cong A_5.$$

□

Exercise 6.4. Show that $\text{PSL}(2, 7) \cong \text{GL}(3, 2)$.

Solution. Let F be the field with seven elements, and put $G = \text{PSL}(2, 7)$. By Proposition 6.5,

$$|G| = \frac{7^3 - 7}{2} = 168.$$

Also

$$|\text{GL}(3, 2)| = (2^3 - 1)(2^3 - 2)(2^3 - 2^2) = 7 \cdot 6 \cdot 4 = 168.$$

We shall construct the same faithful permutation group of degree 7 from G and from $\text{GL}(3, 2)$. First we construct a subgroup of G of index 7. In $\text{SL}(2, F)$, let

$$C = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad D = \begin{pmatrix} 2 & 3 \\ 3 & -2 \end{pmatrix}, \quad R = \begin{pmatrix} 0 & 2 \\ 3 & 1 \end{pmatrix}, \quad E = \begin{pmatrix} 1 & -1 \\ 2 & -1 \end{pmatrix}.$$

Let c, d, r, e be their images in G . A direct calculation gives

$$C^2 = D^2 = (CD)^2 = -I, \quad CD = -DC.$$

Thus

$$V = \{1, c, d, cd\}$$

is a Klein four-subgroup of G . Also $R^3 = -I$ and $E^2 = -I$, so r has order 3 and e has order 2 in G . Conjugation gives

$$rcr^{-1} = d, \quad rdr^{-1} = cd, \quad r(cd)r^{-1} = c,$$

and

$$ece^{-1} = cd, \quad ede^{-1} = d, \quad e(cd)e^{-1} = c.$$

Moreover $ere^{-1} = r^{-1}$, so $\langle r, e \rangle \cong S_3$. Hence r and e act on the three non-identity elements of V as a 3-cycle and a transposition. This action is faithful, while every element of V centralizes V , so $V \cap \langle r, e \rangle = 1$. Therefore the subgroup

$$H = V\langle r, e \rangle$$

has order $4 \cdot 6 = 24$. Since $|G| = 168$, the subgroup H has index 7 in G .

Let G act on the seven left cosets of H . The kernel is normal in G . By Theorem 6.8, the group $G = \text{PSL}(2, 7)$ is simple. The action is not trivial because H is proper, so the kernel is not all of G . Hence the kernel is trivial, and this coset action embeds G in S_7 .

Now take

$$S = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, \quad T = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}.$$

The images of S and T generate G : indeed, $T = X_{12}(1)$, and conjugating powers of T by S gives the lower root subgroup X_{21} , so Proposition 6.2 applies. We now make the coset calculation explicit. Choose

the following representatives g_i for the seven left cosets of H in G , where each displayed matrix denotes its image in G :

$$\begin{array}{c|c} i & g_i \\ \hline 1 & \begin{pmatrix} 2 & -3 \\ 2 & 1 \end{pmatrix} \\ 2 & \begin{pmatrix} 1 & 0 \\ 2 & 1 \end{pmatrix} \\ 3 & \begin{pmatrix} 2 & 3 \\ -2 & 1 \end{pmatrix} \\ 4 & \begin{pmatrix} 3 & 0 \\ 2 & -2 \end{pmatrix} \\ 5 & \begin{pmatrix} 3 & -2 \\ 0 & -2 \end{pmatrix} \\ 6 & \begin{pmatrix} 3 & 2 \\ 0 & -2 \end{pmatrix} \\ 7 & \begin{pmatrix} 1 & -1 \\ 2 & -1 \end{pmatrix}. \end{array}$$

For $X \in \{S, T\}$, the equality $Xg_iH = g_jH$ is equivalent to

$$g_j^{-1}Xg_i \in H.$$

Checking this membership for the displayed representatives gives

i	1	2	3	4	5	6	7
Sg_iH	g_1H	g_5H	g_4H	g_3H	g_2H	g_6H	g_7H
Tg_iH	g_3H	g_5H	g_6H	g_7H	g_4H	g_2H	g_1H

For example,

$$g_5^{-1}Sg_2 = \begin{pmatrix} 1 & -2 \\ -3 & 0 \end{pmatrix} = R^{-1} \in H,$$

so $Sg_2H = g_5H$. Reading the two rows of the table as permutations of the labels $1, \dots, 7$, we get

$$S \mapsto (2\ 5)(3\ 4), \quad T \mapsto (1\ 3\ 6\ 2\ 5\ 4\ 7).$$

Now consider the natural action of $\text{GL}(3, 2)$ on the seven non-zero vectors of $\mathbb{F}_2^3$. Label these vectors by

label	1	2	3	4	5	6	7
vector	e_1	e_2	$e_1 + e_2$	e_3	$e_1 + e_3$	$e_2 + e_3$	$e_1 + e_2 + e_3$

With respect to the basis e_1, e_2, e_3 , the matrices

$$A = \begin{pmatrix} 1 & 1 & 1 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}, \quad B = \begin{pmatrix} 1 & 1 & 1 \\ 1 & 0 & 1 \\ 0 & 1 & 1 \end{pmatrix}$$

are invertible over $\mathbb{F}_2$. A direct calculation on the seven labelled vectors gives

$$A \mapsto (2\ 5)(3\ 4), \quad B \mapsto (1\ 3\ 6\ 2\ 5\ 4\ 7).$$

Thus the two generators of the image of G in S_7 lie in the image of the natural embedding

$$\text{GL}(3, 2) \hookrightarrow S_7.$$

Since S and T generate G , the whole image of G lies in the image of $\text{GL}(3, 2)$. Both images have order 168: the coset action of G is faithful, and the natural action of $\text{GL}(3, 2)$ on the non-zero vectors is faithful. Therefore the two images in S_7 are equal, and hence

$$\text{PSL}(2, 7) \cong \text{GL}(3, 2).$$

□

Exercise 6.5. Show that $\text{PSL}(2, 9) \cong A_6$.

Solution. Let $F = \mathbb{F}_9 = \mathbb{F}_3(i)$, where $i^2 = -1$, and put $G = \text{PSL}(2, 9)$. By Proposition 6.5,

$$|G| = \frac{9^3 - 9}{2} = 360.$$

We first construct a subgroup of G of order 60. In $\text{SL}(2, F)$, let

$$X = \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix}, \quad Y = \begin{pmatrix} i & i \\ 1 & 1-i \end{pmatrix}.$$

Both matrices have determinant 1. Let x and y be their images in G . A direct calculation gives

$$X^2 = -I, \quad Y^3 = -I, \quad (XY)^5 = -I.$$

Moreover X and Y are not scalar, Y^2 is not scalar, and none of $XY, (XY)^2, (XY)^3, (XY)^4$ is scalar. Hence x has order 2, y has order 3, and xy has order 5 in G .

We use the standard presentation

$$A_5 \cong \langle a, b \mid a^2 = 1, b^3 = 1, (ab)^5 = 1 \rangle.$$

Therefore there is a homomorphism

$$A_5 \rightarrow \langle x, y \rangle$$

sending a to x and b to y . This homomorphism is surjective by definition of $\langle x, y \rangle$. Its image is non-trivial, since x has order 2. Also A_5 is simple by Exercise 6.3 and Theorem 6.8. Hence the kernel is not all of A_5 , so the kernel is trivial. Thus

$$H = \langle x, y \rangle \cong A_5,$$

and in particular $|H| = 60$. Since $|G| = 360$, the subgroup H has index 6 in G .

Let G act on the six left cosets of H . This gives a homomorphism

$$G \rightarrow S_6.$$

The kernel is normal in G . By Theorem 6.8, the group $G = \text{PSL}(2, 9)$ is simple. The action is not trivial because H is a proper subgroup, so the kernel cannot be all of G . Hence the kernel is trivial, and G embeds in S_6 . Composing with the sign homomorphism $S_6 \rightarrow \{\pm 1\}$ gives a homomorphism from the simple group G to a group of order 2. This homomorphism must be trivial, since a non-trivial one would have a non-trivial proper kernel. Therefore the image of G lies in A_6 . Since both groups have order 360, the image is all of A_6 , and hence

$$\text{PSL}(2, 9) \cong A_6.$$

□

Exercise 6.6. By Exercise 6.3 and Theorem 6.8, the group A_5 is simple. Attempt to prove this fact directly by mimicking the proof of Theorem 6.8, with A_5 in place of S , A_4 in place of P , and the Klein four-group in place of K .

Solution. Put $S = A_5$, acting naturally on $\{1, 2, 3, 4, 5\}$, and let

$$P = \{s \in S \mid s(5) = 5\}.$$

Then $P \cong A_4$, since its elements are exactly the even permutations of $\{1, 2, 3, 4\}$. In particular $|P| = 12$, so P has index 5 in S and is maximal. Let

$$K = \{1, (12)(34), (13)(24), (14)(23)\}.$$

Then K is the Klein four-subgroup of P . It is normal in P , because it is the identity together with all three double transpositions in A_4 , and conjugation inside A_4 preserves this set.

Let $N \trianglelefteq S$. First suppose that $N \leq P$. Since N is normal in S , for every $s \in S$ we have

$$N = sNs^{-1} \leq sPs^{-1}.$$

The subgroup sPs^{-1} is the stabilizer of the point $s(5)$. Since S acts transitively on $\{1, 2, 3, 4, 5\}$, it follows that N is contained in the stabilizer of every point. Thus every element of N fixes every point, and hence $N = 1$.

Now suppose that $N \not\leq P$. Since P is maximal, the subgroup PN properly contains P , and therefore

$$PN = S.$$

Let $\eta: S \rightarrow S/N$ be the natural map. Since $K \trianglelefteq P$, we have $\eta(K) \trianglelefteq \eta(P)$. But

$$\eta(P) = PN/N = S/N,$$

so $\eta(K) \trianglelefteq S/N$. By Correspondence Theorem, its inverse image

$$\eta^{-1}(\eta(K)) = KN$$

is normal in S . Since KN contains K and is normal in S , it contains every S -conjugate of every element of K . The non-identity elements of K are double transpositions, and their S -conjugates are all the double transpositions in A_5 .

The double transpositions generate A_5 . Indeed, if $(a\ b\ c)$ is a 3-cycle and d, e are the remaining two letters, then

$$(a\ b\ c) = (b\ c)(d\ e)(a\ c)(d\ e),$$

which is a product of two double transpositions. Since A_5 is generated by its 3-cycles, it follows that

$$KN = S.$$

Hence

$$S/N = KN/N \cong K/(K \cap N)$$

by First Isomorphism Theorem. Since K is abelian, the quotient S/N is abelian.

It remains to observe that S has no non-trivial abelian quotient. Let $\varphi: S \rightarrow A$ be a homomorphism to an abelian group. If $\tau = (a\ b\ c)$ is a 3-cycle and d, e are the remaining two letters, then conjugation by the double transposition $(a\ b)(d\ e) \in S$ sends τ to τ^{-1} . Since A is abelian, conjugate elements have the same image, so

$$\varphi(\tau) = \varphi(\tau^{-1}) = \varphi(\tau)^{-1}.$$

Also $\tau^3 = 1$, so $\varphi(\tau)^3 = 1$. Hence $\varphi(\tau) = 1$. Since the 3-cycles generate S , the homomorphism φ is trivial. Applying this to the natural map $S \rightarrow S/N$, we get $S/N = 1$, and hence $N = S$.

Thus every normal subgroup of $S = A_5$ is either 1 or S , so A_5 is simple. $\square$

Further Exercises

Let F be a field, and let $V = F^2$. We define an equivalence relation on $V - \{0\}$ by $\mathbf{v} \sim \mathbf{w}$ iff $\mathbf{v} = \alpha\mathbf{w}$ for some $\alpha \in F^\times$. The set of equivalence classes under this relation is called the *one-dimensional projective space* (or *projective line*) over F and is denoted by $\mathbb{P}^1(F)$. We write elements of V as column vectors, and the element of $\mathbb{P}^1(F)$ that is the equivalence class of $\begin{pmatrix} x \\ y \end{pmatrix}$ shall be written as $\left[\begin{smallmatrix} x \\ y \end{smallmatrix} \right]$.

Exercise 6.7. Show that there is a well-defined action of $\text{PSL}(2, F)$ on $\mathbb{P}^1(F)$ given by

$$\overline{\begin{pmatrix} a & b \\ c & d \end{pmatrix}} \left[\begin{smallmatrix} x \\ y \end{smallmatrix} \right] = \left[\begin{smallmatrix} ax + by \\ cx + dy \end{smallmatrix} \right]$$

and that this action is doubly transitive. (Here $\overline{\begin{pmatrix} a & b \\ c & d \end{pmatrix}}$ denotes the image in $\text{PSL}(2, F)$ of $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{SL}(2, F)$.)

Solution. Let $A \in \text{SL}(2, F)$ and let $0 \neq \mathbf{v} \in V$. Since A is invertible, $A\mathbf{v} \neq 0$, so the displayed formula sends a point of $\mathbb{P}^1(F)$ to another point of $\mathbb{P}^1(F)$. We first check that this point is independent of the chosen vector representative. If $\mathbf{v}' = \lambda\mathbf{v}$ for some $\lambda \in F^\times$, then

$$A\mathbf{v}' = \lambda A\mathbf{v},$$

so $A\mathbf{v}'$ and $A\mathbf{v}$ span the same one-dimensional subspace. Hence the formula is well-defined on projective points.

Next we check that the formula is well-defined on $\text{PSL}(2, F)$. If A' represents the same element of $\text{PSL}(2, F)$ as A , then

$$A' = A(\mu I)$$

for some scalar matrix $\mu I \in Z \cap \text{SL}(2, F)$. In particular $\mu \in F^\times$, and hence

$$A'\mathbf{v} = \mu A\mathbf{v}.$$

Thus $A'\mathbf{v}$ and $A\mathbf{v}$ define the same point of $\mathbb{P}^1(F)$. Therefore the formula depends only on the image $\bar{A} \in \text{PSL}(2, F)$, not on the particular representative $A \in \text{SL}(2, F)$.

The identity element acts as the identity, and for $A, B \in \text{SL}(2, F)$ we have

$$A(B\mathbf{v}) = (AB)\mathbf{v}.$$

Passing to images in $\text{PSL}(2, F)$, this proves that the formula defines an action of $\text{PSL}(2, F)$ on $\mathbb{P}^1(F)$. It remains to show that the action is doubly transitive. Let $p_1, p_2, q_1, q_2 \in \mathbb{P}^1(F)$ with $p_1 \neq p_2$ and $q_1 \neq q_2$. Choose non-zero vectors $\mathbf{u}_1, \mathbf{u}_2, \mathbf{w}_1, \mathbf{w}_2 \in V$ such that

$$p_i = [\mathbf{u}_i], \quad q_i = [\mathbf{w}_i] \quad (i = 1, 2).$$

Since $p_1 \neq p_2$, the vectors $\mathbf{u}_1, \mathbf{u}_2$ are linearly independent; similarly, $\mathbf{w}_1, \mathbf{w}_2$ are linearly independent. Let $\det(\mathbf{u}_1, \mathbf{u}_2)$ denote the determinant of the matrix with columns $\mathbf{u}_1, \mathbf{u}_2$, and similarly for $\mathbf{w}_1, \mathbf{w}_2$. Put

$$\delta = \frac{\det(\mathbf{u}_1, \mathbf{u}_2)}{\det(\mathbf{w}_1, \mathbf{w}_2)} \in F^\times.$$

There is a unique linear map $A: V \rightarrow V$ satisfying

$$A\mathbf{u}_1 = \delta\mathbf{w}_1, \quad A\mathbf{u}_2 = \mathbf{w}_2,$$

because $\mathbf{u}_1, \mathbf{u}_2$ form a basis of V . Its determinant is

$$\det A = \frac{\det(\delta\mathbf{w}_1, \mathbf{w}_2)}{\det(\mathbf{u}_1, \mathbf{u}_2)} = \frac{\delta \det(\mathbf{w}_1, \mathbf{w}_2)}{\det(\mathbf{u}_1, \mathbf{u}_2)} = 1.$$

Thus $A \in \text{SL}(2, F)$. Since scalar multiples represent the same projective point, the image $\bar{A} \in \text{PSL}(2, F)$ sends

$$p_1 \mapsto q_1, \quad p_2 \mapsto q_2.$$

Therefore the action is doubly transitive. $\square$

Exercise 6.8. (cont.) Determine the proper definition of n -dimensional projective space $\mathbb{P}^n(F)$ for arbitrary n , and show that $\text{PSL}(n+1, F)$ acts doubly transitively on $\mathbb{P}^n(F)$.

Solution. Let $V = F^{n+1}$. We define

$$\mathbb{P}^n(F) = (V - \{0\}) / \sim,$$

where

$$\mathbf{v} \sim \mathbf{w} \iff \mathbf{w} = \lambda\mathbf{v} \text{ for some } \lambda \in F^\times.$$

Equivalently, $\mathbb{P}^n(F)$ is the set of one-dimensional subspaces of V . We write the equivalence class of a non-zero vector $\mathbf{v} \in V$ as $[\mathbf{v}]$.

The action is defined by

$$\bar{A}[\mathbf{v}] = [A\mathbf{v}], \quad A \in \text{SL}(n+1, F).$$

This is well-defined for the same reasons as in Exercise 6.7. If $\mathbf{v}' = \lambda\mathbf{v}$ for some $\lambda \in F^\times$, then

$$A\mathbf{v}' = \lambda A\mathbf{v},$$

so $A\mathbf{v}'$ and $A\mathbf{v}$ define the same projective point. Also, if A' represents the same element of $\text{PSL}(n+1, F)$ as A , then $A' = A(\mu I)$ for some scalar matrix $\mu I \in Z \cap \text{SL}(n+1, F)$, and hence

$$A'\mathbf{v} = \mu A\mathbf{v},$$

which again defines the same projective point. Finally, the identity acts as the identity, and

$$A(B\mathbf{v}) = (AB)\mathbf{v}$$

for $A, B \in \text{SL}(n+1, F)$, so the displayed formula defines an action of $\text{PSL}(n+1, F)$ on $\mathbb{P}^n(F)$.

We now prove double transitivity. If $n = 0$, then $\mathbb{P}^0(F)$ has one point, so there are no ordered pairs of distinct points to consider. Assume $n \geq 1$. Let

$$p_1, p_2, q_1, q_2 \in \mathbb{P}^n(F), \quad p_1 \neq p_2, \quad q_1 \neq q_2.$$

Choose non-zero vectors $\mathbf{u}_1, \mathbf{u}_2, \mathbf{w}_1, \mathbf{w}_2 \in V$ such that

$$p_i = [\mathbf{u}_i], \quad q_i = [\mathbf{w}_i] \quad (i = 1, 2).$$

Since $p_1 \neq p_2$, the vectors $\mathbf{u}_1, \mathbf{u}_2$ are linearly independent. Similarly, $\mathbf{w}_1, \mathbf{w}_2$ are linearly independent. Extend these pairs to bases

$$\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3, \dots, \mathbf{u}_{n+1} \quad \text{and} \quad \mathbf{w}_1, \mathbf{w}_2, \mathbf{w}_3, \dots, \mathbf{w}_{n+1}$$

of V . Put

$$\delta = \frac{\det(\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_{n+1})}{\det(\mathbf{w}_1, \mathbf{w}_2, \dots, \mathbf{w}_{n+1})} \in F^\times.$$

There is a unique linear map $A: V \rightarrow V$ satisfying

$$A\mathbf{u}_1 = \delta\mathbf{w}_1, \quad A\mathbf{u}_i = \mathbf{w}_i \quad (2 \leq i \leq n+1).$$

Its determinant is

$$\begin{aligned} \det A &= \frac{\det(\delta\mathbf{w}_1, \mathbf{w}_2, \dots, \mathbf{w}_{n+1})}{\det(\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_{n+1})} \\ &= \frac{\delta \det(\mathbf{w}_1, \mathbf{w}_2, \dots, \mathbf{w}_{n+1})}{\det(\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_{n+1})} = 1. \end{aligned}$$

Thus $A \in \mathrm{SL}(n+1, F)$. Since scalar multiples represent the same projective point, the image $\bar{A} \in \mathrm{PSL}(n+1, F)$ sends

$$p_1 \mapsto q_1, \quad p_2 \mapsto q_2.$$

Therefore the action is doubly transitive. $\square$

Exercise 6.9. Let F be the field of 7 elements. For $0 \leq i \leq 6$, let i represent $\begin{bmatrix} 1 \\ i \end{bmatrix} \in \mathbb{P}^1(F)$; denote the remaining element $\begin{bmatrix} 0 \\ 1 \end{bmatrix}$ of $\mathbb{P}^1(F)$ by ∞ . View S_8 as being the group of permutations of $\{\infty, 0, 1, 2, 3, 4, 5, 6\}$, and let $\varphi: \mathrm{PSL}(2, 7) \rightarrow S_8$ be the monomorphism arising (as in Proposition 3.1) from the action of $\mathrm{PSL}(2, 7)$ on $\mathbb{P}^1(F)$. Show that the image of φ is the subgroup of A_8 generated by $(0 \ 1 \ 2 \ 3 \ 4 \ 5 \ 6)$, $(1 \ 4 \ 2)(3 \ 5 \ 6)$, and $(\infty \ 0)(1 \ 3)(2 \ 5)(4 \ 6)$. (HINT: Show first that $\mathrm{PSL}(2, 7)$ is generated by the images under the natural map of the elements $\begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}$, $\begin{pmatrix} 4 & 0 \\ 0 & 2 \end{pmatrix}$, and $\begin{pmatrix} 0 & 3 \\ 2 & 0 \end{pmatrix}$ of $\mathrm{SL}(2, 7)$.) For the relevance of this exercise, see Exercises 7.13–7.17.

Solution. Let

$$A = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}, \quad B = \begin{pmatrix} 4 & 0 \\ 0 & 2 \end{pmatrix}, \quad C = \begin{pmatrix} 0 & 3 \\ 2 & 0 \end{pmatrix}$$

over F . These are all in $\mathrm{SL}(2, 7)$, since

$$\det A = 1, \quad \det B = 8 = 1, \quad \det C = -6 = 1$$

in F . Let $\bar{A}, \bar{B}, \bar{C}$ be their images in $\mathrm{PSL}(2, 7)$. We first show that these three elements generate $\mathrm{PSL}(2, 7)$. The element $A = X_{21}(1)$, so its powers give every element of the lower root subgroup X_{21} . Also a direct calculation gives

$$CX_{21}(\alpha)C^{-1} = X_{12}(5\alpha) \quad (\alpha \in F).$$

Thus the subgroup generated by $\bar{A}, \bar{B}, \bar{C}$ contains the images of both root subgroups X_{21} and X_{12} . By Proposition 6.2, the root subgroups generate $\mathrm{SL}(2, 7)$, so their images generate $\mathrm{PSL}(2, 7)$.

It remains to compute the corresponding permutations of $\mathbb{P}^1(F)$. For $0 \leq i \leq 6$, we identify i with $\begin{bmatrix} 1 \\ i \end{bmatrix}$ and identify ∞ with $\begin{bmatrix} 0 \\ 1 \end{bmatrix}$. First,

$$A\begin{pmatrix} 1 \\ i \end{pmatrix} = \begin{pmatrix} 1 \\ i+1 \end{pmatrix}, \quad A\begin{pmatrix} 0 \\ 1 \end{pmatrix} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}.$$

Hence $\varphi(\bar{A})$ fixes ∞ and sends i to $i+1$ modulo 7, so

$$\varphi(\bar{A}) = (0 \ 1 \ 2 \ 3 \ 4 \ 5 \ 6).$$

Next,

$$B \begin{pmatrix} 1 \\ i \end{pmatrix} = \begin{pmatrix} 4 \\ 2i \end{pmatrix} = \begin{pmatrix} 1 \\ 4i \end{pmatrix}, \quad B \begin{pmatrix} 0 \\ 1 \end{pmatrix} = \begin{pmatrix} 0 \\ 2 \end{pmatrix}.$$

Thus $\varphi(\overline{B})$ fixes 0 and ∞ and sends i to $4i$ on $F^\times$. Since multiplication by 4 on $F^\times$ has cycles

$$1 \mapsto 4 \mapsto 2 \mapsto 1, \quad 3 \mapsto 5 \mapsto 6 \mapsto 3,$$

we get

$$\varphi(\overline{B}) = (1 \ 4 \ 2)(3 \ 5 \ 6).$$

Finally,

$$C \begin{pmatrix} 0 \\ 1 \end{pmatrix} = \begin{pmatrix} 3 \\ 0 \end{pmatrix}, \quad C \begin{pmatrix} 1 \\ 0 \end{pmatrix} = \begin{pmatrix} 0 \\ 2 \end{pmatrix}.$$

So $\varphi(\overline{C})$ interchanges ∞ and 0. If $i \neq 0$, then

$$C \begin{pmatrix} 1 \\ i \end{pmatrix} = \begin{pmatrix} 3i \\ 2 \end{pmatrix} = \begin{pmatrix} 1 \\ 3i^{-1} \end{pmatrix},$$

because $(3i)^{-1}2 = 3i^{-1}$ in F . Therefore

$$1 \leftrightarrow 3, \quad 2 \leftrightarrow 5, \quad 4 \leftrightarrow 6,$$

and hence

$$\varphi(\overline{C}) = (\infty \ 0)(1 \ 3)(2 \ 5)(4 \ 6).$$

Since $\overline{A}, \overline{B}, \overline{C}$ generate $\mathrm{PSL}(2, 7)$, their images generate $\mathrm{im} \varphi$. Thus

$$\mathrm{im} \varphi = \langle (0 \ 1 \ 2 \ 3 \ 4 \ 5 \ 6), (1 \ 4 \ 2)(3 \ 5 \ 6), (\infty \ 0)(1 \ 3)(2 \ 5)(4 \ 6) \rangle.$$

Each of these generators is an even permutation, so this subgroup lies in A_8 . □

Recall the definition of a BN -pair from the further exercises to Section 4.

Exercise 6.10. Let $G = \mathrm{GL}(n, F)$. Let B be the standard Borel subgroup of G , let N be the subgroup of G of monomial matrices, and let $T = B \cap N$ be the subgroup of diagonal matrices. Let $B_0 = B \cap \mathrm{SL}(n, F)$, let $N_0 = N \cap \mathrm{SL}(n, F)$, and let $T_0 = T \cap \mathrm{SL}(n, F) = B_0 \cap N_0$. Show that B_0 and N_0 form a BN -pair of $\mathrm{SL}(n, F)$, with the associated Weyl group $W_0 = N_0/T_0$ being isomorphic with S_n .

Solution. Put

$$G_0 = \mathrm{SL}(n, F).$$

We verify the defining conditions for a BN -pair. Thus we must show that $G_0 = \langle B_0, N_0 \rangle$, that $T_0 = B_0 \cap N_0$ is normal in N_0 , that $W_0 = N_0/T_0$ is generated by a finite set of involutions, and that the two double-coset conditions hold.

We first record a determinant-one version of the usual Bruhat double cosets. Suppose that $n_0 \in N_0$ and that

$$x \in G_0 \cap B n_0 B.$$

Write $x = b_1 n_0 b_2$ with $b_1, b_2 \in B$. Since x and n_0 have determinant 1, we have

$$\det(b_1) \det(b_2) = 1.$$

Choose a diagonal matrix $d \in T$ with $\det(d) = \det(b_1)^{-1}$. Then

$$x = (b_1 d) n_0 (n_0^{-1} d^{-1} n_0 b_2).$$

The left factor $b_1 d$ lies in B_0 . Also $n_0^{-1} d^{-1} n_0 \in T$, because monomial matrices normalize the diagonal subgroup. Hence the right factor lies in B . Its determinant is

$$\det(d)^{-1} \det(b_2) = \det(b_1) \det(b_2) = 1,$$

so it lies in B_0 . Therefore

$$G_0 \cap B n_0 B \subseteq B_0 n_0 B_0.$$

The reverse inclusion is immediate from $B_0, n_0 \leq G_0$ and $B_0 \leq B$, so

$$G_0 \cap Bn_0B = B_0n_0B_0 \quad (n_0 \in N_0).$$

We now prove that G_0 is generated by B_0 and N_0 . Let $g \in G_0$. By the Bruhat Decomposition Theorem in $\mathrm{GL}(n, F)$, we can write

$$g = b_1 p b_2,$$

where $b_1, b_2 \in B$ and p is a permutation matrix. Choose a diagonal matrix $e \in T$ with $\det(e) = \det(p)^{-1}$. Then

$$n_0 = ep$$

lies in N_0 , and $g = (b_1 e^{-1}) n_0 b_2 \in Bn_0B$. Since $g \in G_0$, the determinant-one double-coset observation gives $g \in B_0 n_0 B_0$. Thus

$$G_0 = B_0 N_0 B_0,$$

and in particular $G_0 = \langle B_0, N_0 \rangle$.

Next,

$$B_0 \cap N_0 = (B \cap \mathrm{SL}(n, F)) \cap (N \cap \mathrm{SL}(n, F)) = T \cap \mathrm{SL}(n, F) = T_0.$$

Also $T_0 \trianglelefteq N_0$, because conjugating a diagonal matrix by a monomial matrix gives another diagonal matrix, and determinant 1 is preserved under conjugation.

We identify the Weyl group. Define

$$\pi: N_0 \rightarrow S_n$$

by sending a monomial matrix to its underlying permutation of the standard basis lines. Equivalently, if

$$m \mathbf{v}_j = \lambda_j \mathbf{v}_{\sigma(j)} \quad (\lambda_j \in F^\times),$$

then $\pi(m) = \sigma$. This is a homomorphism. Its kernel consists exactly of the determinant-one monomial matrices whose underlying permutation is the identity; these are precisely the determinant-one diagonal matrices, so

$$\ker \pi = T_0.$$

The map π is surjective. Indeed, if $\sigma \in S_n$ and p_σ is its permutation matrix, choose $d \in T$ with $\det(d) = \det(p_\sigma)^{-1}$. Then $dp_\sigma \in N_0$ and $\pi(dp_\sigma) = \sigma$. Hence, by the Fundamental Theorem on Homomorphisms,

$$W_0 = N_0/T_0 \cong S_n.$$

For $1 \leq i \leq n-1$, let $\dot{s}_i \in N_0$ be the monomial matrix which is the identity away from rows and columns $i, i+1$ and has block

$$\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$$

on those rows and columns. This block has determinant 1, so $\dot{s}_i \in N_0$. Let $s_i = \dot{s}_i T_0 \in W_0$. Under the isomorphism $W_0 \cong S_n$, the element s_i corresponds to the adjacent transposition $(i \ i+1)$. Also $\dot{s}_i^2 \in T_0$, so $\dot{s}_i^2 = 1$ in W_0 . Thus

$$S_0 = \{s_1, \dots, s_{n-1}\}$$

is a finite set of involutions generating W_0 , because the adjacent transpositions generate S_n .

It remains to check the two double-coset conditions. Let $s_i \in S_0$, let $w \in W_0$, and choose a representative $\dot{w} \in N_0$. From the BN -pair for $\mathrm{GL}(n, F)$ proved in Exercise 4.8, we have

$$\dot{s}_i B \dot{w} \subseteq B \dot{w} B \cup B \dot{s}_i \dot{w} B.$$

Here we may use the determinant-one representatives $\dot{s}_i$ and $\dot{w}$, because changing a representative by an element of T does not change the corresponding B -double coset. Now take $x \in \dot{s}_i B_0 \dot{w}$. Then $x \in G_0$, and the displayed inclusion puts x in either $B \dot{w} B$ or $B \dot{s}_i \dot{w} B$. Since both $\dot{w}$ and $\dot{s}_i \dot{w}$ lie in N_0 , the determinant-one double-coset observation gives

$$x \in B_0 \dot{w} B_0 \quad \text{or} \quad x \in B_0 \dot{s}_i \dot{w} B_0.$$

Therefore

$$\dot{s}_i B_0 \dot{w} \subseteq B_0 \dot{w} B_0 \cup B_0 \dot{s}_i \dot{w} B_0.$$

Finally, we show that

$$\dot{s}_i B_0 \dot{s}_i \not\subseteq B_0.$$

The transvection $X_{i,i+1}(1)$ lies in B_0 . In the $i, i+1$ rows and columns, the product

$$\dot{s}_i X_{i,i+1}(1) \dot{s}_i$$

has block

$$\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} = \begin{pmatrix} -1 & 0 \\ 1 & -1 \end{pmatrix}.$$

This matrix has a non-zero entry below the diagonal, so it is not in B_0 . Hence $\dot{s}_i B_0 \dot{s}_i \not\subseteq B_0$. All the defining conditions are satisfied, so B_0 and N_0 form a BN -pair of $\mathrm{SL}(n, F)$, and the associated Weyl group is $W_0 \cong S_n$. $\square$

Exercise 6.11. Let $P = \mathrm{PSL}(n, F)$, and let $\overline{B_0}$, $\overline{N_0}$, and $\overline{T_0}$ be the images in P of the subgroups B_0 , N_0 , and T_0 of $\mathrm{SL}(n, F)$. Show that $\overline{B_0}$ and $\overline{N_0}$ form a BN -pair of P (with again the Weyl group $\overline{W_0} = \overline{N_0}/\overline{T_0}$ being isomorphic with S_n).

Solution. Let

$$q: \mathrm{SL}(n, F) \rightarrow P = \mathrm{PSL}(n, F)$$

be the natural map, and let

$$K = \ker q.$$

Then K consists of the scalar matrices in $\mathrm{SL}(n, F)$. In particular $K \leq T_0$, since scalar matrices are diagonal. We write $\bar{x} = q(x)$ for $x \in \mathrm{SL}(n, F)$.

By Exercise 6.10, the subgroups B_0 and N_0 form a BN -pair of $\mathrm{SL}(n, F)$. We check that the defining conditions pass to the quotient. First,

$$P = q(\mathrm{SL}(n, F)) = q(\langle B_0, N_0 \rangle) = \langle \overline{B_0}, \overline{N_0} \rangle.$$

Next we compute the intersection. Certainly $\overline{T_0} \leq \overline{B_0} \cap \overline{N_0}$. Conversely, suppose that

$$y \in \overline{B_0} \cap \overline{N_0}.$$

Then $y = \bar{b} = \bar{n}$ for some $b \in B_0$ and $n \in N_0$. Thus

$$bn^{-1} \in K \leq T_0 \leq N_0.$$

Since $n \in N_0$, it follows that $b \in N_0$. But also $b \in B_0$, so

$$b \in B_0 \cap N_0 = T_0.$$

Hence $y = \bar{b} \in \overline{T_0}$, and therefore

$$\overline{B_0} \cap \overline{N_0} = \overline{T_0}.$$

Also $\overline{T_0} \leq \overline{N_0}$, because $T_0 \leq N_0$.

Now we identify the Weyl group. Define

$$\theta: N_0/T_0 \rightarrow \overline{N_0}/\overline{T_0}, \quad \theta(nT_0) = \bar{n}\overline{T_0}.$$

This map is well-defined because replacing n by nt with $t \in T_0$ replaces $\bar{n}$ by $\bar{n}\bar{t}$, which gives the same coset modulo $\overline{T_0}$. It is also surjective. If $\theta(nT_0) = \overline{T_0}$, then $\bar{n} \in \overline{T_0}$, so $\bar{n} = \bar{t}$ for some $t \in T_0$. Hence $nt^{-1} \in K \leq T_0$, and therefore $n \in T_0$. Thus $\ker \theta = 1$, and θ is an isomorphism. Since Exercise 6.10 gives $N_0/T_0 \cong S_n$, we get

$$\overline{W_0} = \overline{N_0}/\overline{T_0} \cong S_n.$$

In particular $\overline{W_0}$ is generated by the images of the same adjacent-transposition generators used in Exercise 6.10. If $\dot{s}_i \in N_0$ is the determinant-one monomial representative with block

$$\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$$

in rows and columns $i, i+1$, then $\overline{\dot{s}_i} \overline{T_0}$ is an involution in $\overline{W_0}$, and these elements generate $\overline{W_0}$.

It remains to verify the two double-coset conditions. Let s_i be one of the above generators, and let

$w \in \overline{W_0}$. Choose representatives $\dot{s}_i \in N_0$ and $\dot{w} \in N_0$ whose images in P represent s_i and w . In $\text{SL}(n, F)$, Exercise 6.10 gives

$$\dot{s}_i B_0 \dot{w} \subseteq B_0 \dot{w} B_0 \cup B_0 \dot{s}_i \dot{w} B_0.$$

Applying q gives

$$\overline{\dot{s}_i} \overline{B_0} \overline{\dot{w}} \subseteq \overline{B_0} \overline{\dot{w}} \overline{B_0} \cup \overline{B_0} \overline{\dot{s}_i} \overline{\dot{w}} \overline{B_0}.$$

This is the first double-coset condition for P .

Finally, we prove the non-containment condition. In Exercise 6.10, the element

$$y = \dot{s}_i X_{i,i+1}(1) \dot{s}_i$$

lies in $\dot{s}_i B_0 \dot{s}_i$ but does not lie in B_0 . We claim that $\overline{y} \notin \overline{B_0}$. Indeed, if $\overline{y} \in \overline{B_0}$, then $\overline{y} = \overline{b}$ for some $b \in B_0$. Hence $y b^{-1} \in K \leq T_0 \leq B_0$, so $y \in B_0$, a contradiction. Therefore

$$\overline{\dot{s}_i} \overline{B_0} \overline{\dot{s}_i} \not\subseteq \overline{B_0}.$$

All the defining conditions are satisfied, so $\overline{B_0}$ and $\overline{N_0}$ form a BN -pair of $P = \text{PSL}(n, F)$, and the Weyl group is again isomorphic with S_n . $\square$

Chapter 3

Local Structure

7 Sylow's Theorem

Setup and notation

- Throughout the section, G is a finite group and p a prime divisor of $|G|$.
- $|G|_p$ denotes the highest power of p dividing $|G|$, so $|G|_p = p^n$ with $p^n \mid |G|$ but $p^{n+1} \nmid |G|$.

p -elements, p -groups, p -subgroups

- $g \in G$ is a p -element if its order is a power of p .
- G is a p -group if $|G|$ is a power of p ; $H \leq G$ is a p -subgroup if $|H|$ is a power of p .
- Every element of a p -group is a p -element. (For infinite groups, p -group means every element is a p -element.)
- $H \leq G$ is a *Sylow p -subgroup* of G if H is a p -subgroup of order $|G|_p$, the maximal possible order of a p -subgroup of G .
- A subgroup of G is a *Sylow subgroup* if it is a Sylow p -subgroup for some prime divisor p of $|G|$.

Motivating example: $\text{GL}(n, p)$

- For $G = \text{GL}(n, p)$, Proposition 4.2 gives $|G| = p^{\frac{n(n-1)}{2}}(p^n - 1) \cdots (p - 1)$, so $|G|_p = p^{\frac{n(n-1)}{2}}$.
- The subgroup U of upper uni-triangular matrices satisfies $|U| = p^{n-1}p^{n-2} \cdots p^2p = p^{\frac{n(n-1)}{2}} = |G|_p$, hence U is a Sylow p -subgroup.
- Every p -element of $\text{GL}(n, p)$ is unipotent (its characteristic polynomial is $(X-1)^n$), and by Kolchin's Theorem any p -subgroup is conjugate with a subgroup of U .
- In particular: $\text{GL}(n, p)$ has a Sylow p -subgroup, all are conjugate, and every p -subgroup is contained in one. Sylow's Theorem extends this to arbitrary finite G .

Theorem (Sylow's Theorem). (i) G has at least one Sylow p -subgroup.

(ii) All the Sylow p -subgroups of G are conjugate.

(iii) Any p -subgroup of G is contained in a Sylow p -subgroup.

(iv) The number of Sylow p -subgroups of G is congruent to 1 modulo p .

Proof. Let $|G| = p^n m$, where $p^n = |G|_p$ and $p \nmid m$. Let X be the set of all subsets of G having p^n elements. Then G acts on X by left multiplication.

We first examine the orbits of this action whose sizes are not divisible by p . Suppose that $\mathcal{O}$ is such an orbit. Choose some $A \in \mathcal{O}$. Since A is non-empty, there is some $a \in A$, and then $a^{-1}A$ lies in the same orbit and contains 1. Replacing A by $a^{-1}A$, we may assume that $1 \in A$. Let P be the stabilizer of A . If $x \in P$, then $xA = A$, so $x = x1 \in A$. Hence $P \subseteq A$, and therefore

$$|P| \leq |A| = p^n.$$

By Corollary 3.5, we have

$$|\mathcal{O}| = |G : P|,$$

and hence $|G| = |P| |\mathcal{O}|$. Since $p \nmid |\mathcal{O}|$, the full p -part p^n of $|G|$ must divide $|P|$. Together with $|P| \leq p^n$, this gives $|P| = p^n$. Thus P is a Sylow p -subgroup of G . Since $P \subseteq A$ and $|P| = |A|$, we have $A = P$, so $\mathcal{O}$ is the coset space G/P .

Conversely, if P is a Sylow p -subgroup of G , then the set of left cosets G/P is contained in X , and it is exactly the orbit of the subset P under left multiplication. This orbit has size

$$|G : P| = m,$$

which is not divisible by p . Therefore the Sylow p -subgroups of G are in bijective correspondence with the orbits in X whose sizes are not divisible by p .

Let X' be the union of all orbits in X whose sizes are not divisible by p , and let r be the number of Sylow p -subgroups of G . Every orbit in $X - X'$ has size divisible by p , so

$$|X| \equiv |X'| \pmod{p}.$$

Each orbit contained in X' has size m , and there are r such orbits. Hence

$$|X'| = rm,$$

so

$$rm \equiv |X| = \binom{p^n m}{p^n} \pmod{p}.$$

This congruence depends only on the order of the group. Now compare with the cyclic group of order $p^n m$. By Theorem 1.4, this cyclic group has exactly one subgroup of order p^n , so the same congruence gives

$$m \equiv \binom{p^n m}{p^n} \pmod{p}.$$

Since $p \nmid m$, we may cancel m modulo p , obtaining

$$r \equiv 1 \pmod{p}.$$

This proves (iv), and in particular $r > 0$, so G has at least one Sylow p -subgroup. This proves (i).

It remains to prove conjugacy and containment. Let P be a Sylow p -subgroup of G , and let Q be an arbitrary p -subgroup of G . Let Y be the set of conjugates of P in G . The group Q acts on Y by conjugation. Each orbit of this action has size equal to the index of a subgroup of the p -group Q , and hence has size a power of p .

By Proposition 3.14,

$$|Y| = |G : N_G(P)|.$$

Since $P \leq N_G(P)$, the index $|G : N_G(P)|$ divides

$$|G : P| = m.$$

Thus $p \nmid |Y|$. Therefore at least one orbit of the action of Q on Y has size 1. Let this orbit be $\{P_1\}$, where P_1 is a conjugate of P . Then $xP_1x^{-1} = P_1$ for every $x \in Q$, so $QP_1 = P_1Q$. By Proposition 1.3, $QP_1 \leq G$. Moreover, by Proposition 3.12,

$$|QP_1| = \frac{|Q||P_1|}{|Q \cap P_1|},$$

so QP_1 is a p -subgroup of G . Since P_1 is already a Sylow p -subgroup and $P_1 \leq QP_1$, we must have $QP_1 = P_1$. Hence $Q \leq P_1$. This proves that every p -subgroup of G is contained in a Sylow p -subgroup, proving (iii).

Finally, if Q itself is a Sylow p -subgroup of G , then $Q \leq P_1$ and $|Q| = |P_1|$, so $Q = P_1$. Since P_1 is a conjugate of P , every Sylow p -subgroup of G is conjugate to P . This proves (ii). $\square$

Corollary 7.1. *The number of Sylow p -subgroups of G divides $|G|/|G|_p$.*

Proof. Let r be the number of Sylow p -subgroups of G , and let P be one of them. By part (ii) of Sylow's Theorem, every Sylow p -subgroup of G is conjugate to P , so r is exactly the number of conjugates of P in G . By Proposition 3.14, this number is

$$r = |G : N_G(P)|.$$

Since $P \leq N_G(P) \leq G$, we have

$$|G : P| = |G : N_G(P)| |N_G(P) : P|.$$

Therefore $r = |G : N_G(P)|$ divides

$$|G : P| = \frac{|G|}{|P|} = \frac{|G|}{|G|_p},$$

as required. $\square$

Theorem (Cauchy's Theorem). *G has an element of order p .*

Proof. By part (i) of Sylow's Theorem, G has a Sylow p -subgroup P . Since p divides $|G|$, this subgroup is non-trivial. Choose some $1 \neq x \in P$. Then $o(x)$ is a power of p , say $o(x) = p^k$ with $k \geq 1$. Therefore

$$o(x^{p^{k-1}}) = p.$$

Thus G has an element of order p . $\square$

(Other proofs of Sylow's Theorem often assume that Cauchy's Theorem is already known, as was the case historically.)

Sylow subgroups of normal subgroups and quotients

- Sylow p -subgroups behave well with respect to a normal subgroup $N \trianglelefteq G$: both PN/N and $P \cap N$ are Sylow p -subgroups (of G/N and N respectively).
- For a non-normal subgroup $H \leq G$, $P \cap H$ need not be Sylow in H , but some conjugate $gPg^{-1} \cap H$ is.

Proposition 7.2. *Let $N \trianglelefteq G$, and let P be a Sylow p -subgroup of G . Then PN/N is a Sylow p -subgroup of G/N , and $P \cap N$ is a Sylow p -subgroup of N .*

Proof. By Proposition 1.7, PN is a subgroup of G . The Correspondence Theorem applied to the natural map $G \rightarrow G/N$ gives

$$|G/N : PN/N| = |G : PN|.$$

Since $P \leq PN \leq G$, the index $|G : PN|$ divides $|G : P|$. But P is a Sylow p -subgroup of G , so

$$|G : P| = \frac{|G|}{|G|_p}$$

is not divisible by p . Hence $p \nmid |G/N : PN/N|$. On the other hand, by the First Isomorphism Theorem,

$$PN/N \cong P/(P \cap N),$$

so PN/N is a p -group. Thus PN/N is a p -subgroup of G/N whose index in G/N is not divisible by p , and therefore PN/N is a Sylow p -subgroup of G/N .

It remains to prove that $P \cap N$ is Sylow in N . Clearly $P \cap N$ is a p -subgroup. By Proposition 3.12,

$$|PN| = \frac{|P||N|}{|P \cap N|},$$

and hence

$$|N : P \cap N| = \frac{|N|}{|P \cap N|} = \frac{|PN|}{|P|} = |PN : P|.$$

Since $P \leq PN \leq G$, the index $|PN : P|$ divides $|G : P|$, and again $p \nmid |G : P|$. Therefore $p \nmid |N : P \cap N|$. Hence $P \cap N$ is a p -subgroup of N whose index in N is not divisible by p , so $P \cap N$ is a Sylow p -subgroup of N . $\square$

Applications: groups of order pq and simplicity of A_5

- Sylow's theorem classifies groups of order pq (distinct primes $p > q$).
- It also gives a clean count of p -elements via the orbit count of Sylow p -subgroups, used to prove A_5 is simple.

Proposition 7.3. *Let p and q be distinct primes, with $p > q$. If $p \not\equiv 1 \pmod{q}$, then any group of order pq is isomorphic with $\mathbb{Z}_{pq}$. If $p \equiv 1 \pmod{q}$, then any abelian group of order pq is isomorphic with $\mathbb{Z}_{pq}$, and there is exactly one isomorphism class of non-abelian groups of order pq .*

Proof. Let G be a group of order pq . Let P be a Sylow p -subgroup of G , and let Q be a Sylow q -subgroup of G . Then $|P| = p$ and $|Q| = q$, so $P \cong \mathbb{Z}_p$ and $Q \cong \mathbb{Z}_q$. Also, $P \cap Q = 1$, and Proposition 3.12 gives

$$|PQ| = \frac{|P||Q|}{|P \cap Q|} = pq = |G|,$$

so $G = PQ$.

We first count the Sylow subgroups. By Sylow's Theorem and Corollary 7.1, the number of Sylow p -subgroups divides q and is congruent to 1 modulo p . Since $p > q$, we have $q \not\equiv 1 \pmod{p}$, so there is exactly one Sylow p -subgroup. Hence $P \trianglelefteq G$. Similarly, the number of Sylow q -subgroups divides p and is congruent to 1 modulo q . Therefore this number is either 1 or p , and the second case can occur only when $p \equiv 1 \pmod{q}$.

If $Q \trianglelefteq G$, then both P and Q are normal in G . Since $G = PQ$ and $P \cap Q = 1$, Lemma 2.7 gives

$$G \cong P \times Q \cong \mathbb{Z}_p \times \mathbb{Z}_q \cong \mathbb{Z}_{pq},$$

where the final isomorphism follows from Lemma 2.8. Thus G is cyclic whenever Q is normal. This proves the first assertion when $p \not\equiv 1 \pmod{q}$, and it also proves that every abelian group of order pq is cyclic, since then Q is automatically normal.

It remains to classify the non-abelian case, so assume $p \equiv 1 \pmod{q}$ and that G is non-abelian. Then Q is not normal, while $P \trianglelefteq G$, $G = PQ$, and $P \cap Q = 1$. Hence $G = P \rtimes Q$. Let

$$\varphi: Q \rightarrow \text{Aut}(P)$$

be the conjugation homomorphism. If $\ker \varphi \neq 1$, then $\ker \varphi = Q$, because Q has prime order. In that case the action is trivial, so $G = P \times Q$, contradicting that G is non-abelian. Therefore φ is injective. Thus every non-abelian group of order pq arises from a monomorphism

$$\mathbb{Z}_q \rightarrow \text{Aut}(\mathbb{Z}_p).$$

Conversely, any such monomorphism gives a non-abelian semidirect product $\mathbb{Z}_p \rtimes_{\varphi} \mathbb{Z}_q$ of order pq .

By Proposition 2.2, we have $\text{Aut}(\mathbb{Z}_p) \cong \mathbb{Z}_{p-1}$. Since $q \mid (p-1)$, Theorem 1.4 shows that $\text{Aut}(\mathbb{Z}_p)$ has a unique subgroup K of order q . Hence a monomorphism $\varphi: \mathbb{Z}_q \rightarrow \text{Aut}(\mathbb{Z}_p)$ exists, and every such monomorphism has image K . If $\psi: \mathbb{Z}_q \rightarrow \text{Aut}(\mathbb{Z}_p)$ is another monomorphism, then $\varphi(\mathbb{Z}_q) = K = \psi(\mathbb{Z}_q)$, so Proposition 2.11 gives

$$\mathbb{Z}_p \rtimes_{\varphi} \mathbb{Z}_q \cong \mathbb{Z}_p \rtimes_{\psi} \mathbb{Z}_q.$$

Therefore, when $p \equiv 1 \pmod{q}$, there is exactly one isomorphism class of non-abelian groups of order pq . $\square$

Theorem 7.4. A_5 is simple.

Proof. We have

$$|A_5| = \frac{5!}{2} = 60 = 2^2 \cdot 3 \cdot 5.$$

We first count the elements of prime-power order that will be needed below. A 5-cycle is even, and the number of 5-cycles in S_5 is $4! = 24$. Each Sylow 5-subgroup of A_5 has order 5 and hence contains 4 non-identity elements, and two distinct subgroups of order 5 intersect trivially. Thus A_5 has

$$\frac{24}{4} = 6$$

Sylow 5-subgroups, and these account for all 24 elements of order 5. Similarly, the 3-cycles are even, and there are

$$\binom{5}{3} \cdot 2 = 20$$

of them. Each Sylow 3-subgroup has two non-identity elements, so A_5 has 10 Sylow 3-subgroups, accounting for all 20 elements of order 3.

Now consider the Sylow 2-subgroups. For $1 \leq i \leq 5$, let $\{a, b, c, d\}$ be the complement of $\{i\}$ in $\{1, 2, 3, 4, 5\}$, and put

$$V_i = \{1, (ab)(cd), (ac)(bd), (ad)(bc)\}.$$

Then V_i is a subgroup of A_5 of order 4, hence a Sylow 2-subgroup. If $i \neq j$, then $V_i \cap V_j = 1$, since a non-identity element of V_i fixes exactly the point i . Moreover, for $\sigma \in A_5$ we have

$$\sigma V_i \sigma^{-1} = V_{\sigma(i)}.$$

Since A_5 acts transitively on $\{1, 2, 3, 4, 5\}$, the five subgroups $V_1, \dots, V_5$ are conjugate. By part (ii) of Sylow's Theorem, these are all the Sylow 2-subgroups of A_5 . Thus A_5 has $5(4 - 1) = 15$ elements of order 2. Also, every element of order 2 in A_5 has a distinct conjugate in A_5 : for instance, if $x = (ab)(cd)$, with i the remaining point, then conjugation by the 3-cycle (acb) sends x to $(ac)(bd) \neq x$.

Now let $N \triangleleft A_5$ be a proper normal subgroup, and put $n = |N|$. By Lagrange's Theorem, n divides 60, and since N is proper we have $n \leq 30$. Suppose first that 5 divides n . Then by Cauchy's Theorem, N contains an element of order 5, and hence contains a Sylow 5-subgroup of A_5 . Since N is normal in A_5 , it contains every conjugate of this Sylow 5-subgroup, so it contains all 24 elements of order 5. This forces $n = 30$. But then 3 divides n , so the same argument with Sylow 3-subgroups shows that N contains all 20 elements of order 3. These are disjoint from the 24 elements of order 5, and with the identity this gives more than 30 elements in N , a contradiction. Therefore $5 \nmid n$.

If 3 divides n , then again by Cauchy's Theorem, N contains an element of order 3, and normality forces N to contain all 20 elements of order 3. But $5 \nmid n$ and $n \mid 60$, so $n \leq 12$, a contradiction. Hence $3 \nmid n$. Therefore n is one of 1, 2, 4. If $n = 4$, then N is a Sylow 2-subgroup of A_5 , but a Sylow 2-subgroup has five conjugates, contradicting the normality of N . If $n = 2$, then $N = \{1, x\}$ for some element x of order 2, but we have just observed that x has a conjugate different from itself, again contradicting normality. Hence $n = 1$.

Thus every proper normal subgroup of A_5 is trivial, and so A_5 is simple. $\square$

Theorem 7.5. *Any simple group of order 60 is isomorphic with A_5 .*

Proof. Let G be a simple group of order 60. We first observe that G has no proper subgroup of index less than 5. Indeed, if $H < G$ has index r , then the action of G on G/H by left multiplication gives a homomorphism

$$G \rightarrow S_r$$

by Proposition 3.1. Since this action is transitive and $r > 1$, it is non-trivial. The kernel is therefore a proper normal subgroup of the simple group G , and hence the kernel is trivial. Thus G embeds in S_r . But if $r < 5$, then

$$|S_r| = r! < 60 = |G|,$$

which is impossible.

We next show that G has a subgroup of index 5. Suppose, for a contradiction, that it does not. Let S be a Sylow 2-subgroup of G . Then $|S| = 4$. By Corollary 7.1, the number of Sylow 2-subgroups divides $60/4 = 15$, and by Proposition 3.14 this number is the index $|G : N_G(S)|$. This number is not 1, since otherwise S would be normal in G , contradicting simplicity. Since G has no proper subgroup of index less than 5 and, by assumption, none of index 5, this number must be 15.

Let S_1 and S_2 be distinct Sylow 2-subgroups. Suppose that there is some $1 \neq t \in S_1 \cap S_2$. Since every group of order 4 is abelian, both S_1 and S_2 lie in $C_G(t)$. Thus $|C_G(t)| > 4$, and since $S_1 \leq C_G(t)$, the order $|C_G(t)|$ is divisible by 4. Hence

$$|G : C_G(t)| \leq 5.$$

The first paragraph and our assumption that G has no subgroup of index 5 force $C_G(t) = G$. Thus $t \in Z(G)$, so $\langle t \rangle$ is a non-trivial proper normal subgroup of G , contradicting simplicity. Therefore distinct Sylow 2-subgroups intersect trivially. Since there are 15 Sylow 2-subgroups, G has

$$15(4 - 1) = 45$$

non-identity elements lying in Sylow 2-subgroups.

Since G is simple, its Sylow 5-subgroup is not normal. By Sylow's Theorem and Corollary 7.1, the number of Sylow 5-subgroups divides $60/5 = 12$ and is congruent to 1 modulo 5, so this number must be 6. These subgroups intersect trivially away from the identity, and hence contribute

$$6(5 - 1) = 24$$

elements of order 5. This is impossible, since $45 + 24 > 60$. Therefore G has a subgroup of index 5.

Let H be a subgroup of index 5. The action of G on G/H gives a homomorphism $G \rightarrow S_5$. As above, simplicity forces this homomorphism to be injective, so we identify G with its image in S_5 . Then $|G| = 60$, and hence $|S_5 : G| = 2$. By Proposition 1.8, $G \trianglelefteq S_5$.

We claim that $G = A_5$. Suppose not. Since $A_5 \trianglelefteq S_5$, Proposition 1.7 gives $GA_5 \leq S_5$. The subgroup GA_5 contains A_5 properly, because $G \neq A_5$ and both groups have order 60. Hence $|GA_5| > 60$, so $GA_5 = S_5$. By Proposition 3.12,

$$|G \cap A_5| = \frac{|G||A_5|}{|GA_5|} = \frac{60 \cdot 60}{120} = 30.$$

Therefore $G \cap A_5$ has index 2 in G , so Proposition 1.8 gives $G \cap A_5 \trianglelefteq G$. This is a non-trivial proper normal subgroup of G , contradicting simplicity. Thus $G = A_5$, and the original group is isomorphic with A_5 . $\square$

Corollary 7.6. $\text{PSL}(2, 4) \cong \text{PSL}(2, 5) \cong A_5$.

Proof. By Proposition 6.5, we have

$$|\text{PSL}(2, 4)| = 4^3 - 4 = 60$$

and

$$|\text{PSL}(2, 5)| = \frac{1}{2}(5^3 - 5) = 60.$$

By Theorem 6.8, both groups are simple, since the exceptional groups in degree 2 occur only over fields of order 2 and 3. Therefore Theorem 7.5 gives

$$\text{PSL}(2, 4) \cong A_5 \quad \text{and} \quad \text{PSL}(2, 5) \cong A_5.$$

Hence $\text{PSL}(2, 4) \cong \text{PSL}(2, 5) \cong A_5$. $\square$

It is a relatively easy exercise to show that the only simple groups of order less than 60 are the cyclic groups of prime order, and hence that A_5 is the non-abelian finite simple group having smallest order. The non-abelian finite simple group of next smallest order is $\text{PSL}(2, 7)$, which has order 168.

Exercises

Throughout these exercises, G is a finite group and p is a prime divisor of $|G|$.

Exercise 7.1. Let $H \leq G$ and let P be a Sylow p -subgroup of G . Prove, without reference to Sylow's Theorem, that there is some conjugate of P whose intersection with H is a Sylow p -subgroup of H .

Solution. $\square$

Exercise 7.2. (cont.) Use Exercise 7.1 to give an alternate proof of part (i) of Sylow's Theorem.

Solution. $\square$

Exercise 7.3. Give an alternate proof of parts (ii) and (iii) of Sylow's Theorem by considering the action of an arbitrary p -subgroup Q of G on the coset space G/P , where P is a Sylow p -subgroup of G .

Solution. $\square$

Exercise 7.4. Prove the following generalization of part (iv) of Sylow's Theorem: If $|G|$ is divisible by p^b , and $H \leq G$ has order p^a where $a \leq b$, then the number of subgroups of G that both contain H and have order p^b is congruent to 1 modulo p .

Solution. $\square$

Exercise 7.5. Show that if P is a Sylow p -subgroup of G , then $N_G(P)$ is a self-normalizing subgroup of G , meaning that $N_G(N_G(P)) = N_G(P)$. (Any subgroup of G that can be written as $N_G(P)$ for some non-trivial p -subgroup P of G is said to be a p -local subgroup of G .)

Solution. □

Exercise 7.6. Given a prime p , find an example of a finite group having exactly $1 + p$ Sylow p -subgroups. Can this be done for $1 + 2p$? $1 + 3p$?

Solution. □

Exercise 7.7. Show that if G has exactly $1 + kp$ Sylow p -subgroups for some $k \in \mathbb{N}$, then there is a subgroup of S_{1+kp} having exactly $1 + kp$ Sylow p -subgroups.

Solution. □

Exercise 7.8. Let $n \geq 6$, and assume by induction that A_{n-1} is simple. Let N be a non-trivial proper normal subgroup of A_n .

- (a) Show that $A_n = N \rtimes A_{n-1}$.
- (b) Let $N^\#$ be the set of non-identity elements of N , and consider the action of A_{n-1} on $N^\#$ given by the conjugation homomorphism from A_{n-1} to $\text{Aut}(N)$. Show that $N^\#$ is isomorphic as an A_{n-1} -set to $\{1, \dots, n-1\}$.
- (c) Show that A_{n-1} acts triply transitively on $N^\#$. (An action of a group G on a set X is *triply transitive* if for every pair of triples (x_1, x_2, x_3) and (y_1, y_2, y_3) , where $x_i, y_i \in X$ for all i and $x_i \neq x_j$ (resp., $y_i \neq y_j$) when $i \neq j$, there is some $g \in G$ such that $gx_i = y_i$ for all i .)
- (d) Derive a contradiction, and conclude that A_n is simple.

Solution. □

The remaining exercises develop a proof that if G is a finite simple group with $|G| \leq 200$, then either $G \cong \mathbb{Z}_p$ for some prime p , or $G \cong A_5$, or $G \cong \text{PSL}(2, 7)$. This was established by Hölder in 1892. We will need to assume two facts, which will be proved in later sections:

- A. If $|G| = p^n$ where $n > 1$, then G is not simple (Section 8).
- B. If G is a p -group and $H < G$, then $H < N_G(H)$ (Section 11).

Exercise 7.9. Show that G is not simple whenever one of the following statements is true, where p is an odd prime:

- (a) $|G| = 2^m p^n$, where $2^k \not\equiv 1 \pmod{p}$ for any $1 \leq k \leq m$.
- (b) $|G| = p^n q$, where $q \neq p$ is prime and $q \not\equiv 1 \pmod{p}$.
- (c) $|G| = ap$, where $p \nmid a$ and $kp + 1 \nmid a$ for any $k \in \mathbb{N}$.

Solution. □

Exercise 7.10. Show that G is not simple whenever one of the following statements is true:

- (a) $|G| = p^a(p+1)$, where $a > 1$.
- (b) $|G| = p^a(p+3)$, where $a > 3$ if $p = 2$ and $a > 1$ if $p > 3$.
- (c) $|G| = p^a(p^2 - 1)$, where $a > 1$ and p is odd.

Solution. □

Exercise 7.11. Show that if $|G| \in \{30, 56, 105, 132\}$, then G is not simple.

Solution. □

Exercise 7.12. Show that G is not simple in the following cases:

- (a) $|G| = 90$. (HINT: Show that if Q is a Sylow 3-subgroup of G and $1 \neq x \in Q$, then $C_G(x) = Q$.)
- (b) $|G| = 112$. (HINT: Use fact B above to show that any two Sylow 2-subgroups of G intersect trivially.)
- (c) $|G| = 120$. (HINT: Show that G imbeds in S_6 , and contradict Exercise 3.8.)
- (d) $|G| = 144$. (HINT: Argue as in part (i) above.)
- (e) $|G| = 180$. (HINT: Show that if Q is a Sylow 3-subgroup of G and $1 \neq x \in Q$, then $|C_G(x) : Q| \leq 2$.)

Solution. □

From fact A above and Exercises 9–12, we can conclude that if G is a simple group with $|G| \leq 200$, then $|G|$ either is prime or is equal to either 60 or 168. (Verify this.) In light of Theorem 7.5 and Theorem 6.8, all we need now show is that any two simple groups of order $168 = 2^3 \cdot 3 \cdot 7$ are isomorphic. We accomplish this in Exercises 13–17 below by showing that any simple group of order 168 is isomorphic with a specific subgroup of A_8 . (Compare with Exercise 6.9.) Let G be a simple group of order 168, let $P = \langle x \rangle$ be a Sylow 7-subgroup of G , and let $H = N_G(P)$.

Exercise 7.13. Show that $H = \{g \in G \mid gxg^{-1} \in P\}$ is a non-abelian group of order 21 which is generated by x and y , where y has order 3 and $xyx^{-1} = x^4$.

Solution. □

Exercise 7.14. (cont.) Show that $N_G(\langle y \rangle)$ is isomorphic with S_3 and is generated by y and z , where z has order 2 and $zyz = z$.

Solution. □

Exercise 7.15. (cont.) Show that $G/H = \{H, zH, xzH, \dots, x^6zH\}$. Conclude that $G = \langle x, y, z \rangle$.

Solution. □

Let $\varphi: G \rightarrow S_8$ be the monomorphism corresponding to the action of G on the coset space G/H . Observe that $\varphi(G) = \varphi(G)' \leq S_8' \leq A_8$. View S_8 as the group of permutations of the set $\{\infty, 0, 1, 2, 3, 4, 5, 6\}$; for each $0 \leq i \leq 6$, associate i with the coset $x^i zH$, and associate ∞ with the coset H .

Exercise 7.16. (cont.) Show that $\varphi(x) = (0\ 1\ 2\ 3\ 4\ 5\ 6)$, that $\varphi(y) = (1\ 4\ 2)(3\ 5\ 6)$, and that $\varphi(z) = z_i$ for some i , where $z_1 = (\infty\ 0)(1\ 3)(2\ 5)(4\ 6)$, $z_2 = (\infty\ 0)(1\ 5)(2\ 6)(3\ 4)$, and $z_3 = (\infty\ 0)(1\ 6)(2\ 3)(4\ 5)$.

Solution. □

Exercise 7.17. (cont.) Let $L_i = \langle \varphi(x), \varphi(y), z_i \rangle \leq A_8$ for $i = 1, 2, 3$. Show that $L_1 = L_2 = L_3$. Conclude that G is uniquely determined up to isomorphism.

Solution. □

8 Finite p -groups

Why study finite p -groups?

- Sylow's Theorem reduces many questions about a finite group to questions about its subgroups of prime-power order.
- Finite p -groups form a rich subject in their own right; this section only scratches the surface.

The center and maximal subgroups

- The center of a non-trivial finite p -group is non-trivial (Theorem 8.1); more generally, every non-trivial normal subgroup meets $Z(P)$ non-trivially.
- Consequently, the only finite simple p -groups are the groups $\mathbb{Z}_p$ of prime order.
- For non-abelian G , the quotient $G/Z(G)$ is never cyclic (Lemma 8.2); in particular $|P : Z(P)| \neq p$.
- Every maximal subgroup of a finite p -group is normal of index p (Proposition 8.3, Corollary 8.4).

Abelian p -groups and the basis theorem

- Every finite abelian p -group is a direct product of cyclic p -groups (Theorem 8.6).
- The non-generators of a cyclic p -group are exactly its p th powers (Lemma 8.5).
- Combining Theorems 8.6 and 8.7 yields the *basis theorem*: every finite abelian group is a direct product of cyclic p -groups (Corollary 8.8).

Nilpotent groups and the Frattini argument

- The Frattini Argument states that $G = N_G(P)N$ whenever P is a Sylow subgroup of a normal subgroup $N \trianglelefteq G$.
- A finite group G is *nilpotent* if every Sylow subgroup of G is normal; equivalently, G is the direct product of its Sylow subgroups, equivalently every maximal subgroup is normal (Theorem 8.7).
- Finite abelian groups and finite p -groups are nilpotent; an alternative definition of nilpotence which extends to infinite groups is taken up in Section 11.

Groups of small order

- Any group of order p is isomorphic with $\mathbb{Z}_p$.
- Any group of order p^2 is abelian (by Theorem 8.1 and Lemma 8.2) and is isomorphic with either $\mathbb{Z}_{p^2}$ or $\mathbb{Z}_p \times \mathbb{Z}_p$.
- Any abelian group of order p^3 is isomorphic with one of $\mathbb{Z}_{p^3}$, $\mathbb{Z}_{p^2} \times \mathbb{Z}_p$, or $\mathbb{Z}_p \times \mathbb{Z}_p \times \mathbb{Z}_p$ (by Theorem 8.6).
- For odd primes p , there are exactly two isomorphism classes of non-abelian groups of order p^3 : one having an element of order p^2 (Proposition 8.10) and one with no such element (Proposition 8.11).
- Lemma 8.9 provides the technical commutator identities supporting the classification.
- For $p = 2$, the non-abelian groups of order 8 are the dihedral group D_8 and the *quaternion group*; the latter is the unique group of order p^3 that cannot be written non-trivially as a semidirect product.

Theorem 8.1. *If P is a non-trivial finite p -group, then $Z(P)$ is also non-trivial. Moreover, if $1 \neq N \trianglelefteq P$, then $N \cap Z(P) \neq 1$.*

Proof.

□

Lemma 8.2. *If G is a non-abelian group, then $G/Z(G)$ cannot be cyclic. In particular, if P is a finite p -group, then $|P : Z(P)| \neq p$.*

Proof. □

Proposition 8.3. *If P is a finite p -group, then every maximal subgroup of P is normal in P .*

Proof. □

Corollary 8.4. *Any maximal subgroup of a finite p -group is of index p .*

Proof. □

Lemma 8.5. *Every non-generator of a cyclic p -group P is a p th power in P .*

Proof. □

Theorem 8.6. *A finite abelian p -group is a direct product of cyclic p -groups.*

Proof. □

Theorem (Fratini Argument). *Let G be a finite group, let $N \trianglelefteq G$, and let P be a Sylow subgroup of N . Then $G = N_G(P)N$.*

Proof. □

Theorem 8.7. *The following statements about a finite group G are equivalent:*

- (1) *Every Sylow subgroup of G is normal in G .*
- (2) *G is the direct product of its Sylow subgroups.*
- (3) *Every maximal subgroup of G is normal in G .*

Proof. □

Corollary 8.8. *A finite abelian group is a direct product of cyclic p -groups.*

Proof. □

Lemma 8.9. *Let G be a group and let $x, y \in G$. If $[x, y] \in Z(G)$, then $[x^n, y] = [x, y]^n$ and $x^n y^n = (xy)^n [x, y]^{\frac{n(n-1)}{2}}$ for any $n \in \mathbb{N}$.*

Proof. □

Proposition 8.10. *If p is an odd prime, then there is exactly one isomorphism class of non-abelian groups of order p^3 having elements of order p^2 .*

Proof. □

Proposition 8.11. *If p is an odd prime, then there is exactly one isomorphism class of non-abelian groups of order p^3 having no elements of order p^2 .*

Proof. □

Exercises

Exercise 8.1. Show that if P is a non-cyclic finite p -group, then P has a normal subgroup N such that $P/N \cong \mathbb{Z}_p \times \mathbb{Z}_p$.

Solution. □

Exercise 8.2. Let P be a group of order p^n . Show that P has a normal subgroup N_a of order p^a for every $0 \leq a \leq n$, and that these subgroups can be chosen so that N_a is contained in N_b whenever $a \leq b$.

Solution. □

Exercise 8.3. Let $G = \text{GL}(n, p)$, and let P be a Sylow p -subgroup of G . What is the order of $Z(P)$? What is the order of $Z(P/Z(P))$? If we let $Z_2(P) \leq P$ be such that $Z_2(P)/Z(P) = Z(P/Z(P))$ and continue in this way, what happens?

Solution. □

Exercise 8.4. Let U be the subgroup of $\text{GL}(n, p)$ consisting of the upper unitriangular matrices, and let Q be the subgroup of U consisting of matrices whose (i, j) -entry is zero whenever $1 < i < j < n$. Determine $Z(Q)$, and show that $Q/Z(Q)$ is abelian.

Solution. □

Exercise 8.5. Show that subgroups and quotient groups of finite nilpotent groups are nilpotent, and that direct products of finite nilpotent groups are nilpotent.

Solution. □

Exercise 8.6. Let U be the subgroup of $\text{GL}(3, p)$ consisting of the upper unitriangular matrices. Show that if p is an odd prime, then U is a non-abelian group of order p^3 having no elements of order p^2 . If $p = 2$, with which group of order 8 is U isomorphic?

Solution. □

Further Exercises

Exercise 8.7. Show that a finite group G has a largest nilpotent normal subgroup, in the sense that it contains all nilpotent normal subgroups of G . (This subgroup is called the *Fitting subgroup* of G .)

Solution. □

The intersection of all maximal subgroups of a finite group G is called the *Frattini subgroup* of G and is denoted by $\Phi(G)$.

Exercise 8.8. Show that $\Phi(G)$ is a nilpotent normal subgroup of G .

Solution. □

Exercise 8.9. Show that $g \in \Phi(G)$ iff whenever $G = \langle S \rangle$ and $g \in S$, then $G = \langle S - \{g\} \rangle$.

Solution. □

Exercise 8.10. Show that if P is a finite p -group, then $P/\Phi(P)$ is an elementary abelian p -group.

Solution. □

9 The Schur-Zassenhaus Theorem

Setting and terminology

- This section is devoted to another of the basic results of finite group theory, Schur–Zassenhaus Theorem.
- A *complement* to a normal subgroup N of a group G is a subgroup H of G such that $G = N \rtimes H$.
- A subgroup H of a finite group G is called a *Hall subgroup* if $|H|$ and $|G : H|$ are coprime.

Theorem (Schur–Zassenhaus Theorem). *Any normal Hall subgroup of a finite group has a complement.*

Proof. □

Cohomological framework

- We now wish to fit some of the ideas developed in the proof of Schur–Zassenhaus Theorem into a more general framework.
- Let H be an arbitrary group, and let A be an abelian group written additively. Suppose that we have an action of H as automorphisms of A , or equivalently a homomorphism from H to $\text{Aut}(A)$.
- A function $f : H \times H \rightarrow A$ satisfying the cocycle identity given in the above proof is called a *2-cocycle* of the pair (H, A) .
- The set of all 2-cocycles of (H, A) is denoted by $Z^2(H, A)$, and if we define $(f + g)(h_1, h_2) = f(h_1, h_2) + g(h_1, h_2)$ for $f, g \in Z^2(H, A)$ and $h_1, h_2 \in H$, then $Z^2(H, A)$ becomes an abelian group.
- A 2-cocycle is called a *2-coboundary* if there is a function $c : H \rightarrow A$ such that $f(h_1, h_2) = c(h_1) + {}^{h_1}c(h_2) - c(h_1h_2)$ for all $h_1, h_2 \in H$.
- The set of 2-coboundaries of (H, A) is denoted by $B^2(H, A)$ and is a subgroup of $Z^2(H, A)$.
- The quotient group $Z^2(H, A)/B^2(H, A)$ is called the *second cohomology group* of (H, A) and is denoted by $H^2(H, A)$. (This terminology comes from algebraic topology. We shall define the n th cohomology group of (H, A) for any $n \in \mathbb{N}$ in the further exercises to Section 12.)
- Now suppose that A is an abelian normal Hall subgroup of a finite group G . We showed during the proof of Schur–Zassenhaus Theorem that any 2-cocycle f of $(G/A, A)$ was a 2-coboundary, or equivalently that $H^2(G/A, A) = 0$. (The difference of sign between the definition of 2-coboundary given above and what was established in the proof is irrelevant.) Furthermore, we showed that this fact implied the existence of a complement to A in G . We attempt to make this connection between the second cohomology group of $(G/A, A)$ and the existence of complements to A in G more transparent in the further exercises below.
- A stronger version of Schur–Zassenhaus Theorem asserts that if N is a normal Hall subgroup of a finite group G , then not only does N have a complement in G , but all such complements are conjugate in G . To prove this, one needs the concept of solvable groups, which we shall introduce in Section 11; see [22, pp. 246–8] for further details.

Exercises

Exercise 9.1. Suppose that G is a finite group such that $G = N \rtimes H$, where H is abelian. Show that if $|N|$ and $|H|$ are relatively coprime, then all complements to N in G are conjugate.

Solution. □

Exercise 9.2. Repeat Exercise 9.1 under the assumption that H is nilpotent rather than abelian.

Solution. □

Further Exercises

These exercises are a continuation of the further exercises to Section 2. Let N and H be groups. A *factor pair* of N by H is a pair (f, φ) of set maps $f : H \times H \rightarrow N$ and $\varphi : H \rightarrow \text{Aut}(N)$ satisfying properties **1**, **2**, and **3** listed on page 27. Let $\mathcal{E}$ be the set of extensions of N by H , and let $\mathcal{F}$ be the set of factor pairs of N by H . In what follows, we will always use “extension” to mean an element of $\mathcal{E}$, and “factor pair” to mean an element of $\mathcal{F}$.

Exercise 9.3. Let (f, φ) be a factor pair, and define

$$(x, \alpha) \cdot (y, \beta) = (x\varphi(\alpha)(y)f(\alpha, \beta), \alpha\beta)$$

for $(x, \alpha), (y, \beta) \in N \times H$. Show that this gives a group structure on $N \times H$; call this group $E_{f, \varphi}$. Show further that $(E_{f, \varphi}, i, \pi)$ is an extension, where $i(x) = (x, 1)$ and $\pi(x, \alpha) = \alpha$, and that (f, φ) is the factor pair arising from some normalized section of $E_{f, \varphi}$. (Observe that this construction generalizes the notion of external semidirect product.)

Solution. □

We have seen in Exercise 2.13 that an extension gives rise to a factor pair via a choice of normalized section, and we have just given an explicit construction of an extension from a given factor pair. We view these processes as giving maps between $\mathcal{E}$ and $\mathcal{F}$, and we now investigate the relationship between these maps. We must first consider the relation between factor pairs arising from different normalized sections of the same extension.

Exercise 9.4. (cont.) Suppose that t and u are normalized sections of an extension E , and let (f, φ) and (g, ρ) be the factor pairs arising from t and u , respectively. Let $c : H \rightarrow N$ be the set map such that $u(\alpha) = c(\alpha)t(\alpha)$ for every $\alpha \in H$. Show that the following properties hold:

4 $\rho(\alpha) = \Psi(c(\alpha))\varphi(\alpha)$ for $\alpha \in H$, where $\Psi(c(\alpha))$ is the inner automorphism of N corresponding to $c(\alpha)$.

5 $g(\alpha, \beta) = c(\alpha)\varphi(\alpha)(c(\beta))f(\alpha, \beta)c(\alpha\beta)^{-1}$ for $\alpha, \beta \in H$.

Solution. □

The above exercise motivates the following definition: We say that two factor pairs (f, φ) and (g, ρ) are *equivalent* if there is a map $c : N \rightarrow H$ such that properties **4** and **5** hold. (Verify that this is an equivalence relation on $\mathcal{F}$.) Let $\overline{\mathcal{F}}$ denote the set of equivalence classes of factor pairs; we will use $[f, \varphi]$ to denote the class of the factor pair (f, φ) . We have a well-defined map from $\mathcal{E}$ to $\overline{\mathcal{F}}$ which sends an extension to the class of a factor pair arising from any normalized section. We must now consider what happens when we pass from $\overline{\mathcal{F}}$ back to $\mathcal{E}$ via the construction in Exercise 9.3. Here we will need to recall the exact definition of an extension.

Exercise 9.5. (cont.) Let (f, φ) and (g, ρ) be factor pairs, with $[f, \varphi] = [g, \rho]$. Let $i : N \rightarrow E_{f, \varphi}$ and $j : N \rightarrow E_{g, \rho}$ be the natural inclusions (of N into the underlying set $N \times H$), and let $\pi : E_{f, \varphi} \rightarrow H$ and $\tau : E_{g, \rho} \rightarrow H$ be the natural projections (of the underlying set $N \times H$ onto H). Show that there is an isomorphism $\xi : E_{f, \varphi} \rightarrow E_{g, \rho}$ such that $\xi \circ i = j$ and $\tau \circ \xi = \pi$.

Solution. □

Motivated by the above exercise, we say that two extensions (E, i, π) and (F, j, τ) are *equivalent* if there is an isomorphism $\xi : E \rightarrow F$ such that $\xi \circ i = j$ and $\tau \circ \xi = \pi$. (Verify that this gives an equivalence relation on $\mathcal{E}$.) We let $\overline{\mathcal{E}}$ denote the set of equivalence classes of extensions, and we let $[E]$ denote the class of an extension E . In this context, Exercise 9.5 asserts that there is a well-defined map from $\overline{\mathcal{F}}$ to $\overline{\mathcal{E}}$, sending $[f, \varphi]$ to $[E_{f, \varphi}]$.

Exercise 9.6. (cont.) If p is an odd prime, show that $\mathbb{Z}_{p^2}$ can be realized in $p - 1$ nonequivalent ways as an extension of $\mathbb{Z}_p$ by $\mathbb{Z}_p$.

Solution. □

Exercise 9.7. (cont.) We have already obtained a map from $\mathcal{E}$ to $\overline{\mathcal{F}}$, sending an extension E to the class of the factor pair arising from any normalized section of E . Show that this map induces a map from $\overline{\mathcal{E}}$ to $\overline{\mathcal{F}}$.

Solution. □

Exercise 9.8. (cont.) Show that the map from $\overline{\mathcal{E}}$ to $\overline{\mathcal{F}}$ obtained in Exercise 9.7 is inverse to the map from $\overline{\mathcal{F}}$ to $\overline{\mathcal{E}}$ sending $[f, \varphi]$ to $[E_{f, \varphi}]$. Conclude that there is a bijective correspondence between the set of equivalence classes of extensions and the set of equivalence classes of factor pairs.

Solution. □

Exercise 9.8 implies that in order to study extensions up to equivalence, it suffices to study equivalence classes of factor pairs. The next two exercises give a slight refinement of the correspondence just obtained.

Exercise 9.9. (cont.) Let $\eta : \text{Aut}(N) \rightarrow \text{Out}(N)$ be the natural map. Show that if (f, φ) and (g, ρ) are any two factor pairs arising from an extension E , then $\eta \circ \varphi = \eta \circ \rho$, and this map from H to $\text{Out}(N)$ (which we denote by ψ_E) is a homomorphism. Conclude that there is a well-defined map from $\mathcal{E}$ to the set of homomorphisms from H to $\text{Out}(N)$, sending E to ψ_E .

Solution. □

Exercise 9.10. (cont.) Let $\psi : H \rightarrow \text{Out}(N)$ be a given homomorphism. Show that there is a bijective correspondence between the set of classes $[E]$ of $\overline{\mathcal{E}}$ for which $\psi_E = \psi$ and the set of classes $[f, \varphi]$ of $\overline{\mathcal{F}}$ for which $\eta \circ \varphi = \psi$, where $\eta : \text{Aut}(N) \rightarrow \text{Out}(N)$ is the natural map.

Solution. □

We now consider the case where the group N is abelian; we write A instead of N , and we will use additive notation for A . Observe that $\text{Out}(A) = \text{Aut}(A)$. We fix a homomorphism $\varphi : H \rightarrow \text{Aut}(A)$, and we write ${}^x a$ in lieu of $\varphi(x)(a)$ for $x \in H$ and $a \in A$. We would like to study those equivalence classes $[E]$ of extensions of N by H for which $\psi_E = \varphi$; we say that such extensions *respect* the action of H on A . By Exercise 9.10, it suffices to study equivalence classes $[f, \varphi]$ of factor pairs of A by H . We suppress φ in our notation, so that we are studying functions $f : H \times H \rightarrow A$ such that $f(x, 1) = f(1, x) = 0$ for all $x \in H$ and which in addition satisfy

$$f(x, y) + f(xy, z) = {}^x f(y, z) + f(x, yz)$$

for all $x, y, z \in H$, with two such functions f and g being equivalent if there is a map $c : H \rightarrow A$ such that $c(1) = 0$ and

$$g(x, y) = f(x, y) + c(x) + {}^x c(y) - c(xy)$$

for all $x, y \in H$. As discussed in the section, the set $H^2(H, A)$ of equivalence classes of such functions forms an abelian group that is called the second cohomology group of (H, A) . It now follows from Exercise 9.10 that there is a bijective correspondence between the group $H^2(H, A)$ and the set of equivalence classes of extensions of A by H which respect the action of H on A . In particular, if $H^2(H, A) = 0$, then every extension of A by H is split.

Exercise 9.11. (cont.) Suppose that H and A are both finite. Show that the order of each element of $H^2(H, A)$ divides both $|H|$ and the exponent of A . (This implies that $H^2(G/A, A) = 0$ when A is an abelian normal Hall subgroup of a finite group G , which we established in proving Schur–Zassenhaus Theorem.)

Solution. □

Chapter 4

Normal Structure

10 Composition Series

Composition Series

- A series of subgroups $G = G_0 > G_1 > \dots > G_r = 1$ of a group G is a *composition series* of G if $G_{i+1} \trianglelefteq G_i$ for every i and if each *successive quotient* G_i/G_{i+1} is simple.
- The composition series above is said to have *length* r .
- The successive quotients of a composition series are called the *composition factors* of the series.
- More generally, a group is said to be a *composition factor* of G if it is isomorphic with one of the composition factors in some composition series of G .
- Example: $S_5 > A_5 > 1$ is the only composition series of S_5 (since A_5 is simple by Theorem 7.4 and is the only non-trivial proper normal subgroup of S_5 by Theorem 7.5), with composition factors $\mathbb{Z}_2$ and A_5 .

Maximal Normal Subgroups

- N is a *maximal normal subgroup* of G if $N \trianglelefteq G$ and there is no proper normal subgroup of G that properly contains N .
- By Correspondence Theorem, N is a maximal normal subgroup of G iff G/N is simple.
- A maximal subgroup that is also normal is clearly a maximal normal subgroup (and must be of prime index), but a maximal normal subgroup need not be a maximal subgroup, for it could be properly contained in a proper subgroup that is not normal.
- For example, $A_5 \times \mathbb{Z}_2$ has $1 \times \mathbb{Z}_2$ as a maximal normal subgroup that is not maximal.

Proposition 10.1. *Finite groups have composition series.*

Proof.

□

- Infinite groups need not have composition series. For example, using Theorem 1.5 we see that every non-trivial subgroup of the infinite cyclic group $\mathbb{Z}$ is isomorphic with $\mathbb{Z}$; as $\mathbb{Z}$ is not simple, $\mathbb{Z}$ has no simple subgroups, and hence we cannot construct a composition series of $\mathbb{Z}$, since the last non-trivial term of such a series must be a simple subgroup of $\mathbb{Z}$.

Lemma 10.2. *Let G be a group that has a composition series, and let $N \trianglelefteq G$. Then N has a composition series.*

Proof.

□

Equivalence of Composition Series

- Let $G = G_0 > G_1 > \dots > G_r = 1$ be a composition series, and suppose $G = H_0 > H_1 > \dots > H_r = 1$ is another composition series of the same length r . These series are *equivalent* if there is some $\rho \in S_r$ such that $G_{i-1}/G_i \cong H_{\rho(i)-1}/H_{\rho(i)}$ for every i .
- Example: $G = \langle x \rangle \cong \mathbb{Z}_6$ with $G_1 = \langle x^2 \rangle$, $H_1 = \langle x^3 \rangle$ gives equivalent composition series $G > G_1 > 1$ and $G > H_1 > 1$, since $G/G_1 \cong H_1/1 \cong \mathbb{Z}_2$ and $G_1/1 \cong G/H_1 \cong \mathbb{Z}_3$. (Here $\rho = (1\ 2) \in S_2$.)
- Up to equivalence, a group has at most one composition series, so a group having a composition series has a well-defined collection of composition factors.
- Since composition factors are simple groups, classifying finite simple groups (completed in 1980) gives comprehensive understanding of finite groups.

Theorem (Jordan–Hölder Theorem). *Suppose that G is a group that has a composition series. Then any two composition series of G have the same length and are equivalent.*

Proof.

□

Subnormal, Normal, and Chief Series

- A series of subgroups $G = G_0 \geq G_1 \geq \dots \geq G_r = 1$ of a group G is called a *subnormal series* of G if $G_{i+1} \trianglelefteq G_i$ for every i .
- A subnormal series is called a *normal series* if $G_i \trianglelefteq G$ for every i .
- Some authors use “normal series” for what we call a subnormal series, and “invariant series” for what we call a normal series.
- Composition series are examples of subnormal series, but subnormal series can have successive quotients that are trivial or that are non-trivial but not simple.
- Two subnormal series of the same length are said to be *equivalent* if they satisfy the condition stated as the definition of equivalence of composition series.
- A *refinement* of a subnormal series is one obtained by interposing additional terms; it is *proper* if at least one of the interposed terms was not already present.
- A composition series is a subnormal series that has no repeated terms and does not admit a proper refinement.
- A *chief series* of G is a normal series of G with no repeated terms and with the additional property that no normal subgroup of G is contained properly between any two terms of the series.
- Analogy: A composition series is a subnormal series having no subnormal series as a proper refinement, and a chief series is a normal series having no normal series as a proper refinement.
- The successive quotients of a chief series are its *chief factors*; a group is a *chief factor* of G if it is isomorphic with a chief factor in some chief series of G .
- The analogue of the Jordan–Hölder Theorem for chief series holds—namely, any two chief series of a group have the same length and are equivalent, with a proof virtually identical to that for composition series. This is a special case of the Jordan–Hölder theorem for groups with operators (see [26, Section 2.3]).

Minimal Normal Subgroups

- N is a *minimal normal subgroup* of G if $1 \neq N \trianglelefteq G$ and there is no non-trivial normal subgroup of G that is properly contained in N .
- Any non-trivial finite group has minimal normal subgroups.
- A simple group has a unique minimal normal subgroup, namely itself.

Proposition 10.3. *Finite groups have chief series.*

Proof. □

- By definition, the composition factors of finite groups are simple groups. We now turn to determining the nature of chief factors of finite groups; the first step is to rephrase the problem in different terms.

Lemma 10.4. *Let G be a group having a chief series. Then every chief factor of G is a minimal normal subgroup of a quotient group of G .*

Proof. □

Theorem 10.5. *A minimal normal subgroup of a finite group is a direct product of mutually isomorphic simple groups.*

Proof. □

Corollary 10.6. *A chief factor of a finite group is a direct product of mutually isomorphic simple groups.*

Proof. □

Exercises

Exercise 10.1. Show that an abelian group has a composition series iff it is finite.

Solution. □

Exercise 10.2. Show that $\text{GL}(n, F)$ has a composition series iff F is finite.

Solution. □

Exercise 10.3. Using Third Isomorphism Theorem (Exercise 2.14), prove the *Schreier refinement theorem*: Any two subnormal series of a given group have equivalent refinements.

Solution. □

Exercise 10.4. (cont.) Use the Schreier refinement theorem to give an alternate proof of Jordan–Hölder Theorem.

Solution. □

Exercise 10.5. Determine all composition series and all chief series of S_n for $n \geq 2$. (HINT: Use Exercise 3.8.)

Solution. □

Exercise 10.6. Show that any group having a composition series has a chief series.

Solution. □

Exercise 10.7. (cont.) Show that any chief factor of a group having a composition series is a direct product of mutually isomorphic simple groups.

Solution. □

Exercise 10.8. Show that any finite group that has no proper non-trivial characteristic subgroups is a direct product of mutually isomorphic simple groups. Use this to give a new proof of Theorem 10.5.

Solution. □

11 Solvable Groups

The derived series and solvability

- We define a series $G^{(k)}$ of subgroups of a group G by setting $G^{(0)} = G$ and taking $G^{(k)}$ to be the derived group of $G^{(k-1)}$ for $k \in \mathbb{N}$. This series is called the *derived series* of G .
- Since $G^{(k+1)}$ is a characteristic subgroup of $G^{(k)}$ for each k by Lemma 2.5, we see via Lemma 2.4 that the derived series is a normal series of G . Moreover, we see from Proposition 2.6 that each successive quotient $G^{(k)}/G^{(k+1)}$ of the derived series is abelian.
- We say that a group is *solvable* if its derived series terminates in the identity. (Authors from British Commonwealth countries often write “soluble” in lieu of “solvable” and “insoluble” in lieu of “non-solvable.”)
- Since a group is abelian iff its derived group is trivial, we see that abelian groups are solvable. However, not all groups are solvable; for example, A_5 is non-abelian and simple by Theorem 7.4, so $A_5^{(k)} = A_5$ for all k .
- A group whose derived group is itself is said to be *perfect*; any non-abelian simple group is perfect, and in particular is not solvable. However, a perfect group need not be simple: $\text{SL}(n, F)$ is perfect but has non-trivial center, except when $n = 2$ and $|F| = 3$.
- Solvable groups are to be thought of as the opposite of simple groups: simple groups have very few normal subgroups while solvable groups are rife with them.

Proposition 11.1. *A simple solvable group has prime order.*

Proof. □

Proposition 11.2. *The following statements about a group G are equivalent:*

- (1) G is solvable.
- (2) G has a normal series in which every successive quotient is abelian.
- (3) G has a subnormal series in which every successive quotient is abelian.

Proof. □

Proposition 11.3. (i) *If G is solvable and $H \leq G$, then H is solvable.*

(ii) *If G is solvable and $N \trianglelefteq G$, then G/N is solvable.*

(iii) *If $N \trianglelefteq G$ and both N and G/N are solvable, then G is solvable.*

(iv) *If G and H are solvable, then $G \times H$ is solvable.*

Proof. □

Proposition 11.4. *A group having a composition series is solvable iff all of its composition factors have prime order. (In particular, such groups are finite.)*

Proof. □

Examples and consequences

- There are groups, such as S_5 , which are neither simple nor solvable, and there are non-abelian groups, such as S_3 (which has the composition series $S_3 > A_3 > 1$), which are solvable.
- Proposition 11.4 also implies that an infinite abelian group, being solvable, cannot possess a composition series (which was Exercise 10.1).

Corollary 11.5. *Finite p -groups are solvable.*

Proof. □

Proposition 11.6. *A group having a composition series is solvable iff all of its chief factors are elementary abelian.*

(Any group having a composition series also has a chief series by Exercise 10.6. Alternately, a solvable group having a composition series is finite by Proposition 11.4 and hence has a chief series by Proposition 10.3.)

Proof. □

Corollary 11.7. *A group having a composition series is solvable iff it has a normal series in which every successive quotient is a p -group.*

Proof. □

Supersolvable groups and Hall's theorems

- Those finite groups whose chief factors all have prime order are said to be *supersolvable*; finite supersolvable groups are solvable by Proposition 11.6.
- It follows from Exercise 8.2 that finite p -groups are supersolvable. However, not all finite solvable groups are supersolvable. The series $S_4 > A_4 > K > 1$, where K is the Klein four-group, is a chief series of S_4 , and we have $S_4/A_4 \cong \mathbb{Z}_2$, $A_4/K \cong \mathbb{Z}_3$, and $K \cong \mathbb{Z}_2 \times \mathbb{Z}_2$, which by Proposition 11.6 shows that S_4 is solvable but not supersolvable.
- There is an equivalent definition of supersolvability which extends to infinite groups; see Exercise 11.6.
- Solvable groups possess a number of properties beyond those which hold for arbitrary groups. The following theorem, due to Philip Hall, is a generalization for finite solvable groups of part (i) of Sylow's Theorem.

Theorem 11.8. *Let G be a finite solvable group of order mn , where m and n are coprime. Then G has a subgroup of order m .*

(In other words, this theorem asserts that a finite solvable group possesses Hall subgroups of all possible orders.)

Proof. □

More on solvability criteria

- Theorem 11.8 does not hold for all finite groups; for instance, we observed in the proof of Theorem 7.5 that the group A_5 of order $60 = 20 \cdot 3$ has no subgroup of order 20.
- P. Hall also proved the following generalization for finite solvable groups of parts (ii) and (iii) of Sylow's Theorem, which the reader is asked to prove in the exercises.

Theorem 11.9. *Let G be a finite solvable group of order mn , where m and n are coprime. Then any two subgroups of G of order m are conjugate, and any subgroup of G whose order divides m is contained in a subgroup of order m .*

Proof. □

Classical theorems on solvability

- We now state, without proof, a number of well-known theorems giving conditions under which a finite group is solvable. We start with a classical result that generalizes Corollary 11.5.

Theorem (Burnside's Theorem). *If p and q are primes, then any group of order $p^a q^b$ is solvable.*

Proof. □

- Burnside's Theorem is the best possible result, in the sense that a finite group whose order has exactly three prime divisors need not be solvable, with A_5 providing a counterexample.

- Burnside proved this theorem in 1904 using the methods of character theory, which will be discussed in Chapter 6; we present a character-theoretic proof of Burnside's theorem in the Appendix (Burnside's Theorem). Until the 1960s, there was no proof of this theorem that did not involve character theory.

Theorem (Feit–Thompson Theorem). *All finite groups of odd order are solvable.*

Proof. □

- This result, also known as the “odd order theorem,” had first been conjectured by Burnside in 1911. As a corollary, we see that the only simple groups of odd order are cyclic groups of prime order.
- Once it had finally been established that all non-abelian finite simple groups are of even order, the movement toward a classification of all finite simple groups gathered steam. Feit and Thompson completed their proof of this theorem during a special “Finite Group Theory Year,” held at the University of Chicago in the school year 1960–61. Their original proof [12] was 255 pages long, and even at that length the proof requires too much background knowledge for it to be readily comprehensible to anyone but the specialists at the time at which the proof was published. In recent years there has been a program, of which [8] represents a major part, to produce a more accessible proof of this fundamental result.
- The following result is a converse to Theorem 11.8 and is also due to P. Hall. While this theorem generalizes Burnside's Theorem, its proof relies on the fact that groups of order $p^a q^b$ are solvable.

Theorem 11.10. *Let G be a finite group. If G has a subgroup of order m whenever m and n are coprime numbers such that $|G| = mn$, then G is solvable.*

Proof. □

- This next result was conjectured by P. Hall in the same paper [14] in which he proved Theorem 11.10; it was later proved by Thompson in [27].

Theorem 11.11. *A finite group G is not solvable iff there exist non-trivial elements x, y, z of G of pairwise coprime orders a, b, c such that $xy = z$.*

Proof. □

- For example, in A_5 this criterion for non-solvability is satisfied by taking $a = 5$, $b = 2$, $c = 3$ and $x = (1\ 2\ 3\ 4\ 5)$, $y = (1\ 2)(3\ 4)$, $z = (1\ 3\ 5)$. (Compare with Exercise 1.5.)

Galois's theorem and the affine group

- The concept of solvable groups originated with Galois around 1830 in his work on the solution by radicals of polynomial equations; this in fact is the origin of the term “solvable” for this class of groups.
- Let V be a vector space over a field F . For $v \in V$, let T_v be the self-map of V corresponding to translation by v , and let $\mathcal{T}(V)$ be the set of all such translations; this is a group isomorphic with the abelian group V , and if we regard $\mathcal{T}(V)$ and $\text{GL}(V)$ as subgroups of the group of all invertible self-maps of V , then $\text{GL}(V) \cap \mathcal{T}(V)$ is trivial.
- We easily verify that $ST_vS^{-1} = T_{S(v)}$ for any $S \in \text{GL}(V)$ and $v \in V$. This gives rise to a homomorphism from $\text{GL}(V)$ to $\text{Aut}(\mathcal{T}(V))$, and hence we have an (internal) semidirect product $\mathcal{T}(V) \rtimes \text{GL}(V)$. This group is called the *affine group* of V and is denoted $\text{Aff}(V)$; its elements are called the *affine transformations* of V .

Theorem 11.12. *Let G be a finite, solvable, primitive permutation group on a set X . Then X can be given the structure of a vector space over the field $\mathbb{Z}/p\mathbb{Z}$ for some prime p in such a way that G is isomorphic with a subgroup of $\text{Aff}(X)$ that contains $\mathcal{T}(X)$.*

Proof. □

Nilpotent groups

- A normal series $G = G_0 \geq G_1 \geq \dots \geq G_r = 1$ of a group G is said to be a *central series* of G if, for each i , G_i/G_{i+1} is contained in the center of G/G_{i+1} .
- A group G is said to be *nilpotent* if it has a central series. An abelian group G has the central series $G > 1$, and hence abelian groups are nilpotent.

Proposition 11.13. *Nilpotent groups are solvable.*

Proof.

□

- There are solvable groups that are not nilpotent. For example, the group S_3 cannot have a central series, as the penultimate term of such a series would have to be a non-trivial subgroup of $Z(S_3) = 1$.

Lemma 11.14. *Finite p -groups are nilpotent.*

Proof.

□

- If H and K are subgroups of a group G , we define a new subgroup $[H, K]$ of G by $[H, K] = \langle \{[h, k] \mid h \in H, k \in K\} \rangle$. Observe that $G' = [G, G]$.
- We now reconcile the above definition of nilpotence with that made in Section 8 for finite groups. Recall that a subgroup H of a group G is said to be *self-normalizing* if $N_G(H) = H$.

Theorem 11.15. *The following statements about a finite group G are equivalent:*

- (1) G is nilpotent.
- (2) G has no proper self-normalizing subgroups.
- (3) Every Sylow subgroup of G is normal in G .
- (4) G is the direct product of its Sylow subgroups.
- (5) Every maximal subgroup of G is normal in G .

Proof.

□

Exercises

Exercise 11.1. Show that any finite group has a largest solvable normal subgroup.

Solution.

□

Exercise 11.2. Let N be a solvable normal Hall subgroup of a finite group G . Show that any two complements to N in G are conjugate. (The strong version of Schur–Zassenhaus Theorem asserts that this is true for any normal Hall subgroup N , whether solvable or not. The proof uses Feit–Thompson Theorem to argue that if N is not solvable, then G/N must be solvable.)

Solution.

□

Exercise 11.3. Complete the following sketch, which gives a proof of Theorem 11.8 that is independent of Schur–Zassenhaus Theorem. Here G is a finite solvable group of order mn , where m and n are coprime, and we are to show that G has a subgroup of order m .

- Using induction, reduce to the case where G has a unique minimal normal subgroup N of order n .
- Let M/N be a minimal normal subgroup of G/N ; since G/N is solvable by Proposition 11.3, it follows from Proposition 11.6 that $|M/N|$ is a power of some prime divisor p of m . If P is a Sylow p -subgroup of M , show that $|N_G(P)| = m$.

Solution.

□

Exercise 11.4. Prove Theorem 11.9. If your proof involves Schur–Zassenhaus Theorem, then construct a second proof that is independent of Schur–Zassenhaus by mimicking Exercise 11.3 above.

Solution. □

Exercise 11.5. Let V be a finite-dimensional vector space over the field of p elements. Suppose that G is a solvable subgroup of $\text{Aff}(V)$ which contains $\mathcal{T}(V)$ and that the stabilizer of the origin under the action of G on V does not leave any non-zero proper subspace of V invariant. Show that G is a primitive permutation group on V .

Solution. □

Exercise 11.6. Show that a finite group is supersolvable iff it has a normal series having cyclic successive quotients. (An arbitrary group is called supersolvable if this latter condition holds.)

Solution. □

Exercise 11.7. Show that finite nilpotent groups are supersolvable. (If we define supersolvability for arbitrary groups as in Exercise 11.6, is an infinite nilpotent group necessarily supersolvable?)

Solution. □

Exercise 11.8. (cont.) Give an example of a finite supersolvable group that is not nilpotent. (Hence we have a proper ascending chain of classes of finite groups, starting with cyclic groups, then abelian, then nilpotent, then supersolvable, and ending with solvable.)

Solution. □

Further Exercises

Let G be a group. Let $\Gamma_1 = G$, and for $n \in \mathbb{N}$ let $\Gamma_{n+1} = [\Gamma_n, G] \leq G$. Let $Z_0 = 1$, and for $n \in \mathbb{N}$ let Z_n be the unique subgroup of G such that $Z_n/Z_{n-1} = Z(G/Z_{n-1})$. (Observe that $\Gamma_2 = G'$ and $Z_1 = Z(G)$.) We call the series $G = \Gamma_1 \geq \Gamma_2 \geq \Gamma_3 \geq \dots$ the *lower central series* of G and the series $1 = Z_0 \leq Z_1 \leq Z_2 \leq \dots$ the *upper central series* of G .

Exercise 11.9. Show that each Γ_n and Z_n is a characteristic subgroup of G .

Solution. □

Exercise 11.10. (cont.) Show that G is nilpotent iff $\Gamma_r = 1$ for some r iff $Z_s = G$ for some s .

Solution. □

Exercise 11.11. (cont.) Suppose that G is nilpotent. Show that the lower and upper central series of G have the same length, say c , and that no central series of G can have length less than c . (This number c is called the *nilpotency class* of G . The groups of nilpotency class 1 are exactly the abelian groups.)

Solution. □

Chapter 5

Semisimple Algebras

12 Modules and Representations

From Group Actions to Linear Actions

- In Section 3 we studied actions of groups on sets; if a group acts on a set carrying additional algebraic structure, it is the structure-respecting actions that are most interesting.
- Here we focus on actions of groups on vector spaces that respect the vector space structure.
- Let F be a field and let G be a group acting on an F -vector space V . The action is *linear* if
 - $g(v + w) = gv + gw$ for all $g \in G$ and $v, w \in V$;
 - $g(\alpha v) = \alpha(gv)$ for all $g \in G$, $\alpha \in F$, and $v \in V$.
- Proposition 3.1 gave the bijection between group actions on a set and homomorphisms to symmetric groups; we now deduce its analogue for linear actions.

Proposition 12.1. *There is a bijective correspondence between the set of linear actions of a group G on an F -vector space V and the set of homomorphisms from G to $\text{GL}(V)$.*

Proof.

□

Linear Representations and Modules

- A homomorphism $\rho: G \rightarrow \text{GL}(V)$, where G is a group and V a vector space, is called a *linear representation* of G in V .
- By Proposition 12.1, the study of linear representations is equivalent to the study of linear actions; with emphasis on finite groups and finite-dimensional vector spaces, this area originated in the late nineteenth century.
- The modern approach to the representation theory of finite groups involves yet another equivalent concept, that of finitely generated modules over group algebras. It is therefore necessary at this juncture to review some elementary module theory.
- As in Section 1, we will omit most proofs on the assumption that the reader will have seen this material previously; however, our account requires only a mild knowledge of rings, fields, and vector spaces.
- Let R be a ring with unit, and let M be an abelian group written additively. Then M is a *left R -module* if there is a map $R \times M \rightarrow M$, $(r, m) \mapsto rm$, satisfying:
 - $1m = m$ for all $m \in M$;
 - $r(m + n) = rm + rn$ for all $r \in R$ and $m, n \in M$;
 - $(r + s)m = rm + sm$ for all $r, s \in R$ and $m \in M$;
 - $r(sm) = (rs)m$ for all $r, s \in R$ and $m \in M$.

- If $R = \mathbb{Z}$, then $\mathbb{Z}$ -modules are exactly abelian groups; if F is a field, then F -modules are precisely F -vector spaces. A module is the natural generalisation of a vector space to an arbitrary ring.
- One similarly defines right R -modules; if R is commutative, every left module is canonically a right module. In general, a left R -module is a right module over the *opposite ring* R^{op} .
- Let S be another ring with unit. An abelian group M that is both a left R -module and a right S -module is an (R, S) -bimodule provided $r(ms) = (rm)s$ for all $r \in R, m \in M, s \in S$. Any left R -module is an $(R, \mathbb{Z})$ -bimodule, and R itself is an (R, R) -bimodule.
- An R -module M is *finitely generated* if every element is an R -linear combination of elements from some finite subset of M . Unlike vector spaces, minimal generating sets need not have a common size.
- If N is a subgroup of M closed under the R -action, N is an R -submodule. The submodules of the regular module R are exactly the left ideals. A nonzero module with no submodules other than 0 and itself is *simple*.
- Quotient modules M/N are defined by $r(m + N) = rm + N$.
- Sums, intersections, internal and external direct sums of submodules are defined as for groups; N is a *direct summand* of M if $M = N \oplus N'$ for some submodule N' . We write nM or M^n for $\underbrace{M \oplus \cdots \oplus M}_n$.
- A *composition series* of M is a descending series of submodules ending at 0 in which each successive quotient is simple. Modules need not have one (any infinite $\mathbb{Z}$ -module is an example), but when they do, Jordan–Hölder Theorem holds.
- Every composition factor of any submodule or quotient module of an R -module M must also be a composition factor of M , as any composition series of a submodule (resp., quotient module) can be extended (resp., lifted) to give a composition series of M .
- An R -module homomorphism $\varphi: M \rightarrow N$ is a group homomorphism with $\varphi(rm) = r\varphi(m)$ for all $r \in R, m \in M$. Mono-, epi-, iso-, endo-, automorphisms are defined as for groups; the kernel is a submodule, the image is a submodule, and Fundamental Theorem on Homomorphisms, Correspondence Theorem, and the two isomorphism theorems (First Isomorphism Theorem and Second Isomorphism Theorem) all carry over.

Lemma (Schur’s Lemma). *Any non-zero homomorphism between simple R -modules is an isomorphism.*

Proof. □

Hom-Modules, Balanced Maps, and Tensor Products

- Given R -modules M and N , we write $\text{Hom}_R(M, N)$ for the set of R -module homomorphisms $M \rightarrow N$, and $\text{End}_R(M)$ for $\text{Hom}_R(M, M)$. Pointwise addition makes $\text{Hom}_R(M, N)$ an abelian group.
- If S is another ring, M an (R, S) -bimodule, and N an R -module, then $\text{Hom}_R(M, N)$ is an S -module via $(s\varphi)(m) = \varphi(ms)$. In particular, when F is a field and U, V are F -vector spaces, $\text{Hom}_F(U, V)$ is an F -vector space.
- The map $\text{Hom}_R(R, M) \rightarrow M$ sending φ to $\varphi(1)$ is an R -module isomorphism.
- Let M be an (R, S) -bimodule and N a left S -module. A map $f: M \times N \rightarrow U$ to an R -module U is *balanced* if
 - $f(m_1 + m_2, n) = f(m_1, n) + f(m_2, n);$
 - $f(m, n_1 + n_2) = f(m, n_1) + f(m, n_2);$
 - $f(ms, n) = f(m, sn);$
 - $f(rm, n) = rf(m, n).$

- The *tensor product* $M \otimes_S N$ is the R -module equipped with a balanced map $\eta: M \times N \rightarrow M \otimes_S N$ such that any balanced $f: M \times N \rightarrow U$ factors uniquely through η : there is a unique R -module homomorphism $\alpha: M \otimes_S N \rightarrow U$ with $f = \alpha \circ \eta$. We write $m \otimes n = \eta(m, n)$.
- Concretely, $M \otimes_S N$ is generated by the symbols $\{m \otimes n \mid m \in M, n \in N\}$ subject to the relations
 - $(m_1 + m_2) \otimes n = m_1 \otimes n + m_2 \otimes n$;
 - $m \otimes (n_1 + n_2) = m \otimes n_1 + m \otimes n_2$;
 - $(ms) \otimes n = m \otimes (sn)$;
 - $(rm) \otimes n = r(m \otimes n)$.

A general element of $M \otimes_S N$ is a *sum* of elementary tensors, not just one.

- Special case: if F is a field and U, V are finite-dimensional with bases $\{u_1, \dots, u_r\}$ and $\{v_1, \dots, v_s\}$, then $U \otimes_F V$ has basis $\{u_i \otimes v_j\}$ and dimension rs . For $u = \sum_i a_i u_i$ and $v = \sum_j b_j v_j$, $u \otimes v = \sum_{i,j} a_i b_j (u_i \otimes v_j)$.
- (An understanding of this special case will, strictly speaking, suffice in reading the remainder of the book; however, an understanding of the general case will be beneficial in Section 16.)

Proposition 12.2. *Let R be a ring with unit, and let M be an R -module. Then M and $R \otimes_R M$ are isomorphic R -modules.*

Proof. □

Proposition 12.3. *Let R and S be rings with unit, let $M_1, \dots, M_r$ be (R, S) -bimodules, and let N be an S -module. Then $(\bigoplus_i M_i) \otimes_S N$ and $\bigoplus_i M_i \otimes_S N$ are isomorphic R -modules.*

Proof. □

Proposition 12.4. *Let F be a field, and let U and V be F -vector spaces, with $\dim_F(U) < \infty$. Let $U^* = \text{Hom}_F(U, F)$. Then the map $\Gamma: U^* \otimes_F V \rightarrow \text{Hom}_F(U, V)$ defined by setting $\Gamma(\varphi \otimes v)(u) = \varphi(u)v$ and extending linearly is an isomorphism of F -vector spaces.*

Proof. □

Group Rings and Group Algebras

- Let R be a ring and G a group. The *group ring* of G over R , denoted RG , consists of all finite formal R -linear combinations of elements of G , with the obvious addition and with multiplication extending that in G :

$$\left(\sum_{x \in G} \alpha_x x \right) \left(\sum_{y \in G} \beta_y y \right) = \sum_{x \in G} \sum_{y \in G} \alpha_x \beta_y xy = \sum_{x \in G} \left(\sum_{g \in G} \alpha_g \beta_{g^{-1}x} \right) x.$$

- The unit element is the identity of G .
- When $R = F$ is a field and G is finite, FG is also an F -vector space with basis G , hence has finite dimension $|G|$. In this case FG is called the *group algebra*.
- An *algebra* over F (or *F -algebra*) is a set A carrying both a ring structure and an F -vector space structure that share the same addition, with $(\lambda a)b = a(\lambda b) = \lambda(ab)$ for $\lambda \in F, a, b \in A$. We do *not* assume an algebra has a ring unit.
- An algebra is *finite-dimensional* if it has finite dimension as an F -vector space; the matrix ring $M_n(F)$ is a finite-dimensional F -algebra, and FG is finite-dimensional whenever G is finite.
- A homomorphism of F -algebras is a ring homomorphism that is also F -linear.

Lemma 12.5. *If F is a field and G a finite group, then an FG -module is finitely generated iff it has finite dimension as an F -vector space.*

Proof. □

Proposition 12.6. *If F is a field and G a finite group, then there is a bijective correspondence between finitely generated FG -modules and linear actions of G on finite-dimensional F -vector spaces.*

Proof. □

Examples and Constructions of FG -Modules

- In what follows G is a finite group, F a field, and all F -vector spaces are finite-dimensional; all FG -modules will be finitely generated and hence finite-dimensional by Lemma 12.5.
- It is enough to specify the action of G on V ; the FG -action then extends linearly.
- *Trivial module*: F regarded as an FG -module via $g\lambda = \lambda$.
- *Permutation module*: if G acts on a finite set $X = \{x_1, \dots, x_n\}$, the F -vector space FX with basis X is an FG -module via $g(\sum_i c_i x_i) = \sum_i c_i(gx_i)$.
- Direct sums of FG -modules: $g(u, v) = (gu, gv)$.
- *Tensor product* $U \otimes_F V$ with the diagonal action $g(u \otimes v) = gu \otimes gv$.
- $\text{Hom}_F(U, V)$ with action $(g\varphi)(u) = g(\varphi(g^{-1}u))$; one checks $(g_1 g_2)\varphi = g_1(g_2\varphi)$.
- Taking $V = F$ trivial gives the *dual module* $U^* = \text{Hom}_F(U, F)$ with $(g\varphi)(u) = \varphi(g^{-1}u)$.

Proposition 12.7. *Let U and V be FG -modules. Then $U^* \otimes_F V$ and $\text{Hom}_F(U, V)$ are isomorphic FG -modules.*

Proof. □

Maschke's Theorem and Semisimplicity

- We now reach the most basic result of the representation theory of finite groups, discovered in 1898 by Heinrich Maschke at the University of Chicago.

Theorem (Maschke's Theorem). *Let G be a group, and suppose that the characteristic of F is either zero or coprime to $|G|$. If U is an FG -module and V is an FG -submodule of U , then V is a direct summand of U as FG -modules.*

Proof. □

- We summarise the hypothesis by saying that the characteristic of F *does not divide* $|G|$, even though this is a slight abuse.
- A module is *semisimple* if it is a direct sum of simple modules.

Corollary 12.8. *Let G be a group, and let F be a field whose characteristic does not divide $|G|$. Then every non-zero FG -module is semisimple.*

Proof. □

- In Chapter 6 we shall be concerned only with the case $F = \mathbb{C}$, called *ordinary* representation theory; since $\mathbb{C}$ has characteristic zero, by Corollary 12.8 every non-zero $\mathbb{C}G$ -module is semisimple.
- The next section concentrates on algebras with this property. The case where the characteristic of F divides $|G|$, in which FG -modules need not be semisimple, is the subject of *modular* representation theory (originated by Dickson in 1907; mostly developed by Brauer from the 1930s onward).

Exercises

Throughout these exercises, let F be a field and let G be a finite group.

Exercise 12.1. Let U and V be FG -modules having the same dimension n , and let $\rho: G \rightarrow \text{GL}(U)$ and $\tau: G \rightarrow \text{GL}(V)$ be the corresponding representations. By fixing F -bases for U and V , we consider ρ and τ as homomorphisms from G to $\text{GL}(n, F)$. Show that $U \cong V$ iff there exists some $M \in \text{GL}(n, F)$ such that $\rho(g)M = M\tau(g)$ for every $g \in G$. (Two representations are said to be *equivalent* if this latter condition holds; hence two representations are equivalent iff their corresponding modules over the group algebra are isomorphic.)

Solution. □

Exercise 12.2. Let $\sigma = \sum_{g \in G} g \in FG$. Show that the subspace $F\sigma$ of FG is the unique submodule of FG that is isomorphic with the trivial module.

Solution. □

Exercise 12.3. (cont.) Let $\epsilon: FG \rightarrow F$ be the FG -module epimorphism defined by $\epsilon(g) = 1$ for all $g \in G$, and let $\Delta = \ker \epsilon$. (We call Δ the *augmentation ideal* of FG .) Show that Δ is the unique submodule of FG whose quotient is isomorphic with the trivial module.

Solution. □

Exercise 12.4. (cont.) Suppose that the characteristic of F divides $|G|$. Show that $F\sigma \subseteq \Delta$. Conclude that Δ is not a direct summand of FG and hence that the FG -module FG is not semisimple. (This shows that the converse of Corollary 12.8 is true.)

Solution. □

Further Exercises

In Section 9 we defined the second cohomology group of a pair (H, A) , where H is a group and A an abelian group, with respect to a given homomorphism from H to $\text{Aut}(A)$. It is easy to verify that, for an abelian group A , specifying a homomorphism from a group H to $\text{Aut}(A)$ is exactly the same as specifying a $\mathbb{Z}H$ -module structure on A . Therefore, it is customary to talk about the second cohomology group of a pair (H, A) where H is a group and A a $\mathbb{Z}H$ -module. (We anticipated this development by writing xa instead of $\varphi(x)(a)$ on page 86.) We will now generalize the second cohomology group.

Let H be a group, let H^n be the set of n -tuples of elements of H for some $n \in \mathbb{N}$, and let A be a $\mathbb{Z}H$ -module. A *normalized n -cochain* of (H, A) is a set function $f: H^n \rightarrow A$ such that $f(h_1, \dots, h_n) = 0$ whenever $h_i = 1$ for some i ; the set of normalized n -cochains of (H, A) is denoted $C^n(H, A)$. We give $C^n(H, A)$ an abelian group structure by defining

$$(f + g)(h_1, \dots, h_n) = f(h_1, \dots, h_n) + g(h_1, \dots, h_n).$$

We define a homomorphism $\delta^n: C^{n-1}(H, A) \rightarrow C^n(H, A)$ by

$$(\delta^n f)(h_1, \dots, h_n) = h_1 f(h_2, \dots, h_n) + \sum_{i=1}^{n-1} (-1)^i f(h_1, \dots, h_i h_{i+1}, \dots, h_n) + (-1)^n f(h_1, \dots, h_{n-1}).$$

The image of δ^n is called the set of *n -coboundaries* and is denoted $B^n(H, A)$; the kernel of δ^{n+1} is called the set of *n -cocycles* and is denoted $Z^n(H, A)$. For example, $\ker \delta^3$ consists of all normalized 2-cochains f satisfying $h_1 f(h_2, h_3) - f(h_1 h_2, h_3) + f(h_1, h_2 h_3) - f(h_1, h_2) = 0$ for all $h_1, h_2, h_3 \in H$, which agrees with our definition of a 2-cocycle in Section 9. Both $B^n(H, A)$ and $Z^n(H, A)$ are abelian subgroups of $C^n(H, A)$. We find that $\delta^{n+1} \circ \delta^n$ is the zero map, or equivalently that $B^n(H, A) \subseteq Z^n(H, A)$. (Verify this.) The quotient group $Z^n(H, A)/B^n(H, A)$ is denoted $H^n(H, A)$ and is called the *n th cohomology group* of (H, A) .

While $H^3(H, A)$ plays some role in the theory of extensions of a non-abelian group N by H , where $Z(N) = A$ (see [20, pp. 124–131]), the groups $H^n(H, A)$ for $n > 3$ have no known group-theoretic meaning. Nonetheless, higher cohomology groups are studied for their own sake, and in recent years connections between the cohomology of groups and the representation theory of groups have attracted much attention. (As the title of one of the first author's papers [6] proclaims, “Cohomology is representation theory.”) We developed the group-theoretic meaning of $H^2(H, A)$ in the further exercises to Section 9, and in what follows we develop the meaning of $H^1(H, A)$.

Exercise 12.5. (cont.) Define a homomorphism $\varphi: H \rightarrow \text{Aut}(A)$ by $\varphi(x)(a) = xa$. Show that there is a bijective correspondence between the set of splitting maps of $A \rtimes_\varphi H$ and $Z^1(H, A)$. (Recall that a homomorphism $t: H \rightarrow A \rtimes_\varphi H$ is called a splitting map if $\eta \circ t$ is the identity map on H , where $\eta: A \rtimes_\varphi H \rightarrow H$ is the natural map. The natural inclusion of H into $A \rtimes_\varphi H$ is a splitting map, but there may be others.)

Solution. □

Exercise 12.6. (cont.) We say that two splitting maps t and u of $A \rtimes_{\varphi} H$ are *conjugate* if there is some $a \in A$ such that $u(x) = at(x)a^{-1}$ for all $x \in H$. (Verify that this is an equivalence relation on the set of splitting maps.) Show that there is a bijective correspondence between the set of conjugacy classes of splitting maps of $A \rtimes_{\varphi} H$ and $H^1(H, A)$.

Solution.

□

13 Wedderburn Theory

Setting and Goal

- Maschke (Corollary 12.8) shows that if $\text{char}(F) \nmid |G|$ then every finitely generated FG -module decomposes as a direct sum of simple modules.
- This section identifies the finite-dimensional F -algebras with this property and uses the resulting structure theorems to finish Kolchin's theorem on unipotent subgroups.
- Throughout, all algebras are finite-dimensional F -algebras (with unit unless stated), and all modules are finitely generated, equivalently finite-dimensional as F -vector spaces (Lemma 12.5).
- The classification is due to J. H. M. Wedderburn.

Semisimplicity for Modules

- Three equivalent formulations of semisimplicity for an A -module: every submodule is a direct summand; the module is a direct sum of simple submodules; the module is a sum of simple submodules.
- Semisimplicity is preserved under submodules and quotient modules.
- An algebra A is *semisimple* if every non-zero A -module is semisimple; equivalently, A is semisimple as an A -module over itself.

Lemma 13.1. *The following statements about an A -module M are equivalent:*

1. *Any submodule of M is a direct summand of M .*
2. *M is semisimple.*
3. *M is a sum (but not, a priori, a direct sum) of simple submodules.*

Proof.

□

Lemma 13.2. *Submodules and quotient modules of semisimple modules are semisimple.*

Proof.

□

Lemma 13.3. *The algebra A is semisimple iff the A -module A is semisimple.*

Proof.

□

Counting Simple Summands

- Decomposing $A \cong S_1 \oplus \cdots \oplus S_r$ into simple summands captures all isomorphism types of simple A -modules.
- Multiplicities of simple summands in a semisimple module are determined uniquely (Jordan–Hölder Theorem).

Proposition 13.4. *Let A be a semisimple algebra, and suppose that as A -modules we have $A \cong S_1 \oplus \cdots \oplus S_r$, where the S_i are simple submodules of A . Then any simple A -module is isomorphic with some S_i .*

Proof.

□

Proposition 13.5. *Suppose that A is a semisimple algebra, and let $S_1, \dots, S_r$ be a collection of simple A -modules such that every simple A -module is isomorphic with exactly one S_i . Let M be an A -module, and write $M \cong n_1 S_1 \oplus \cdots \oplus n_r S_r$ for some non-negative integers n_i . Then the n_i are uniquely determined.*

Proof.

□

Matrix Algebras over Division Algebras

- For a finite-dimensional F -algebra D , the matrix algebra $\mathcal{M}_n(D)$ has dimension $n^2 \dim_F D$.
- $E_{ij}(\alpha)$ denotes the matrix with sole non-zero entry α in position (i, j) .
- D^n is the natural left $\mathcal{M}_n(D)$ -module of column vectors.
- A *division algebra* is an algebra whose non-zero elements form a group under multiplication; field extensions of F are division algebras, but non-commutative examples also exist.

Theorem 13.6. *Let D be a division algebra, and let $n \in \mathbb{N}$. Then any simple $\mathcal{M}_n(D)$ -module is isomorphic with D^n , and $\mathcal{M}_n(D)$ is isomorphic as $\mathcal{M}_n(D)$ -modules with the direct sum of n copies of D^n . In particular, $\mathcal{M}_n(D)$ is a semisimple algebra.*

Proof. □

Simple Algebras

- An algebra is *simple* if its only two-sided ideals are itself and the zero ideal.
- Simple algebras are semisimple, and matrix algebras over division algebras are simple.

Lemma 13.7. *Simple algebras are semisimple.*

Proof. □

Theorem 13.8. *Let D be a division algebra, and let $n \in \mathbb{N}$. Then $\mathcal{M}_n(D)$ is a simple algebra.*

Proof. □

Direct Sums of Algebras

- The *external direct sum* $B = B_1 \oplus \cdots \oplus B_r$ has the Cartesian product of underlying sets with componentwise operations; each B_i embeds as an ideal.
- A B -module M becomes a B_i -module via $b_i m = (0, \dots, b_i, \dots, 0)m$, and is simple/semisimple as a B_i -module iff it is simple/semisimple as a B -module.
- An *internal direct sum* of ideals $B_1, \dots, B_r$ (vector-space direct sum) is isomorphic to the external direct sum.
- Two-sided ideals of a finite direct sum decompose componentwise.

Lemma 13.9. *Let $B = B_1 \oplus \cdots \oplus B_n$ be a direct sum of algebras. Then the two-sided ideals of B are exactly the sets of the form $J_1 \oplus \cdots \oplus J_n$, where J_i is a two-sided ideal of B_i for each i .*

Proof. □

Theorem 13.10. *Let $r \in \mathbb{N}$. For each $1 \leq i \leq r$, let D_i be a division algebra over F , let $n_i \in \mathbb{N}$, and let $B_i = \mathcal{M}_{n_i}(D_i)$. Let B be the external direct sum of the B_i . Then B is a semisimple algebra having exactly r isomorphism classes of simple modules and exactly 2^r two-sided ideals, namely every sum of the form $\bigoplus_{j \in J} B_j$, where J is a subset of $\{1, \dots, r\}$.*

Proof. □

Endomorphism and Opposite Algebras

- For an A -module M , $\text{End}_A(M)$ is an F -algebra under composition; we call it the *endomorphism algebra* of M .
- The *opposite algebra* B^{op} has the same underlying additive structure but reversed multiplication: $a \cdot b := ba$. Note $(B^{\text{op}})^{\text{op}} = B$, and the opposite of a division algebra is again a division algebra.
- These two notions are linked by $B^{\text{op}} \cong \text{End}_B(B)$: every B -endomorphism of B is right multiplication by some element.
- Endomorphism algebras decompose along A -module direct sums into block form, and over a single simple module they yield a matrix algebra.

Lemma 13.11. *Let B be an algebra. Then $B^{\text{op}} \cong \text{End}_B(B)$.*

Proof.

□

Lemma 13.12. *Let $S_1, \dots, S_r$ be the distinct simple A -modules; for each i , let U_i be a direct sum of copies of S_i , and let $U = U_1 \oplus \dots \oplus U_r$. Then we have $\text{End}_A(U) \cong \text{End}_A(U_1) \oplus \dots \oplus \text{End}_A(U_r)$.*

Proof.

□

Lemma 13.13. *If S is a simple A -module, then for any $n \in \mathbb{N}$ we have $\text{End}_A(nS) \cong \mathcal{M}_n(\text{End}_A(S))$.*

Proof.

□

Lemma 13.14. *Suppose that F is algebraically closed, and let S be a simple A -module. Then $\text{End}_A(S) \cong F$.*

Proof.

□

Lemma 13.15. *Let B be an algebra. Then $\mathcal{M}_n(B)^{\text{op}} \cong \mathcal{M}_n(B^{\text{op}})$ for any $n \in \mathbb{N}$.*

Proof.

□

Wedderburn's Structure Theorems

- Wedderburn's main structure theorem (Theorem 13.16): a finite-dimensional algebra is semisimple iff it is isomorphic to a direct sum of matrix algebras over division algebras.
- Equivalently, semisimple iff a direct sum of simple algebras, confirming the consistency of terminology.
- An algebra is simple iff it is isomorphic to a single matrix algebra over a division algebra (Theorem 13.17).
- Over an algebraically closed field, every semisimple algebra is a direct sum of full matrix algebras over F itself (Theorem 13.18).

Theorem 13.16. *The algebra A is semisimple iff it is isomorphic with a direct sum of matrix algebras over division algebras.*

Proof.

□

Theorem 13.17. *The algebra A is simple iff it is isomorphic with a matrix algebra over a division algebra.*

Proof.

□

Theorem 13.18. *Suppose that the field F is algebraically closed. Then any semisimple algebra is isomorphic with a direct sum of matrix algebras over F .*

Proof.

□

Nilpotent Ideals and the Radical

- In this part, A is an algebra possibly without unit. An element $x \in A$ is *nilpotent* if $x^n = 0$ for some $n \in \mathbb{N}$.
- An ideal I is *nilpotent* if $I^n = 0$ for some n , where I^n is the ideal spanned by all products of n elements; an algebra is nilpotent if some such n exists for $I = A$.
- A nilpotent algebra cannot have a unit.
- The sum of two nilpotent ideals is nilpotent, and (using finite-dimensionality) a unique largest nilpotent ideal exists.
- Submodules with semisimple quotient are closed under finite intersection, so a smallest such submodule of A exists.
- For an A -module M , $\text{Ann}(M) = \{a \in A \mid aM = 0\}$ is the *annihilator* of M , an ideal of A ; the *common annihilator* of a family $\{M_t\}_{t \in \mathcal{T}}$ is $\bigcap_{t \in \mathcal{T}} \text{Ann}(M_t)$.
- Wedderburn (Theorem 13.23): the largest nilpotent ideal, the common annihilator of all simple A -modules, and the smallest submodule of A with semisimple quotient all coincide. This common object is the *radical* $\text{rad}(A)$, also called the Jacobson radical $J(A)$.

Lemma 13.19. *Let I and J be nilpotent ideals of A . Then $I + J$ is also a nilpotent ideal of A .*

Proof. □

Corollary 13.20. *A has a largest nilpotent ideal.*

Proof. □

Lemma 13.21. *Let M and N be submodules of A such that A/M and A/N are semisimple A -modules. Then $A/(M \cap N)$ is a semisimple A -module.*

Proof. □

Corollary 13.22. *A has a smallest submodule having semisimple quotient.*

Proof. □

Theorem 13.23. *Let A be an algebra with unit. Then the following are equal:*

- *The largest nilpotent ideal of A .*
- *The common annihilator of all simple A -modules.*
- *The smallest submodule of A having semisimple quotient.*

Proof. □

Corollary 13.24. *Let A be an algebra with unit. Then A is semisimple iff A has no non-zero nilpotent ideals.*

Proof. □

Algebras Without Unit; Nilpotence

- Any algebra (possibly without unit) embeds as an ideal in an algebra with unit, with quotient F .
- Consequence: an algebra without unit having radical zero cannot exist—no such algebras occur.
- Wedderburn: an algebra generated as an F -vector space by nilpotent elements is itself nilpotent. This finishes Kolchin's Theorem.

Lemma 13.25. *Let A be an algebra, possibly without unit. Then there exists an algebra B with unit and an ideal I of B such that, as algebras, we have $I \cong A$ and $B/I \cong F$.*

Proof. □

Theorem 13.26. *Let A be a non-zero algebra, possibly without unit. If A has no non-zero nilpotent ideals, then A is an algebra with unit.*

Proof. □

Theorem 13.27. *Suppose that the algebra A can be generated as an F -vector space by a set consisting of nilpotent elements. Then A is nilpotent.*

Proof. □

Proposition 13.28. *Let $V_n(F)$ be the space of column vectors of length $n \in \mathbb{N}$ with entries from a field F , and let H be a unipotent subgroup of $\mathrm{GL}(n, F)$. Then there exists some $0 \neq \mathbf{v} \in V_n(F)$ such that $x\mathbf{v} = \mathbf{v}$ for all $x \in H$.*

Proof. □

Exercises

Throughout these exercises, A denotes a finite-dimensional algebra with unit over a field F , and all A -modules are finitely generated.

Exercise 13.1. Let $n \in \mathbb{N}$. Let V be the n -dimensional subspace of A^n spanned by the identity elements of the summands. Show that if M is an A -module and $T: V \rightarrow M$ is a linear transformation, then there is a unique A -module homomorphism from A^n to M which extends T .

Solution. □

Exercise 13.2. (cont.) Suppose that B is an A -module having an n -dimensional subspace U such that whenever M is an A -module and $T: U \rightarrow M$ is a linear transformation, there is a unique extension of T to an A -module homomorphism from B to M . Show that B and A^n are isomorphic A -modules. (Modules of the form A^n for some $n \in \mathbb{N}$ are called *free* A -modules; hence this exercise provides an alternate characterization of freeness, which we shall generalize for group algebras in the exercises to Section 16.)

Solution. □

Exercise 13.3. Let $n \in \mathbb{N}$, and let $T_n(F)$ be the algebra of upper triangular $n \times n$ matrices. Show that the set $V_n(F)$ of column vectors over F of length n is a $T_n(F)$ -module that has a unique composition series in which every simple $T_n(F)$ -module appears exactly once as a composition factor.

Solution. □

Exercise 13.4. (cont.) Show that the $T_n(F)$ -module $T_n(F)$ is isomorphic with the direct sum of all non-zero submodules of $V_n(F)$.

Solution. □

Exercise 13.5. (cont.) Find the largest nilpotent ideal of $T_n(F)$, and show without reference to Theorem 13.23 that it is also the common annihilator of all simple $T_n(F)$ -modules and the smallest submodule of $T_n(F)$ having semisimple quotient.

Solution. □

Exercise 13.6. Let U be an A -module, let $n \in \mathbb{N}$, and let U^n be the set of column vectors of length n with entries from U , considered in the obvious way as an $\mathcal{M}_n(A)$ -module. Show that U is a simple A -module iff U^n is a simple $\mathcal{M}_n(A)$ -module.

Solution. □

Exercise 13.7. (cont.) Show that $\mathrm{Hom}_A(U, V) \cong \mathrm{Hom}_{\mathcal{M}_n(A)}(U^n, V^n)$ for any A -modules U and V .

Solution. □

Exercise 13.8. (cont.) Show that every $\mathcal{M}_n(A)$ -module is isomorphic with a module of the form U^n for some A -module U . (What Exercises 13.6–13.8 show is that the module theory of the algebras A and $\mathcal{M}_n(A)$ is, in a sense that can be made precise, the same. In the language of category theory, we say that the categories of A -modules and $\mathcal{M}_n(A)$ -modules are *Morita equivalent*.)

Solution. □

Exercise 13.9. Show that A is a simple A -module iff A is a division algebra.

Solution. □

Exercise 13.10. We can clearly regard an $A/\text{rad } A$ -module as being an A -module; on the other hand, a simple A -module is annihilated by $\text{rad}(A)$ and hence can be considered as an $A/\text{rad}(A)$ -module. Show that an A -module is simple iff it is a simple $A/\text{rad}(A)$ -module. (Since $A/\text{rad}(A)$ is a semisimple algebra, this implies that the determination of simple modules over arbitrary algebras reduces to the case of semisimple algebras.)

Solution. □

Exercise 13.11. Let B be a finite-dimensional F -algebra, possibly without unit. Show that if B is a simple algebra whose multiplication is not identically zero, then B is isomorphic with a matrix algebra over a division algebra.

Solution. □

Further Exercises

Let A be as above, and let M be a finitely generated A -module.

Exercise 13.12. Show that the following objects (exist and) are equal: the intersection of all maximal submodules of M ; the smallest submodule of M having semisimple quotient; and $\text{rad}(A)M$. This submodule of M is called the *radical* of M and is denoted $\text{rad}(M)$. (Observe that our two definitions of the symbol $\text{rad}(A)$ agree.)

We define a descending series $\text{rad}^n(M)$ of submodules of M by setting $\text{rad}^0(M) = M$ and $\text{rad}^n(M) = \text{rad}(\text{rad}^{n-1}(M))$ for $n \in \mathbb{N}$. It follows from Exercise 13.12 that $\text{rad}^n(M) = (\text{rad}(A))^n M$ for any $n \in \mathbb{N}$. This series is called the *radical series* of M . Observe that $\text{rad}^n(M) = 0$ for some n , since $\text{rad}(A)$ is a nilpotent ideal, and that each successive quotient of the radical series is a semisimple module, with M being semisimple iff $\text{rad}(M) = 0$.

Solution. □

Exercise 13.13. (cont.) Show that M has a unique largest semisimple submodule and that this submodule is equal to $\{m \in M \mid \text{rad}(A)m = 0\}$. We call this submodule the *socle* of M , and we denote it by $\text{soc}(M)$.

Solution. □

Exercise 13.14. (cont.) We define an ascending series $\text{soc}^n(M)$ of submodules of M as follows: We let $\text{soc}^0(M) = 0$ and $\text{soc}^1(M) = \text{soc}(M)$, and for $n > 1$ we let $\text{soc}^n(M)$ be the submodule of M such that $\text{soc}^n(M)/\text{soc}^{n-1}(M) = \text{soc}(M/\text{soc}^{n-1}(M))$. Show that we have $\text{soc}^n(M) = \{m \in M \mid \text{rad}(A)^n m = 0\}$ for any $n \in \mathbb{N}$. (This series is called the *socle series* of M . As with the radical series, we observe that $\text{soc}^n(M) = M$ for some n , since $\text{rad}(A)$ is nilpotent, and that each successive quotient of the socle series is semisimple, with M being semisimple iff $\text{soc}(M) = M$.)

Solution. □

Exercise 13.15. (cont.) Show that $\text{rad}^n(M) = 0$ iff $\text{soc}^n(M) = M$; conclude that the radical and socle series of M have the same finite length. If this common length is r , then show that $\text{rad}^n(M) \subseteq \text{soc}^{r-n}(M)$ for every $0 \leq n \leq r$.

Solution. □

Chapter 6

Group Representations

14 Characters

Standing assumptions

- In this chapter G denotes a finite group.
- All $\mathbb{C}G$ -modules are finitely generated, equivalently, finite-dimensional as $\mathbb{C}$ -vector spaces.
- $\mathbb{C}$ is the field of complex numbers.

Wedderburn structure of $\mathbb{C}G$

- Since $\mathbb{C}$ is algebraically closed of characteristic zero, Wedderburn theory pins down the structure of $\mathbb{C}G$ precisely.
- The simple summands $\mathcal{M}_{f_i}(\mathbb{C})$ correspond bijectively to isomorphism classes of simple $\mathbb{C}G$ -modules S_i , with $\dim_{\mathbb{C}} S_i = f_i$.

Theorem 14.1. *There is some $r \in \mathbb{N}$ and some $f_1, \dots, f_r \in \mathbb{N}$ such that $\mathbb{C}G \cong \mathcal{M}_{f_1}(\mathbb{C}) \oplus \dots \oplus \mathcal{M}_{f_r}(\mathbb{C})$ as $\mathbb{C}$ -algebras. Furthermore, there are exactly r isomorphism classes of simple $\mathbb{C}G$ -modules, and if we let $S_1, \dots, S_r$ be representatives of these r classes, then we can order the S_i so that $\mathbb{C}G \cong f_1 S_1 \oplus \dots \oplus f_r S_r$ as $\mathbb{C}G$ -modules, where $\dim_{\mathbb{C}} S_i = f_i$ for each i . Any $\mathbb{C}G$ -module can be written uniquely in the form $a_1 S_1 \oplus \dots \oplus a_r S_r$, where the a_i are non-negative integers.*

Proof.

□

Degrees of G

- The $\mathbb{C}$ -dimensions $f_1, \dots, f_r$ of the r simple $\mathbb{C}G$ -modules are called the *degrees* of G .
- The trivial $\mathbb{C}G$ -module $\mathbb{C}$ is one-dimensional and hence simple, so G always has at least one degree equal to 1; by convention we set $f_1 = 1$.

Corollary 14.2. *We have $\sum_{i=1}^r f_i^2 = |G|$.*

Proof.

□

Degrees and conjugacy classes

- In fact the degrees of G must divide $|G|$; we shall not use this fact, but we provide a proof of it in the Appendix.
- The next theorem links the number of simple $\mathbb{C}G$ -modules to the conjugacy structure of G via the center of $\mathbb{C}G$.

Theorem 14.3. *The number r of simple $\mathbb{C}G$ -modules is equal to the number of conjugacy classes of G .*

Proof.

□

Defining characters

- In general there is no natural bijective correspondence between the conjugacy classes of G and the simple $\mathbb{C}G$ -modules. However, if G is a symmetric group, then there is such a correspondence, although we shall not develop it here; for an overview, see [13, Section 4.1].
- If U is a $\mathbb{C}G$ -module, then each $g \in G$ defines an invertible linear transformation of U that sends $u \in U$ to gu . The *character* of U is the function $\chi_U: G \rightarrow \mathbb{C}$, where $\chi_U(g)$ is the trace of this linear transformation of U defined by g .
- For example, $\chi_U(1) = \dim_{\mathbb{C}} U$, since the identity element of G induces the identity transformation on U .
- If $\rho: G \rightarrow \text{GL}(U)$ is the representation corresponding to U , then $\chi_U(g)$ is just the trace of the map $\rho(g)$.
- Isomorphic $\mathbb{C}G$ -modules have equal characters.
- For any $g, h \in G$ the linear transformations of U defined by g and hgh^{-1} are similar and hence have the same trace; so any character of G is constant on each conjugacy class of G .
- For example, let $U = \mathbb{C}G$ and let $g \in G$. With respect to the basis G of $\mathbb{C}G$, $\chi_U(g)$ equals the number of elements $x \in G$ for which $gx = x$. Therefore $\chi_U(1) = |G|$ and $\chi_U(g) = 0$ for every $1 \neq g \in G$. This character is called the *regular character* of G .
- The theory of characters was developed by Frobenius and others, starting in 1896. At first nothing was known about linear representations, let alone modules over the group algebra; Frobenius defined characters as being functions from G to $\mathbb{C}$ satisfying certain properties, but it turned out that his characters were exactly the trace functions of finitely generated $\mathbb{C}G$ -modules.
- We will denote the characters of the r simple $\mathbb{C}G$ -modules by $\chi_1, \dots, \chi_r$; these characters will be referred to as the *irreducible characters* of G . Whenever we say that $S_1, \dots, S_r$ are the distinct simple $\mathbb{C}G$ -modules, we implicitly order them so that $\chi_{S_i} = \chi_i$ for each i .
- In keeping with the convention that f_1 denotes the degree of the trivial representation, we let χ_1 be the character of the trivial representation; we call χ_1 the *principal character* of G , and we have $\chi_1(g) = 1$ for all $g \in G$.
- A character of a one-dimensional $\mathbb{C}G$ -module is called a *linear character*. Since one-dimensional modules are simple, all linear characters are irreducible. If χ arises from a one-dimensional $\mathbb{C}G$ -module U , then for $g, h \in G$ and $u \in U$ we have $gu = \chi(g)u$ and $hu = \chi(h)u$, so $\chi(gh)u = (gh)u = \chi(g)\chi(h)u$; hence χ is a homomorphism from G to $\mathbb{C}^\times$. Conversely, given a homomorphism $\varphi: G \rightarrow \mathbb{C}^\times$, we can define a one-dimensional $\mathbb{C}G$ -module U by $gu = \varphi(g)u$ for $g \in G$ and $u \in U$, and we then have $\chi_U = \varphi$. Therefore, linear characters of G are exactly the same as group homomorphisms from G to $\mathbb{C}^\times$.

Proposition 14.4. *Let U be a $\mathbb{C}G$ -module, let $\rho: G \rightarrow \text{GL}(U)$ be the representation corresponding to U , and let $g \in G$ be of order n . Then:*

- $\rho(g)$ is diagonalizable.
- $\chi_U(g)$ equals the sum (with multiplicities) of the eigenvalues of $\rho(g)$.
- $\chi_U(g)$ is a sum of $\chi_U(1)$ n th roots of unity.
- $\chi_U(g^{-1}) = \overline{\chi_U(g)}$. (Here $\bar{z}$ denotes the complex conjugate of $z \in \mathbb{C}$.)
- $|\chi_U(g)| \leq \chi_U(1)$.
- $\{x \in G \mid \chi_U(x) = \chi_U(1)\}$ is a normal subgroup of G .

Proof.

□

Algebra of characters

- If χ and ψ are characters of G , we can define new functions $\chi + \psi$ and $\chi\psi$ from G to $\mathbb{C}$ by $(\chi + \psi)(g) = \chi(g) + \psi(g)$ and $(\chi\psi)(g) = \chi(g)\psi(g)$ for $g \in G$.
- These new functions obtained from characters are not, a priori, characters themselves.
- Given a scalar $\lambda \in \mathbb{C}$, we can also define a new function $\lambda\chi: G \rightarrow \mathbb{C}$ by $(\lambda\chi)(g) = \lambda\chi(g)$, so we view characters of G as elements of a $\mathbb{C}$ -vector space of functions from G to $\mathbb{C}$.

Proposition 14.5. *The irreducible characters of G are, as functions from G to $\mathbb{C}$, linearly independent over $\mathbb{C}$.*

Proof. □

Lemma 14.6. $\chi_{U \oplus V} = \chi_U + \chi_V$ for any $\mathbb{C}G$ -modules U and V .

Proof. □

Characters distinguish modules

- Characters of $\mathbb{C}G$ -modules suffice to distinguish $\mathbb{C}G$ -modules up to isomorphism.

Theorem 14.7. *If $S_1, \dots, S_r$ are the distinct simple $\mathbb{C}G$ -modules, then the character of the $\mathbb{C}G$ -module $a_1 S_1 \oplus \dots \oplus a_r S_r$ (where the a_i are non-negative integers) is $a_1 \chi_1 + \dots + a_r \chi_r$. Consequently, two $\mathbb{C}G$ -modules are isomorphic iff their characters are equal.*

Proof. □

Tensor products, duals, and Hom

- Although each character determines a $\mathbb{C}G$ -module up to isomorphism by Theorem 14.7, there is no generic way to construct a module from its corresponding character, and so in some sense information is lost by studying characters in lieu of modules.
- Characters turn out to be an efficient means of translating information about the ordinary representation theory of G to information about G itself; from now on we focus on characters rather than modules.
- Recall from Section 12 that if U and V are $\mathbb{C}G$ -modules, then $U \otimes_{\mathbb{C}} V$ and $\text{Hom}_{\mathbb{C}}(U, V)$, written here as $U \otimes V$ and $\text{Hom}(U, V)$, admit natural $\mathbb{C}G$ -module structures.

Proposition 14.8. *Let U and V be $\mathbb{C}G$ -modules. Then:*

- (i) $\chi_{U \otimes V} = \chi_U \chi_V$.
- (ii) $\chi_{U^*} = \overline{\chi_U}$. (Recall that $U^* = \text{Hom}(U, F)$.)
- (iii) $\chi_{\text{Hom}(U, V)} = \overline{\chi_U} \chi_V$.

Proof. □

Virtual characters and class functions

- A *virtual character* of G is a $\mathbb{Z}$ -linear combination of the irreducible characters of G . (Some authors prefer the term “generalized character.”) Characters are virtual characters by Theorem 14.7.

Corollary 14.9. *The virtual characters of G form a ring.*

Proof. □

Class functions

- A *class function* on G is a function from G to $\mathbb{C}$ whose value within any conjugacy class is constant.
- Characters of $\mathbb{C}G$ -modules are class functions.
- The set of class functions on G forms a $\mathbb{C}$ -vector space of dimension r , where r is the number of conjugacy classes of G ; an obvious basis is the set of functions taking the value 1 on a single conjugacy class and 0 on all other classes.

Proposition 14.10. *The irreducible characters of G form a basis for the space of class functions on G .*

Proof. □

Inner product on class functions

- If α and β are class functions on G , then their *inner product* is the complex number

$$(\alpha, \beta) = \frac{1}{|G|} \sum_{g \in G} \alpha(g) \overline{\beta(g)}.$$

- This function $(\cdot, \cdot)$ is, as the name suggests, an inner product on the space of class functions:
 - $(\alpha, \alpha) \geq 0$ for all α , and $(\alpha, \alpha) = 0$ iff $\alpha = 0$.
 - $(\alpha, \beta) = \overline{(\beta, \alpha)}$ for all α, β .
 - $(\lambda\alpha, \beta) = \lambda(\alpha, \beta)$ for all α, β and all $\lambda \in \mathbb{C}$.
 - $(\alpha_1 + \alpha_2, \beta) = (\alpha_1, \beta) + (\alpha_2, \beta)$ for all $\alpha_1, \alpha_2, \beta$.
- The following are easy consequences of the above properties:
 - $(\alpha, \lambda\beta) = \overline{\lambda}(\alpha, \beta)$ for all α, β and all $\lambda \in \mathbb{C}$.
 - $(\alpha, \beta_1 + \beta_2) = (\alpha, \beta_1) + (\alpha, \beta_2)$ for all α, β_1, β_2 .
- We need a small piece of notation: if U is a $\mathbb{C}G$ -module, the set of elements of U on which G acts trivially is a $\mathbb{C}G$ -submodule, denoted $U^G = \{u \in U \mid gu = u \text{ for all } g \in G\}$.

Lemma 14.11. *If U is a $\mathbb{C}G$ -module, then $\dim_{\mathbb{C}} U^G = \frac{1}{|G|} \sum_{g \in G} \chi_U(g)$.*

Proof. □

Theorem 14.12. *We have $(\chi_U, \chi_V) = \dim_{\mathbb{C}} \text{Hom}_{\mathbb{C}G}(U, V)$ for any $\mathbb{C}G$ -modules U and V .*

Proof. □

Exercises

Throughout these exercises, G denotes a finite group, and all modules are finitely generated.

Exercise 14.1. Let A be a semisimple $\mathbb{C}$ -algebra. Let $[A, A]$ be the subspace of A spanned by the set $\{ab - ba \mid a, b \in A\}$. Show that the codimension of $[A, A]$ in A (which is just the dimension of $A/[A, A]$) is equal to the number of isomorphism classes of simple A -modules.

Solution. □

Exercise 14.2. (cont.) Let $A = \mathbb{C}G$, and let $g_1, \dots, g_r$ be representatives of the r conjugacy classes of G . Show that $\{g_i + [A, A] \mid 1 \leq i \leq r\}$ is a basis of $A/[A, A]$.

Solution. □

Exercise 14.3. Show that if S is a simple $\mathbb{C}G$ -module and U is a one-dimensional $\mathbb{C}G$ -module, then $S \otimes U$ is simple.

Solution. □

Exercise 14.4. Show that the set of isomorphism classes of the one-dimensional $\mathbb{C}G$ -modules forms a group under the taking of tensor products.

Solution. □

Exercise 14.5. Let χ be an irreducible character of G . Show that if λ is any $|G|$ th root of unity, then $\{x \in G \mid \chi(x) = \lambda\chi(1)\} \trianglelefteq G$.

Solution. □

Exercise 14.6. Show that a $\mathbb{C}G$ -module U is simple iff its dual U^* is simple. Conclude that the complex conjugate of an irreducible character is again an irreducible character.

Solution. □

If F is any field and U is any FG -module, then we define the character of U to be an F -valued function on U whose value at $g \in G$ is the trace of the F -endomorphism of U induced by g , exactly as was done in this section in the case $F = \mathbb{C}$.

Exercise 14.7. Suppose that F has characteristic $p > 0$. Give an example to demonstrate that two FG -modules that have the same character need not be isomorphic.

Solution. □

Exercise 14.8. Suppose that F has characteristic $p > 0$. Show that if U is an FG -module then we have $\chi_U(g) = \chi_U(g_{p'})$ for any $g \in G$, where $g_{p'}$ is the p' -part of g as defined in Exercise 1.4.

Solution. □

15 The Character Table

Definition of the Character Table

- Any character of G is a $\mathbb{Z}$ -linear combination of the r irreducible characters $\chi_1, \dots, \chi_r$, where r is the number of conjugacy classes of G (Theorem 14.3).
- Each irreducible character is determined by its values on the r conjugacy classes, so the irreducible characters are completely determined by an $r \times r$ array.
- The *character table* of G is this $r \times r$ array $\mathcal{X} = (\chi_i(g_j))_{1 \leq i, j \leq r}$, where $g_1, \dots, g_r$ are conjugacy class representatives.
- The character table is well-defined only up to reordering of rows and columns.
- By convention, $g_1 = 1$, so the first column consists of the degrees of G .
- We write $\mathcal{X}$ in the form

$$\begin{array}{c|cccc}
 & 1 & k_2 & \cdots & k_r \\
 & 1 & g_2 & \cdots & g_r \\
 \hline
 \chi_1 & 1 & 1 & \cdots & 1 \\
 \chi_2 & f_2 & \chi_2(g_2) & \cdots & \chi_2(g_r) \\
 \vdots & \vdots & \vdots & \ddots & \vdots \\
 \chi_r & f_r & \chi_r(g_2) & \cdots & \chi_r(g_r)
 \end{array}$$

where the f_i are the degrees of G , and $k_i = |G : C_G(g_i)|$ is the order of the conjugacy class of g_i (Proposition 3.13).

- By convention χ_1 is always the principal character.

Orthogonality of Rows

Theorem (Row Orthogonality Theorem). $(\chi_i, \chi_j) = \delta_{ij}$ for any i and j .

Proof.

□

In other words, this theorem asserts that

$$\delta_{ij} = \frac{1}{|G|} \sum_{g \in G} \chi_i(g) \overline{\chi_j(g)} = \frac{1}{|G|} \sum_{t=1}^r k_t \chi_i(g_t) \overline{\chi_j(g_t)}$$

for any i and j , where the g_t are conjugacy class representatives and the k_t are the orders of the conjugacy classes. The rows of the character table, considered as vectors in $\mathbb{C}^r$, are orthogonal with respect to an inner product that differs slightly from the standard one.

Corollary 15.1. *The irreducible characters of G form an orthonormal basis for the vector space of class functions on G .*

Proof.

□

Corollary 15.2. *If $\alpha = \sum_i a_i \chi_i$ and $\beta = \sum_j b_j \chi_j$ are virtual characters of G , then $(\alpha, \beta) = \sum_i a_i b_i$.*

Proof.

□

Corollary 15.3. *If α is a character of G and $n \in \{1, 2, 3\}$, then $(\alpha, \alpha) = n$ iff α is a sum of n irreducible characters.*

Proof.

□

Corollary 15.4. *If α is a virtual character of G , then each χ_j appears with coefficient (α, χ_j) in the unique expression of α as a $\mathbb{Z}$ -linear combination of the irreducible characters of G .*

Proof.

□

Proposition 15.5. *If α is a linear character of G and χ is an irreducible character of G , then $\alpha\chi$ is an irreducible character of G .*

Proof.

□

Orthogonality of Columns

Theorem (Column Orthogonality Theorem). *If $g_1, \dots, g_r$ are a set of conjugacy class representatives of G , and $k_1, \dots, k_r$ are the orders of the conjugacy classes, then for any $1 \leq i, j \leq r$ we have*

$$\sum_{t=1}^r \chi_t(g_i) \overline{\chi_t(g_j)} = \frac{|G|}{k_i} \delta_{ij} = |C_G(g_i)| \delta_{ij}.$$

Proof. □

Characters and Quotient Groups

Lemma 15.6. *Let $N \trianglelefteq G$, and let U be a $\mathbb{C}(G/N)$ -module. Then U admits a canonical $\mathbb{C}G$ -module structure, with a subspace of U being a $\mathbb{C}G$ -submodule iff it is a $\mathbb{C}(G/N)$ -submodule. If ψ is the character of the $\mathbb{C}(G/N)$ -module U , then the character of the $\mathbb{C}G$ -module U is $\psi \circ \eta$, where $\eta: G \rightarrow G/N$ is the natural map.*

Proof. □

We define $K_\chi = \{x \in G \mid \chi(x) = \chi(1)\}$ for any character χ of G ; we have $K_\chi \trianglelefteq G$ by part (vi) of Proposition 14.4. We call K_χ the *kernel* of χ , as it is the kernel of the corresponding representation. We write K_i instead of K_{χ_i} .

Proposition 15.7. *The normal subgroups of G are exactly the sets of the form $\bigcap_{i \in I} K_i$ for some $I \subseteq \{1, \dots, r\}$.*

Proof. □

Corollary 15.8. *G is simple iff the only irreducible character χ_i for which $\chi_i(g) = \chi_i(1)$ for some $1 \neq g \in G$ is the principal character χ_1 .*

Proof. □

Corollary 15.9. *The character table of G can be used to determine whether or not G is solvable.*

Proof. □

We define $Z_\chi = \{x \in G \mid |\chi(x)| = \chi(1)\}$ for any character χ of G . Again we write Z_i instead of Z_{χ_i} .

Lemma 15.10. *$Z_\chi \leq G$ for any character χ of G , and if in addition χ is irreducible, then $Z_\chi/K_\chi = Z(G/K_\chi)$.*

Proof. □

Corollary 15.11. *If G is non-abelian and simple, then $Z_i = 1$ for all $i > 1$.*

Proof. □

Proposition 15.12. $Z(G) = \bigcap_{i=1}^r Z_i$.

Proof. □

Lemma 15.13. *If $N \trianglelefteq G$, then the irreducible characters of G/N can be determined from those of G .*

Proof. □

Corollary 15.14. *The character table of G can be used to determine whether or not G is nilpotent.*

Proof. □

Observe that in Corollary 15.9 we assumed that we knew both the irreducible characters of G and the orders of the conjugacy classes, whereas in Corollary 15.14 we did not need to know the orders of the classes. As evidenced by Lemma 15.13, there may be circumstances in which we can construct a character table without knowing the orders of the conjugacy classes. There is a theorem of G. Higman (see [15, p. 136]) which asserts that if we know the irreducible characters of G , but not the orders of the conjugacy classes, then we can at least determine the sets of prime divisors of the orders of the conjugacy classes.

Examples: Constructing Character Tables

The remainder of this section consists of a series of examples in which we develop methods of finding characters and use these methods to construct the character tables of various groups.

Example 15.1. Let $G = \langle g \rangle \cong \mathbb{Z}_n$ for some $n \in \mathbb{N}$, and let λ be a primitive n th root of unity. For each $1 \leq i \leq n$, let V_i be a one-dimensional $\mathbb{C}$ -vector space, and let g act on V_i by multiplication by λ^{i-1} ; since G is cyclic, this definition completely determines a $\mathbb{C}G$ -module structure on V_i . Each V_i , being one-dimensional, is a simple $\mathbb{C}G$ -module. If χ_i is the character of V_i , then $\chi_i(g) = \lambda^{i-1}$, and hence $\chi_i(g^a) = \lambda^{a(i-1)}$ for any a . The characters $\chi_1, \dots, \chi_n$ are n distinct linear characters of G , and since G can have at most $|G| = n$ irreducible characters, we see that the χ_i are precisely the irreducible characters of G . Thus, the character table of G is

	1	1	1	$\dots$	1
	1	g	g^2	$\dots$	g^{n-1}
χ_1	1	1	1	$\dots$	1
χ_2	1	λ	λ^2	$\dots$	λ^{n-1}
χ_3	1	λ^2	λ^4	$\dots$	λ^{n-2}
$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\ddots$	$\vdots$
χ_n	1	λ^{n-1}	λ^{n-2}	$\dots$	λ

Observe that the set $\{\chi_1, \dots, \chi_n\}$ is a cyclic group under multiplication of characters, with generator χ_2 ; we have $\chi_i = \chi_2^{i-1}$ for any i .

Example 15.2. Let G and H be groups with respective irreducible characters $\chi_1, \dots, \chi_r$ and $\psi_1, \dots, \psi_s$. We wish to determine the irreducible characters of $G \times H$. We see that two elements (x, y) and (x', y') of $G \times H$ are conjugate iff x and x' are conjugate in G and y and y' are conjugate in H . Since by Theorem 14.3, G and H have r and s conjugacy classes, respectively, we conclude that $G \times H$ has rs conjugacy classes, with each class of $G \times H$ being the product of a class of G and a class of H ; hence by Theorem 14.3, $G \times H$ has rs irreducible characters.

Let $S_1, \dots, S_r$ and $T_1, \dots, T_s$ be the distinct simple $\mathbb{C}G$ - and $\mathbb{C}H$ -modules, respectively. For each i and j , we give $S_i \otimes T_j$ the structure of a $\mathbb{C}(G \times H)$ -module by $(g, h)(s \otimes t) = gs \otimes ht$ and linear extension. Let τ_{ij} be the character of $S_i \otimes T_j$ for each i and j . By imitating the proof of part (i) of Proposition 14.8, we find that $\tau_{ij}(g, h) = \chi_i(g)\psi_j(h)$; we adopt the notation $\tau_{ij} = \chi_i \times \psi_j$. We note that the τ_{ij} are distinct and that we can recover the χ_i and ψ_j from the τ_{ij} by appropriate restriction. Now for any i, i', j, j' we have

$$\begin{aligned}
 (\tau_{ij}, \tau_{i'j'}) &= \frac{1}{|G \times H|} \sum_{(g,h) \in G \times H} \tau_{ij}(g, h) \overline{\tau_{i'j'}(g, h)} \\
 &= \frac{1}{|G||H|} \sum_{g \in G} \sum_{h \in H} \chi_i(g) \psi_j(h) \overline{\chi_{i'}(g) \psi_{j'}(h)} \\
 &= \left(\frac{1}{|G|} \sum_{g \in G} \chi_i(g) \overline{\chi_{i'}(g)} \right) \left(\frac{1}{|H|} \sum_{h \in H} \psi_j(h) \overline{\psi_{j'}(h)} \right) \\
 &= (\chi_i, \chi_{i'}) (\psi_j, \psi_{j'}) = \delta_{ii'} \delta_{jj'}
 \end{aligned}$$

by Row Orthogonality Theorem. In particular, we have $(\tau_{ij}, \tau_{ij}) = 1$ for each i and j , which by Corollary 15.3 implies that τ_{ij} is an irreducible character. Therefore the τ_{ij} are the rs irreducible characters of $G \times H$; that is, the set of irreducible characters of $G \times H$ is $\{\chi_i \times \psi_j\}_{i,j}$.

Example 15.3. Suppose that G is abelian. Then every conjugacy class of G contains exactly one element, and consequently G has $|G|$ irreducible characters by Theorem 14.3. But $\sum_{i=1}^{|G|} f_i^2 = |G|$ by Corollary 14.2, so we must have $f_i = 1$ for each i . Therefore all irreducible characters of G are linear. By Corollary 8.8, G is a direct product of cyclic p -groups; let these cyclic p -groups be of orders $p_1^{a_1}, \dots, p_t^{a_t}$ with respective generators $g_1, \dots, g_t$. We could determine the character table of G from those of the cyclic p -groups using Examples 15.1 and 15.2, but there is an alternate method, which we now describe. A linear character χ of G is just a homomorphism from G to $\mathbb{C}^\times$, and hence to define such a χ it suffices to specify $\chi(g_i)$ for each i , with the only restriction on $\chi(g_i)$ being that it is a $p_i^{a_i}$ -th root of unity.

Therefore, there is a bijective correspondence between irreducible characters of G and ordered t -tuples $(\lambda_1, \dots, \lambda_t)$ in which each λ_i is a $p_i^{a_i}$ -th root of unity.

As a simple example, let $G = \langle a, b \rangle \cong \mathbb{Z}_2 \times \mathbb{Z}_2$. Both a and b have order 2, so by the above paragraph we see that the four irreducible characters of G correspond to the ordered pairs $(1, 1)$, $(1, -1)$, $(-1, 1)$, and $(-1, -1)$. Therefore, the character table for G is

	1	1	1	1
	1	a	b	ab
χ_1	1	1	1	1
χ_2	1	1	-1	-1
χ_3	1	-1	1	-1
χ_4	1	-1	-1	1

Here the second and third columns correspond to the ordered pairs given above, and the fourth column is wholly determined by the previous two columns.

Example 15.4. Let X be a finite G -set. We saw in Section 12 that the vector space $\mathbb{C}X$ having basis X is a $\mathbb{C}G$ -module. We wish to determine the character π of $\mathbb{C}X$. Consider the matrix, with respect to the basis X , of the linear transformation of $\mathbb{C}X$ defined by $g \in G$. Since $gx \in X$ for every $x \in X$, this matrix is a permutation matrix; but $\pi(g)$ is the sum of the diagonal entries of this matrix, so we conclude that $\pi(g)$ equals the number of elements of X which are fixed by g .

Let χ_1 be the principal character of G . We have

$$\begin{aligned}
 (\pi, \chi_1) &= \frac{1}{|G|} \sum_{g \in G} \pi(g) \chi_1(g) = \frac{1}{|G|} \sum_{g \in G} \pi(g) \\
 &= \frac{1}{|G|} \sum_{g \in G} |\{x \in X \mid gx = x\}| \\
 &= \frac{1}{|G|} |\{(g, x) \in G \times X \mid gx = x\}| \\
 &= \frac{1}{|G|} \sum_{x \in X} |\{g \in G \mid gx = x\}|.
 \end{aligned}$$

For each $x \in X$, let G_x be the stabilizer of x ; using Corollary 3.5, we now have

$$(\pi, \chi_1) = \sum_{x \in X} \frac{|G_x|}{|G|} = \sum_{\text{orbits } \mathcal{O}} \sum_{x \in \mathcal{O}} \frac{1}{|G : G_x|} = \sum_{\text{orbits } \mathcal{O}} \sum_{x \in \mathcal{O}} \frac{1}{|\mathcal{O}|} = \sum_{\text{orbits } \mathcal{O}} |\mathcal{O}| \cdot \frac{1}{|\mathcal{O}|} = \sum_{\text{orbits } \mathcal{O}} 1,$$

which allows us to conclude that (π, χ_1) is equal to the number of orbits of X under the action of G . (This result is often attributed to Burnside, although as argued in [21] it would be more accurate to call this result the Cauchy-Frobenius lemma.) In particular, if G acts transitively on X , then we see via Corollary 15.4 that the unique expression of the character $\pi - \chi_1$ as a linear combination of the irreducible characters of G does not involve χ_1 .

Example 15.5. The group G/G' is abelian by Proposition 2.6, and hence (as seen in Example 15.3) all of its irreducible characters are linear. It follows from Lemma 15.6 that each linear character of G/G' can be lifted to a linear character of G , and that the lifts of distinct characters are distinct. Now let χ be a linear character of G , so that $\chi: G \rightarrow \mathbb{C}^\times$ is a homomorphism whose kernel is K_χ . Then G/K_χ is abelian, being isomorphic with X , a subgroup of $\mathbb{C}^\times$, and hence $G' \leq K_\chi$ by Proposition 2.6. We can now define a linear character ψ of G/G' whose lift is χ by $\psi(gG') = \chi(g)$. We conclude that every linear character of G is the lift of a linear character of G/G' .

Example 15.6. Consider S_3 . The conjugacy classes of S_3 are

$$\{1\}, \quad \{(12), (13), (23)\}, \quad \{(123), (132)\}.$$

This follows from Proposition 1.10. We have $S_3/A_3 \cong \mathbb{Z}_2$, so S_3 has two linear characters, which are lifted from the characters of $\mathbb{Z}_2$ as in Lemma 15.6. Letting these characters be χ_1 and χ_2 , we have

	1	3	2
	1	(1 2)	(1 2 3)
χ_1	1	1	1
χ_2	1	-1	1
χ_3			

Since $6 = |S_3| = f_1^2 + f_2^2 + f_3^2 = 2 + f_3^2$ by Corollary 14.2, we have $f_3 = 2$. Column Orthogonality Theorem now gives

$$0 = \sum_{i=1}^3 f_i \chi_i((12)) = 1 \cdot 1 + 1 \cdot (-1) + 2\chi_3((12))$$

and

$$0 = \sum_{i=1}^3 f_i \chi_i((123)) = 1 \cdot 1 + 1 \cdot 1 + 2\chi_3((123)),$$

from which we see that the character table of S_3 is

	1	3	2
	1	(12)	(123)
χ_1	1	1	1
χ_2	1	-1	1
χ_3	2	0	-1

Example 15.7. Consider S_4 . Recall that S_4 has a normal subgroup of order 4, namely

$$K = \{1, (12)(34), (13)(24), (14)(23)\}.$$

The subgroup S_3 of S_4 intersects K trivially, and $|S_4| = |S_3||K|$; therefore $S_4 = K \rtimes S_3$. In particular, we have $S_4/K \cong S_3$, and thus we can obtain irreducible characters of S_4 from the irreducible characters of S_3 as in Lemma 15.6.

By Proposition 1.10, each conjugacy class of S_4 consists of all elements having a given cycle structure. Hence S_4 has 5 classes, with representatives 1, $(12)(34)$, (123) , (12) , and (1234) ; the orders of the classes are 1, 3, 8, 6, and 6, respectively.

Let χ_1, χ_2 , and χ_3 be the irreducible characters of S_4 obtained from those of S_3 . To determine these characters, we need information about the images in S_4/K of the class representatives. We easily see that the images of 1 and $(12)(34)$ are trivial, that the image of (123) has order 3, and that the images of (12) and (1234) have order 2.

Using this observation and the character table for S_3 obtained in Example 15.6, we obtain the following partial character table for S_4 :

	1	3	8	6	6
	1	(12)(34)	(123)	(12)	(1234)
χ_1	1	1	1	1	1
χ_2	1	1	1	-1	-1
χ_3	2	2	-1	0	0
χ_4					
χ_5					

Now S_4 acts transitively on the set $X = \{1, 2, 3, 4\}$; let π be the character of the $\mathbb{C}G$ -module $\mathbb{C}X$. Since $\pi(g)$ equals the number of fixed points under the action of g on X by Example 15.4, we have

	1	3	8	6	6
	1	(12)(34)	(123)	(12)	(1234)
π	4	0	1	2	0
$\pi - \chi_1$	3	-1	0	1	-1

We also have

$$(\pi - \chi_1, \pi - \chi_1) = \frac{1}{|S_4|} \sum_{i=1}^5 k_i [(\pi - \chi_1)(g_i)]^2 = \frac{1}{24} (1 \cdot 9 + 3 \cdot 1 + 8 \cdot 0 + 6 \cdot 1 + 6 \cdot 1) = 1,$$

and hence $\pi - \chi_1$ is irreducible by Corollary 15.3. Let $\chi_4 = \pi - \chi_1$. Now $\chi_2 \chi_4 \neq \chi_4$, and $\chi_2 \chi_4$ is irreducible by Proposition 15.5; therefore $\chi_5 = \chi_2 \chi_4$, and hence the character table of S_4 is

	1	3	8	6	6
	1	(12)(34)	(123)	(12)	(1234)
χ_1	1	1	1	1	1
χ_2	1	1	1	-1	-1
χ_3	2	2	-1	0	0
χ_4	3	-1	0	1	-1
χ_5	3	-1	0	-1	1

Example 15.8. Let G be a non-abelian group of order 8. We will show that this information completely determines the character table of G . Since there are two isomorphism classes of non-abelian groups of order 8 (see page 79), this will imply that the character table does not specify a group up to isomorphism.

We have $|Z(G)| = 2$ by Theorem 8.1 and Lemma 8.2, and hence $Z(G) \cong \mathbb{Z}_2$. Since G is non-abelian and $G/Z(G)$ is abelian (being a group of order 4), this forces $G' = Z(G)$ by Proposition 2.6. Now $|G/G'| = 4$, and by Lemma 8.2 G/G' cannot be cyclic since G is non-abelian and $G' = Z(G)$, so we must have $G/G' \cong \mathbb{Z}_2 \times \mathbb{Z}_2$. By Examples 15.3 and 15.5, we now know that G has exactly 4 linear characters. By Corollary 14.2, we have $\sum_{i=1}^r f_i^2 = 8$, where $f_i = 1$ for $1 \leq i \leq 4$ and $f_i > 1$ for $i > 4$; this forces $r = 5$ and $f_5 = 2$.

Let x be the generator of G' . As $G' = Z(G)$, we see that 1 and x are the only elements of G lying in single-element conjugacy classes, and hence each of the other three classes must have order 2. Let a, b , and c be representatives of the multi-element classes. Since G/G' also has four conjugacy classes, we see that the images of a, b , and c in G/G' must lie in distinct conjugacy classes. Using the character table for $\mathbb{Z}_2 \times \mathbb{Z}_2$ determined in Example 15.3, we can now obtain a partial character table for G by lifting:

	1	1	2	2	2
	1	x	a	b	c
χ_1	1	1	1	1	1
χ_2	1	1	1	-1	-1
χ_3	1	1	-1	1	-1
χ_4	1	1	-1	-1	1
χ_5	2				

We see via Column Orthogonality Theorem that $\chi_5(a) = \chi_5(b) = \chi_5(c) = 0$ and that $\chi_5(x) = -2$, completing the character table.

Example 15.9. Consider the group A_5 . Using Proposition 1.10, we find that the elements of A_5 form 4 conjugacy classes in S_5 , with representatives 1, $(12)(34)$, (123) , and (12345) and respective orders 1, 15, 20, and 24. However, elements of A_5 that are conjugate in S_5 may not be conjugate in A_5 .

Let $x \in A_5$, and let $\mathcal{K}$ be the conjugacy class in S_5 of x . We see from Proposition 3.13 that $|\mathcal{K}| = |S_5 : C_{S_5}(x)|$ and that the conjugacy class in A_5 of x is contained in $\mathcal{K}$ and has order $|A_5 : C_{A_5}(x)|$. Suppose first that $C_{S_5}(x) \not\leq A_5$. This forces $A_5 C_{S_5}(x) = S_5$ since A_5 is a maximal subgroup of S_5 . The first isomorphism theorem gives $S_5/A_5 = A_5 C_{S_5}(x)/A_5 \cong C_{S_5}(x)/C_{S_5}(x) \cap A_5 = C_{S_5}(x)/C_{A_5}(x)$, and hence

$$\begin{aligned} |A_5 : C_{A_5}(x)| &= \frac{|S_5 : C_{A_5}(x)|}{|S_5 : A_5|} = \frac{|S_5 : C_{A_5}(x)|}{|C_{S_5}(x) : C_{A_5}(x)|} = |S_5 : C_{S_5}(x)| \\ &= |\mathcal{K}|. \end{aligned}$$

Thus $\mathcal{K}$ is the conjugacy class in A_5 of x . Now suppose $C_{S_5}(x) \leq A_5$. Then we have $C_{A_5}(x) = C_{S_5}(x) \cap A_5 = C_{S_5}(x)$, and hence

$$|A_5 : C_{A_5}(x)| = |A_5 : C_{S_5}(x)| = \frac{1}{2} |S_5 : C_{S_5}(x)| = \frac{1}{2} |\mathcal{K}|,$$

from which we see that $\mathcal{K}$ splits into two conjugacy classes in A_5 of equal cardinality.

We easily see that each of the elements 1, $(12)(34)$, and (123) commutes with some odd permutation in S_5 ; for example, (45) and (123) commute. Hence by the above paragraph, the conjugacy classes in A_5 of these elements are the respective classes in S_5 . Since $24 \nmid 60$, we can now conclude that the conjugacy class in S_5 of (12345) splits into two classes in A_5 , and hence that the conjugacy classes of A_5 have orders 1, 15, 20, 12, and 12. (Using this fact and the remark made after Proposition 3.13, we could now give a slick proof of the simplicity of A_5 .) We will now show that $x = (12345)$ and $x^2 = (13524)$ are not conjugate in A_5 , thereby establishing that the conjugacy class in A_5 of x splits in A_5 into the class containing x and the class containing x^2 .

Suppose that $g \in A_5$ is such that $gxg^{-1} = x^2$. Then we have $g\langle x \rangle g^{-1} = \langle x^2 \rangle = \langle x \rangle$, and hence $g \in N_{A_5}(\langle x \rangle)$. Now $\langle x \rangle$ is a Sylow 5-subgroup of A_5 , and we observed in the proof of Theorem 7.4 that A_5 has 6 Sylow 5-subgroups. Thus $|N_{A_5}(\langle x \rangle)| = 10$, and we find that $N_{A_5}(\langle x \rangle) = \langle x \rangle \rtimes \langle t \rangle \cong D_{10}$, where $t = (25)(34)$. But $txt^{-1} = x^{-1} \neq x^2$; from this, we see that there is no element $g \in N_{A_5}(\langle x \rangle)$ such that $gxg^{-1} = x^2$. Contradiction; therefore x and x^2 are not conjugate in A_5 .

We now see that A_5 has 5 conjugacy classes, with representatives 1, $(12)(34)$, (123) , (12345) , and (13524) and respective orders 1, 15, 20, 12, and 12. Since A_5 is simple, we cannot produce irreducible

characters of A_5 by lifting irreducible characters of quotient groups, as we have done in previous examples. However, A_5 is a group of permutations, so we have as a starting point the characters of certain obvious permutation modules.

Let $X = \{1, 2, 3, 4, 5\}$; A_5 acts transitively on X in the obvious way. Let π be the character of the $\mathbb{C}A_5$ -module $\mathbb{C}X$. By Example 15.4, we see that $\pi(g)$ equals the number of fixed points of X under the action of $g \in A_5$, and hence we have

	1	15	20	12	12
	1	(12)(34)	(123)	(12345)	(13524)
π	5	1	2	0	0
$\pi - \chi_1$	4	0	1	-1	-1

We also have

$$(\pi - \chi_1, \pi - \chi_1) = \frac{1}{|A_5|} \sum_{i=1}^5 k_i [(\pi - \chi_1)(g_i)]^2 = \frac{1}{60} (1 \cdot 16 + 15 \cdot 0 + 20 \cdot 1 + 12 \cdot 1 + 12 \cdot 1) = 1,$$

and hence $\pi - \chi_1$ is irreducible by Corollary 15.3. Let $\chi_2 = \pi - \chi_1$.

Let Y be the set of two-element subsets of X ; this is a transitive A_5 -set under the obvious action. Let ψ be the character of the $\mathbb{C}A_5$ -module $\mathbb{C}Y$. We have

	1	15	20	12	12
	1	(12)(34)	(123)	(12345)	(13524)
ψ	10	2	1	0	0
$\psi - \chi_1$	9	1	0	-1	-1

Now

$$(\psi - \chi_1, \psi - \chi_1) = \frac{1}{|A_5|} \sum_{i=1}^5 k_i [(\psi - \chi_1)(g_i)]^2 = \frac{1}{60} (1 \cdot 81 + 15 \cdot 1 + 20 \cdot 0 + 12 \cdot 1 + 12 \cdot 1) = 2,$$

so by Corollary 15.3, $\psi - \chi_1$ is the sum of two irreducible characters; as noted in Example 15.4, neither of these two characters is χ_1 , since Y is transitive. We have

$$(\psi - \chi_1, \chi_2) = \frac{1}{|A_5|} \sum_{i=1}^5 k_i (\psi - \chi_1)(g_i) \chi_2(g_i) = \frac{1}{60} (1 \cdot 9 \cdot 4 + 15 \cdot 1 \cdot 0 + 20 \cdot 0 \cdot 1 + 24 \cdot (-1)(-1)) = 1,$$

and therefore by Corollary 15.4 we see that $\psi - \chi_1 - \chi_2$ is an irreducible character that is neither χ_1 nor χ_2 . Let $\chi_3 = \psi - \chi_1 - \chi_2$, and observe that $f_3 = 5$.

By Corollary 14.2 we have $\sum_{i=1}^5 f_i^2 = |A_5| = 60$, which gives $f_4^2 + f_5^2 = 18$ and hence $f_4 = f_5 = 3$. Thus we have the following partial character table:

	1	15	20	12	12
	1	(12)(34)	(123)	(12345)	(13524)
χ_1	1	1	1	1	1
χ_2	4	0	1	-1	-1
χ_3	5	1	-1	0	0
χ_4	3				
χ_5	3				

Let $a = \chi_4((12)(34))$ and $b = \chi_5((12)(34))$. By Column Orthogonality Theorem, we have

$$0 = \sum_{i=1}^5 f_i \chi_i((12)(34)) = 1 \cdot 1 + 4 \cdot 0 + 5 \cdot 1 + 3a + 3b,$$

and hence $a + b = -2$. By part (iii) of Proposition 14.4, we see that each of a and b is a sum of three square roots of unity; therefore $a, b \in \{-3, -1, 1, 3\}$. But by Column Orthogonality Theorem we have

$$4 = \frac{60}{15} = \frac{|A_5|}{k_2} = \sum_{i=1}^5 |\chi_i((12)(34))|^2 = 1 + 0 + 1 + a^2 + b^2,$$

and hence $a^2 + b^2 = 2$. It now follows that $a = b = -1$.

By Column Orthogonality Theorem, we have

$$3 = \frac{60}{20} = \frac{|A_5|}{k_3} = \sum_{i=1}^5 |\chi_i((1\ 2\ 3))|^2 = 1 + 1 + 1 + |\chi_4((1\ 2\ 3))|^2 + |\chi_5((1\ 2\ 3))|^2;$$

thus $|\chi_4((1\ 2\ 3))|^2 + |\chi_5((1\ 2\ 3))|^2 = 0$, and therefore we conclude that $\chi_4((1\ 2\ 3)) = \chi_5((1\ 2\ 3)) = 0$. We now have the following partial character table:

	1	15	20	12	12
	1	(1 2)(3 4)	(1 2 3)	(1 2 3 4 5)	(1 3 5 2 4)
χ_1	1	1	1	1	1
χ_2	4	0	1	-1	-1
χ_3	5	1	-1	0	0
χ_4	3	-1	0		
χ_5	3	-1	0		

Let $x = (1\ 2\ 3\ 4\ 5)$. Recall that in proving that x and x^2 are not conjugate in A_5 , we found that x and x^{-1} are conjugate in A_5 ; hence $\chi_4(x) = \chi_4(x^{-1})$, and thus $\chi_4(x)$ is real-valued by part (iv) of Proposition 14.4. The same argument shows that $\chi_4(x^2)$, $\chi_5(x)$, and $\chi_5(x^2)$ are also real-valued. Let $c = \chi_4(x)$ and $d = \chi_4(x^2)$. Then by Row Orthogonality Theorem, we have

$$0 = \sum_{i=1}^5 k_i \chi_4(g_i) = 1 \cdot 3 + 15 \cdot (-1) + 20 \cdot 0 + 12c + 12d,$$

which gives $c + d = 1$, and also (since χ_4 is real-valued)

$$60 = \sum_{i=1}^5 k_i \chi_4(g_i)^2 = 1 \cdot 9 + 15 \cdot 1 + 20 \cdot 0 + 12c^2 + 12d^2,$$

which gives $c^2 + d^2 = 3$. Hence $c^2 - c - 1 = 0$, so without loss of generality we have $c = \frac{1+\sqrt{5}}{2}$ and $d = 1 - c = \frac{1-\sqrt{5}}{2}$. We can similarly show that we must have $\chi_5(x), \chi_5(x^2) \in \{\frac{1+\sqrt{5}}{2}, \frac{1-\sqrt{5}}{2}\}$; we leave it to the reader to verify that $\chi_5(x) = \frac{1-\sqrt{5}}{2}$ and $\chi_5(x^2) = \frac{1+\sqrt{5}}{2}$, which completes the character table of A_5 .

Exercises

Throughout these exercises, G denotes a finite group, and $\mathcal{X}$ denotes the character table of G .

Exercise 15.1. What is the determinant of $\mathcal{X}$?

Solution.

□

Exercise 15.2. Show that the sum of the entries in any row of $\mathcal{X}$ is a non-negative integer.

Solution.

□

Exercise 15.3. Compute the character of the G -set G , where G acts via conjugation, and determine the multiplicities in this character of the irreducible characters.

Solution.

□

Exercise 15.4. Observe that the complex conjugate of an irreducible character is an irreducible character. (This is either Exercise 14.6 or an easy consequence of part (ii) of Proposition 14.8 and Corollary 15.3.) Let P be the $r \times r$ permutation matrix such that $P\mathcal{X}$ is the matrix obtained from $\mathcal{X}$ by interchanging two rows if their corresponding characters are conjugate. Let Q be the $r \times r$ permutation matrix such that $\mathcal{X}Q$ is the matrix obtained from $\mathcal{X}$ by interchanging two columns if their corresponding conjugacy classes are inverse, in the sense that one class consists of the inverses of the elements of the other class. Show that $P\mathcal{X} = \mathcal{X}Q$.

Solution.

□

Exercise 15.5. Suppose that $|G|$ is odd. Show that the only irreducible character that is real-valued is the principal character.

Solution.

□

Exercise 15.6. Let χ be an irreducible character of G , and let $e \in \mathbb{C}G$ be the identity element of the corresponding matrix summand of $\mathbb{C}G$. Show that

$$e = \frac{\chi(1)}{|G|} \sum_{g \in G} \chi(g^{-1})g.$$

Solution.

□

Exercise 15.7. If $N \trianglelefteq G$ and $g \in G$, show that $|C_G(g)| \geq |C_{G/N}(gN)|$.

Solution.

□

Exercise 15.8. Show that if G acts doubly transitively on X , then the character of $\mathbb{C}X$ is the sum of the principal character and exactly one other irreducible character.

Solution.

□

Exercise 15.9. Determine the character table of D_{2n} for any $n \in \mathbb{N}$.

Solution.

□

Exercise 15.10. Determine the character table of S_5 .

Solution.

□

16 Induction

Overview

- Induced modules play a critical role in the representation theory of finite groups.
- Two equivalent definitions of induction are presented: one via tensor products over arbitrary rings (slick but heavier), the other using only tensor products over fields (clumsier but more elementary). A third definition is developed in Exercise 16.1.
- Let $H \leq G$ and let V be an FH -module. Since the group algebra FG is an (FG, FH) -bimodule, we form the FG -module $FG \otimes_{FH} V$, the *induced* FG -module of V (or the *induction* of V to G), denoted $\text{Ind}_H^G V$. Other notations: V^G , $V \uparrow^G$, and $V \uparrow_H^G$.
- For an FG -module U , regarding U as an FH -module yields the *restriction* $\text{Res}_H^G U$. Other notations: U_H , $U \downarrow_H$, and $U \downarrow_H^G$.

Lemma 16.1. *We have $\dim_F \text{Ind}_H^G V = |G : H| \dim_F V$ for V an FH -module. Moreover, as F -vector spaces we have*

$$\text{Ind}_H^G V = \bigoplus_{t \in T} t \otimes V,$$

where T is a transversal for H in G and $t \otimes V = \{t \otimes v \mid v \in V\}$.

Proof. □

Theorem (Frobenius Reciprocity Theorem). *Let U be an FG -module, let $H \leq G$, and let V be an FH -module. Then as F -vector spaces we have $\text{Hom}_{FH}(V, \text{Res}_H^G U) \cong \text{Hom}_{FG}(\text{Ind}_H^G V, U)$.*

Proof. □

Induced Characters

- Restrict attention to $F = \mathbb{C}$. If V is a $\mathbb{C}H$ -module with character φ , the character of $\text{Ind}_H^G V$ is denoted φ^G and called an *induced character*; we have $\varphi^G(1) = |G : H| \varphi(1)$.
- If U is a $\mathbb{C}G$ -module with character χ , the character of $\text{Res}_H^G U$ is denoted $\chi|_H$, and $\chi|_H(1) = \chi(1)$.
- The form of “Frobenius reciprocity” actually known to Frobenius (Theorem 16.2) follows from Frobenius Reciprocity Theorem and Theorem 14.12.

Theorem 16.2. *Let $H \leq G$, let U be a $\mathbb{C}G$ -module having character χ , and let V be a $\mathbb{C}H$ -module having character φ . Then $(\varphi, \chi|_H)_H = (\varphi^G, \chi)_G$.*

Proof. □

Induction–Restriction Tables

- If $H \leq G$ and $\varphi_1, \dots, \varphi_r$ and $\chi_1, \dots, \chi_s$ are the irreducible characters of H and G respectively, the *induction–restriction table* of (G, H) is the $r \times s$ matrix whose (i, j) -entry gives the common value of $(\varphi_i^G, \chi_j)_G$ and $(\varphi_i, \chi_j|_H)_H$.
- By Corollary 15.4, the i th row gives the multiplicities with which the χ_j appear in the decomposition of φ_i^G , while the j th column gives the multiplicities with which the φ_i appear in the decomposition of $\chi_j|_H$.
- One can compute the table by calculating the restrictions of the χ_j and reading off the expressions for the φ_i^G in terms of the χ_j .
- For example, with $G = S_3$ and $H = S_2 \leq G$, inspection gives $\chi_1|_H = \varphi_1$, $\chi_2|_H = \varphi_2$, $\chi_3|_H = \varphi_1 + \varphi_2$, so the induction–restriction table yields $\varphi_1^G = \chi_1 + \chi_3$ and $\varphi_2^G = \chi_2 + \chi_3$.

Proposition 16.3. *Let $H \leq G$, and let V be a $\mathbb{C}H$ -module having character χ . If T is a transversal for H in G , then for any $g \in G$ we have*

$$\chi^G(g) = \sum_{\substack{t \in T, \\ t^{-1}gt \in H}} \chi(t^{-1}gt).$$

Proof. □

Corollary 16.4. *Let χ be a character of $H \leq G$, and let $g \in G$. Then*

$$\chi^G(g) = \frac{1}{|H|} \sum_{\substack{x \in G, \\ x^{-1}gx \in H}} \chi(x^{-1}gx).$$

Proof. □

Proposition 16.5. *Let χ be a character of $H \leq G$, and let $g \in G$, and let s be the number of conjugacy classes of H whose members are conjugate in G to g . If $s = 0$, then $\chi^G(g) = 0$. Otherwise, if we let $h_1, \dots, h_s$ be representatives of these s conjugacy classes of H , then*

$$\chi^G(g) = \sum_{i=1}^s \frac{|C_G(g)|}{|C_H(h_i)|} \chi(h_i).$$

Proof. □

Corollary 16.6. *Let χ be a character of $H \leq G$. Let $g \in G$, and suppose that the number s of conjugacy classes of H whose members are conjugate in G to g is positive. Let ℓ be the order of the conjugacy class in G of g ; let $h_1, \dots, h_s$ be representatives of these s conjugacy classes of H , and let $k_1, \dots, k_s$ be the orders of these classes. Then*

$$\chi^G(g) = \sum_{i=1}^s |G : H| \frac{k_i}{\ell} \chi(h_i).$$

Proof. □

Frobenius Groups

- We end the section by presenting an important theorem on finite groups which, quite remarkably, has no known proof that does not use character theory.
- A *Frobenius group* is a group G having a subgroup H such that $H \cap gHg^{-1} = 1$ for every $g \in G - H$. We call H a *Frobenius complement* of G .
- If G is a finite Frobenius group, applying Frobenius' Theorem yields a normal subgroup N , called the *Frobenius kernel* of G . One has $G = N \rtimes H$, $N \cap H = 1$, and $|N||H| = |G|$.
- For $G = N \rtimes H$ a finite Frobenius group with Frobenius kernel N and complement H , the conjugation map $\varphi : H \rightarrow \text{Aut}(N)$ has the property that for each $1 \neq h \in H$, $\varphi(h)$ is a *fixed-point-free automorphism* of N : $\varphi(h)(n) = n$ forces $n = 1$. In particular, φ is injective.
- Conversely, given finite groups N and H and a homomorphism $\varphi : H \rightarrow \text{Aut}(N)$ for which every $\varphi(h)$, $h \neq 1$, is fixed-point-free, $N \rtimes_{\varphi} H$ is a Frobenius group.
- Example: if p, q are distinct primes with $p \equiv 1 \pmod{q}$, then any monomorphism $\varphi : \mathbb{Z}_q \rightarrow \text{Aut}(\mathbb{Z}_p)$ is fixed-point-free, and hence the unique non-abelian group of order pq is a Frobenius group.
- A theorem of Thompson (his 1959 Chicago Ph.D. thesis) states that any finite group having a fixed-point-free automorphism of prime order is nilpotent. Hence the Frobenius kernel of a finite Frobenius group is nilpotent.

Theorem (Frobenius' Theorem). *Let G be a transitive permutation group on a finite set X , and suppose that each non-identity element of G fixes at most one element of X . Then the union of the identity element and the set of elements of G that have no fixed points is a normal subgroup of G .*

Proof. □

Exercises

Throughout these exercises, G is a finite group, H is a subgroup of G , F is a field, and all FG -modules are finitely generated.

Exercise 16.1. Let V be an FH -module, and let W be an FG -module containing V . Suppose that W has the property that for any FG -module U and any $\varphi \in \text{Hom}_{FH}(V, U)$, there is a unique FG -module homomorphism from W to U which extends φ . (In this case, we say that W is relatively H -free with respect to V ; compare with Exercises 13.1–13.2.) Show that $W \cong \text{Ind}_H^G V$.

Solution. □

Exercise 16.2. Let W be an FG -module that is generated by an FH -submodule V , and suppose that $\dim_F W = |G : H| \dim_F V$. Show that we must have $W \cong \text{Ind}_H^G V$.

Solution. □

Exercise 16.3. Show that $\text{Hom}_{FG}(U, \text{Ind}_H^G V) \cong \text{Hom}_{FH}(\text{Res}_H^G U, V)$ as F -vector spaces for any FG -module U and FH -module V . (HINT: Show first that if $\varphi \in \text{Hom}_{FH}(\text{Res}_H^G U, V)$, then the map sending $u \in U$ to $\sum_{t \in T} t \otimes \varphi(t^{-1}u)$, where T is a transversal for H in G , lies in $\text{Hom}_{FG}(U, \text{Ind}_H^G V)$.)

Solution. □

Exercise 16.4. Show that if X is a transitive G -set, then $FX \cong \text{Ind}_{G_x}^G F$ for any $x \in X$.

Solution. □

Exercise 16.5. Compute the induction-restriction table of (A_5, A_4) .

Solution. □

Exercise 16.6. Compute the character table of the unique non-abelian group of order pq , where p and q are distinct primes and $p \equiv 1 \pmod{q}$.

Solution. □

Exercise 16.7. Suppose that G is a Frobenius group having Frobenius kernel N . Show that the irreducible characters of G either are lifts of irreducible characters of G/N or are induced from the non-principal irreducible characters of N . When is it true that the inductions to G of two distinct non-principal irreducible characters of N are equal?

Solution. □

Further Exercises

In this set of exercises, we develop the character table of $G = \text{GL}(2, q)$, where q is any prime power. Recall from Proposition 4.2 that we have $|G| = q(q-1)^2(q+1)$. Our first task is to determine the conjugacy classes of G . Here we sketch a comparatively “hands-on” approach, which uses only some standard topics from linear algebra. (A slicker approach might, for instance, make use of the invariant factor formulation of rational canonical form.) We denote by $m_g(X)$ the minimal polynomial of an element $g \in G$. Our strategy is to exploit the following:

Exercise 16.8. Show that conjugate elements of G have the same minimal polynomial.

Solution. □

Since $m_g(X)$ is a factor of the characteristic polynomial of g , which is the quadratic polynomial $\det(XI - g)$, we see that $m_g(X)$ is either linear, or a product of two like or unlike linear factors, or an irreducible monic quadratic.

Exercise 16.9. (cont.) Show that if $m_g(X) = X - a$ for some $a \in \mathbb{F}_q^\times$, then $g = \begin{pmatrix} a & 0 \\ 0 & a \end{pmatrix} \in Z(G)$.

Solution. □

Exercise 16.10. (cont.) Show that if $m_g(X) = (X - a)^2$ for some $a \in \mathbb{F}_q^\times$, then g is conjugate with $\begin{pmatrix} a & 1 \\ 0 & a \end{pmatrix}$, and the conjugacy class of g has order $q^2 - 1$.

Solution. □

Exercise 16.11. (cont.) Show that if $m_g(X) = (X - a)(X - b)$ for some distinct $a, b \in \mathbb{F}_q^\times$, then g is conjugate with $\begin{pmatrix} a & 0 \\ 0 & b \end{pmatrix}$, and the conjugacy class of g has order $q^2 + q$.

Solution. □

Exercise 16.12. (cont.) Show that if $r, s \in \mathbb{F}_q$ are such that $X^2 - rX - s$ is irreducible, then the conjugacy class of $\begin{pmatrix} 0 & 1 \\ s & r \end{pmatrix}$ has order $q^2 - q$.

Solution. □

(Hint for Exercises 16.10–16.12: Compute centralizers and use Proposition 3.13.)

Exercise 16.13. (cont.) Verify that what follows is a complete list of the conjugacy classes of G : $q - 1$ classes of 1 element each, indexed by elements of $\mathbb{F}_q^\times$; $q - 1$ classes of $q^2 - 1$ elements each, indexed by elements of $\mathbb{F}_q^\times$; $(q - 1)(q - 2)/2$ classes of $q^2 + q$ elements each, indexed by unordered pairs of distinct elements of $\mathbb{F}_q^\times$; and $q(q - 1)/2$ classes of $q^2 - q$ elements each, indexed by irreducible monic quadratic polynomials with coefficients in $\mathbb{F}_q$.

Solution. □

Having determined the classes of G , we now turn to finding irreducible characters. Recall that we have an epimorphism $\det : G \rightarrow \mathbb{F}_q^\times$. Consequently, we can lift characters of $\mathbb{F}_q^\times$ to characters of G , as in Lemma 15.6. Now $\mathbb{F}_q^\times$ is an abelian group of order $q - 1$, and hence it has $q - 1$ linear characters, which we denote by $\tau_1, \dots, \tau_{q-1}$. (In fact, it is true (see [17, p. 132]) that $\mathbb{F}_q^\times \cong \mathbb{Z}_{q-1}$, and so we can, upon fixing a generator of $\mathbb{F}_q^\times$, completely specify the τ_i .) For each $1 \leq i \leq q - 1$, let $\chi_i = \tau_i \circ \det$. We have

$$\begin{array}{c|cccc} & \begin{pmatrix} a & 0 \\ 0 & a \end{pmatrix} & \begin{pmatrix} a & 1 \\ 0 & a \end{pmatrix} & \begin{pmatrix} a & 0 \\ 0 & b \end{pmatrix} & \begin{pmatrix} 0 & 1 \\ s & r \end{pmatrix} \\ \hline \chi_i & \tau_i(a)^2 & \tau_i(a)^2 & \tau_i(a)\tau_i(b) & \tau_i(-s) \end{array}$$

The characters χ_i are linear, and χ_1 is indeed the principal character of G .

As in Chapter 2, let B , U , and T be the subgroups of G consisting, respectively, of all upper triangular, all upper unitriangular, and all diagonal elements of G . We have $T \cong \mathbb{F}_q^\times \times \mathbb{F}_q^\times$, and so we see from Example 15.2 that the irreducible characters of T have the form $\tau_i \times \tau_j$ (after making appropriate identifications). Now $B = U \rtimes T$ by Proposition 5.1, so we can lift each character $\tau_i \times \tau_j$ of $T \cong B/U$ to

obtain a character θ_{ij} of B . The epimorphism from B to T sends $\begin{pmatrix} x & y \\ 0 & z \end{pmatrix}$ to $\begin{pmatrix} x & 0 \\ 0 & z \end{pmatrix}$, and consequently

we have $\theta_{ij} \begin{pmatrix} x & y \\ 0 & z \end{pmatrix} = (\tau_i \times \tau_j) \begin{pmatrix} x & 0 \\ 0 & z \end{pmatrix} = \tau_i(x)\tau_j(z)$. We now wish to consider the induced characters θ_{ij}^G .

Exercise 16.14. (cont.) Show that

$$\begin{array}{c|cccc} & \begin{pmatrix} a & 0 \\ 0 & a \end{pmatrix} & \begin{pmatrix} a & 1 \\ 0 & a \end{pmatrix} & \begin{pmatrix} a & 0 \\ 0 & b \end{pmatrix} & \begin{pmatrix} 0 & 1 \\ s & r \end{pmatrix} \\ \hline \theta_{ij}^G & (q+1)\tau_i(a)\tau_j(a) & \tau_i(a)\tau_j(a) & \tau_i(a)\tau_j(b) + \tau_i(b)\tau_j(a) & 0 \end{array}$$

Observe that $\theta_{ij}^G = \theta_{ji}^G$.

Solution. □

Exercise 16.15. (cont.) Show that $(\theta_{ij}^G, \theta_{ij}^G) = 1 + \delta_{ij}$ and that $(\theta_{ii}^G, \chi_i) = 1$. Conclude that $\theta_{ii}^G - \chi_i$ is an irreducible character of G for each i and that θ_{ij}^G is an irreducible character of G whenever i and j are distinct.

Solution. □

At this point, we have constructed $q-1$ distinct irreducible characters of degree 1 (the χ_i), $q-1$ distinct irreducible characters of degree q (the $\theta_{ii}^G - \chi_i$), and $(q-1)(q-2)/2$ distinct irreducible characters of degree $q+1$ (the θ_{ij}^G for $i < j$). Our partial character table of G is now

	$\begin{pmatrix} a & 0 \\ 0 & a \end{pmatrix}$	$\begin{pmatrix} a & 1 \\ 0 & a \end{pmatrix}$	$\begin{pmatrix} a & 0 \\ 0 & b \end{pmatrix}$	$\begin{pmatrix} 0 & 1 \\ s & r \end{pmatrix}$
χ_i	$\tau_i(a)^2$	$\tau_i(a)^2$	$\tau_i(a)\tau_i(b)$	$\tau_i(-s)$
$\theta_{ii}^G - \chi_i$	$q\tau_i(a)^2$	0	$\tau_i(a)\tau_i(b)$	$-\tau_i(-s)$
θ_{ij}^G	$(q+1)\tau_i(a)\tau_j(a)$	$\tau_i(a)\tau_j(a)$	$\tau_i(a)\tau_j(b) + \tau_i(b)\tau_j(a)$	0

To complete the character table, it will be convenient to work with a different set of representatives for the conjugacy classes of those elements whose minimal polynomials are irreducible quadratics. (We shall refer to these classes as being the “fourth-column” classes, owing to their placement in our partial character table above.) By Exercise 4.4, G has a subgroup C which is isomorphic with $\mathbb{F}_{q^2}^\times$, where $\mathbb{F}_{q^2}$ is the field of q^2 elements. As we remarked before, we have $C \cong \mathbb{Z}_{q^2-1}$. It is true, although we are not in a position to prove it, that any subgroup of G that is isomorphic with $\mathbb{F}_{q^2}^\times$ is a conjugate of C . (The proof of this fact requires some familiarity with field theory, but the essential point is that all fields of order q^2 are isomorphic; see [19, Corollary XIV.2.7].) It is also true that C must contain $Z(G) \cong \mathbb{F}_q^\times$, and so we will consider $\mathbb{F}_q^\times$ as being a subgroup of C .

Exercise 16.16. (cont.) Show that the elements of $C - \mathbb{F}_q^\times$ form a double set of representatives of the fourth-column classes, with $x \in C - \mathbb{F}_q^\times$ being conjugate with (but unequal to) x^q . (HINT: Argue that each element of G whose minimal polynomial is an irreducible quadratic determines a subgroup of G that is isomorphic with $\mathbb{F}_{q^2}^\times$.)

Solution. □

We now take $x \in C - \mathbb{F}_q^\times$ as our arbitrary fourth-column class representative. We have $\chi_i(x) = \tau_i(\det x)$, $(\theta_{ii}^G - \chi_i)(x) = -\tau_i(\det x)$, and $\theta_{ij}^G(x) = 0$.

Let $\beta_1, \dots, \beta_{q^2-1}$ be the irreducible characters of C . (By choosing a generator for C , we can specify the β_i .) We will consider the induced characters β_k^G .

Exercise 16.17. (cont.) Show

	$\begin{pmatrix} a & 0 \\ 0 & a \end{pmatrix}$	$\begin{pmatrix} a & 1 \\ 0 & a \end{pmatrix}$	$\begin{pmatrix} a & 0 \\ 0 & b \end{pmatrix}$	$x \in C - \mathbb{F}_q^\times$
β_k^G	$(q^2 - q)\beta_k(a)$	0	0	$\beta_k(x) + \beta_k(x)^q$

Observe that $(\beta_k^q)^G = \beta_k^G$, where by β_k^q we mean the linear character of C defined by $\beta_k^q(x) = \beta_k(x)^q$.

Solution. □

It is not hard to show that there are exactly $q-1$ values of k for which $\beta_k^q = \beta_k$. (This follows, for instance, from the observation made in Example 15.1 that the linear characters of a cyclic group of order n themselves form a cyclic group of order n .)

Choose some k such that $\beta_k^q \neq \beta_k$, and let i_k be such that the restriction of β_k to $\mathbb{F}_q^\times \subseteq C$ is τ_{i_k} . Let $\psi_k = (\theta_{11}^G - \chi_1)\theta_{i_k 1}^G - \theta_{i_k 1}^G - \beta_k^G$; this is a virtual character.

Exercise 16.18. (cont.) Show that

	$\begin{pmatrix} a & 0 \\ 0 & a \end{pmatrix}$	$\begin{pmatrix} a & 1 \\ 0 & a \end{pmatrix}$	$\begin{pmatrix} a & 0 \\ 0 & b \end{pmatrix}$	$x \in C - \mathbb{F}_q^\times$
ψ_k	$(q-1)\tau_{i_k}(a)$	$-\tau_{i_k}(a)$	0	$-\beta_k(x) - \beta_k(x)^q$

Solution. □

Exercise 16.19. (cont.) If k is such that $\beta_k^q \neq \beta_k$, show that $(\psi_k, \psi_k) = 1$. Conclude that we have constructed $q(q-1)/2$ distinct irreducible characters of degree $q-1$.

Solution. □

The number of irreducible characters that we have constructed is now equal to the number of conjugacy classes of G , and so by Theorem 14.3 we are done.

Appendix A

Algebraic Integers and Characters

17 Algebraic Integers and Characters

Setting and motivation

- This appendix continues character theory with a different flavour from Chapter 6, leaning on Galois theory and a higher level of algebraic background.
- Logically the appendix follows Section 15.
- Throughout, G is a finite group.
- We say that $x \in \mathbb{C}$ is an *algebraic integer* if there is some monic polynomial $f \in \mathbb{Z}[X]$ such that $f(x) = 0$.

Lemma 17.1. *The set of rational numbers that are also algebraic integers is exactly the set of ordinary integers.*

Proof.

□

Algebraic integers form a ring

- The next standard result is used freely; its proof is sketched in the exercises.

Proposition 17.2. *The algebraic integers form a subring of $\mathbb{C}$.*

Proof.

□

Characters take algebraic-integer values

- The relevance of algebraic integers to representation theory of finite groups is established by the following basic fact.

Proposition 17.3. *Let χ be a character of G . Then $\chi(g)$ is an algebraic integer for any $g \in G$.*

Proof.

□

Class-sum integrality

- The next result is necessary for both intended applications.

Lemma 17.4. *If χ is an irreducible character of G and $g \in G$, then $|G : C_G(g)|\chi(g)/\chi(1)$ is an algebraic integer.*

Proof.

□

Degrees of irreducible characters divide $|G|$

- This proves the first main result of the appendix, a fact already mentioned in Section 14.

Proposition 17.5. *Let χ be an irreducible character of G . Then $\chi(1)$ divides $|G|$.*

Proof. □

Toward Burnside's $p^a q^b$ theorem

- The goal of the remainder of the appendix is to prove Burnside's Theorem on the solvability of groups of order $p^a q^b$, mentioned in Section 11.

Lemma 17.6. *Let χ be a character of G , let $g \in G$, and define $\gamma = \chi(g)/\chi(1)$. If γ is a non-zero algebraic integer, then $|\gamma| = 1$.*

Proof. □

A classical theorem of Burnside

- We now prove a classical theorem due to Burnside.

Theorem 17.7. *If G has a conjugacy class of non-trivial prime power order, then G is not simple.*

Proof. □

Burnside's famous $p^a q^b$ theorem

- Finally, we have Burnside's famed $p^a q^b$ theorem.

Theorem (Burnside's Theorem). *If $|G| = p^a q^b$, where p and q are primes, then G is solvable.*

Proof. □

Beyond two primes

- It was observed on page 100 that there are non-solvable groups of order $p^a q^b r^c$, where p , q , and r are distinct primes.
- In fact, up to isomorphism there are eight such groups.
- Six of these groups are projective special linear groups. (Recall that A_5 and A_6 are, by Exercises 6.2 and 6.5, isomorphic with projective special linear groups.)
- The remaining two groups are members of a family of groups called the projective special unitary groups (see [24, p. 245]).

Exercises

Exercise 17.1. For $x \in \mathbb{C}$, let $\mathbb{Z}[x] = \{f(x) \mid f(X) \in \mathbb{Z}[X]\}$; this is an abelian subgroup of $\mathbb{C}$. Show that x is an algebraic integer iff $\mathbb{Z}[x]$ is finitely generated.

Solution. □

Exercise 17.2. (cont.) Show that the set of algebraic integers is a subring of $\mathbb{C}$. (HINT: Use Exercise 17.1 and the well-known fact (see [1, p. 145]) that any subgroup of a finitely generated abelian group is finitely generated.)

Solution. □

Exercise 17.3. Let G be a finite group, and define a function $\psi: G \rightarrow \mathbb{C}$ by setting $\psi(g) = |\{(x, y) \in G \times G \mid [x, y] = g\}|$ for $g \in G$. Show that

$$\psi = \sum_{i=1}^r \frac{|G|}{\chi_i(1)} \chi_i,$$

which in light of Proposition 17.5 shows that ψ is a character. (Compare with Exercise 3.6.)

Solution. □