



# Uniqueness Theory of Meromorphic Functions Sharing Weighted Values with its $k$ -th derivative

---

Zhanglong CAO

- **Self introduction**
- **Preliminaries**
- **Background**
- **Proof & Conclusion**
- **References**


- **Self introduction**

- Zhanglong CAO, known as Jerome
- 1st year PhD student in statistics, University of Otago
- Worked in Beijing as Data analyst
- Masters in Complex Analysis, USST
- Bachelors in Mathematics, JSNU

- **Preliminaries**
- **Meromorphic Functions**

In the mathematical field of complex analysis, a meromorphic function on an open subset  $D$  of the complex plane is a function that is holomorphic on all  $D$  except a set of isolated points (the poles of the function), at each of which the function must have a Laurent series.

The gamma function is meromorphic in the whole complex plane.



- **Sharing Values**

Let  $f$  and  $g$  be two non-constant meromorphic functions and let  $a$  be a finite complex number.

We say that  $f$  and  $g$  share the value  $a$  CM, provided that  $f - a$  and  $g - a$  have the same zeros with the same multiplicities.

Similarly, we say that  $f$  and  $g$  share the value  $a$  IM, provided that  $f - a$  and  $g - a$  have the same zeros ignoring multiplicities.

- **Weighted Values**

Let  $k$  be a nonnegative integer or infinity. For any  $a \in \mathbb{C} \cup \{\infty\}$ , we denote by  $E_k(a, f)$  the set of all  $a$ -points of  $f$ , where an  $a$ -point of multiplicity  $m$  is counted  $m$  times if  $m \leq k$ , and  $k + 1$  times if  $m > k$ .

If  $E_k(a, f) = E_k(a, g)$ , we say that  $f$  and  $g$  share the value  $a$  with weight  $k$ .

We write  $f, g$  share  $(a, k)$  to mean that  $f, g$  share the value  $a$  with weight  $k$ .

We also note that  $f, g$  share a value  $a$  IM or CM if and only if  $f, g$  share  $(a, 0)$  or  $(a, \infty)$ , respectively.



- **Background**

- In 1980, Gundersen proved the following theory.
- If  $f$  is a non-constant meromorphic function,  $b(\neq 0)$  is a finite value. If  $f$  and  $f'$  share 0 and  $b$  CM, then  $f \equiv f'$ .

- In 1986, Frank-Ohlenroth proved the following,
- $f$  is a non-constant meromorphic function. If  $f$  and  $f^{(k)}$  share two distinct non-zero finite complex values  $a$  and  $b$  CM, then  $f \equiv f^{(k)}$ .
- In the same year, Frank-Weissenborn proved that, for any finite distinct complex values  $a$  and  $b$ , if  $f$  and  $f^{(k)}$  share them CM, then  $f \equiv f^{(k)}$ .

Here comes a conjecture,

Will the equation  $f \equiv f^{(k)}$  still hold if the condition  $f$  and  $f^{(k)}$  share  $b$  CM, changes to  $f$  and  $f^{(k)}$  share  $b$  IM?

Unfortunately, the conjecture is false.


Luckily, I studied in this area and was interested in this topic.

So the theorem is as following,

Let  $f(z)$  be a non-constant meromorphic function,  $k$  is a positive integer and the multiplicity of poles  $\geq 2k + 4$ . If  $f(z)$  and  $f^{(k)}(z)$  share  $(a, \infty)$ ,  $(b, 1)$ , where  $a$  and  $b$  are distinct finite complex numbers, then we have

$$f(z) \equiv f^{(k)}(z)$$

# • Proof



# • References

- [1] Yang, C.J. Yi, H.X. Uniqueness Theory of Meromorphic Functions. Beijing: science press, 2003.
- [2] 杨乐. 值分布论及其新研究. 北京:科学出版社, 1982.
- [3] Nevanlinna, R. Le Theoreme de Picard-borel et la theorie des fonctions meromorphes, Paris, 1929.
- [4] Yi, H.X. Lü, W.R. Unicity theorems of meromorphic functions, Proc. Japan Acad., Ser .A. 2002, 78:152-156.
- [5] Lahiri, I. Weighted sharing and uniqueness of meromorphic functions, Nagoya Math. J. 2001, 161:193-206.
- [6] Lahiri, I. Weighted sharing of three values and uniqueness of meromorphic functions, Kodai Math. J. 2001, 24:421-435.
- [7] Lahiri, I. Weighted value sharing and uniqueness of meromorphic functions, Complex Variables Theory Appl. 2001.46:241-253.
- [8] Frank, G. Weissenborn, G. Meromorphe Funktionen, die mit einer ihrer Ableitungen Werte teilen, Complex Variables, 1986, 7:33-43.
- [9] 陈春芳. 亚纯函数及其导数的唯一性. 南京师大学报(自然科学版) 2004, **27**(3): 36-39.
- [10] Zhang, Q.C. Meromorphic functions sharing three values[J]. India J. Pure Appl. Math, 1999, **30**(7):667-682.
- [11] 李效敏, 肖永火. 具有2个公共小函数的整函数及其k阶导数的唯一性. 中国海洋大学学报. 2008, **38**(2):335-339.
- [12] Lahiri, I. On a result of Ozawa concerning uniqueness of meromorphic functions II, J. Math. Anal. Appl. 2003, 283:66-76.
- [13] Hayman, W.K. Meromorphic Functions, Clarendon, Oxford, 1964.
- [14] Yi, H.X. Unicity theorems for meromorphic functions that share three values, Kodai Math. J. 1995, 18:300-314.
- [15] Han, Q. Yi, H.X. On meromorphic functions that share three values of finite weights. J. Math. Anal. Appl. 2007, 334:314-332
- [16] Hu, P.C. Li, P. Yang, C.C. Unicity of Meromorphic Mappings[M], Kluwer Academic, The Netherlands, 2003.
- [17] Ueda, H. Unicity theorems for meromorphic or entire functions II. Kodai Math. J., 1983, 3:26-36
- [18] Brosch, G. Eindeutigkeitsätze für meromorphe funktions. Thesis, Technical University of Aschen. 1989.
- [19] Gundersen, G.G. Meromorphic functions that share two finite values with their derivatives, Pacific J. Math. 1983, 105:299-309.
- [20] Frank, G. Ohlenroth, W. Meromorphe Funktionen, die mit einer ihrer Ableitungen Werte teilen, Complex Variables, 1986, 6:23-37.



**Thanks to Xinli Wang,  
lead me on the path of research.**

**Thanks to Matthew Parry,  
the supervisor gave chance to start  
my PhD study.**

A faint, light blue background image of a Gothic cathedral tower, likely St. Mark's Basilica in Venice, is visible on the left side of the slide. The tower features intricate carvings and two large circular windows.

**Thank You!**