

# APPC

## TP 6 SVD & NNMF

Thomas ROBERT

### SVD

First, let's try to factorize  $X$  matrix using SVD method.

We get a  $K$  rank factorization, however, how to interpret negative values ?

```
1 % compute 16 first SVD
2 K = 16;
3 [U, S, V] = svds(X, K);
4
5 % estimate
6 Xhat = U*S*V';
7
8 % plot
9 i = 1;
10 for k = randperm(n, 8)
11     if (i <= 4) subplot(4,4,i);
12     else subplot(4,4,4 + i); end
13     imshow(reshape(X(k,:), 50, 50), [])
14     title(['Original n°' int2str(k)])
15     if (i <= 4) subplot(4,4,4 + i);
16     else subplot(4,4,8 + i); end
17     imshow(reshape(Xhat(k,:), 50, 50), [])
18     title(['Estimé n°' int2str(k)])
19     i = i + 1;
20 end
21
22 figure
23 for i = 1:K
24     subplot(4,ceil(K/4),i)
25     imshow(reshape(V(:,i), 50, 50), [])
26     title(['Composante V_{' int2str(i) '}'])
27 end
```




Figure 1 : First components of the SVD.



Figure 2 : Results of the estimates.

## NNMF

To solve this problem of negative values, let's use Non-negative Matrix Factorization of the same data.

We will use proximal gradient method or this.

Because of this constrain of positiveness, we can no longer find components with values that cancels each other. Therefore, the new components contains more null values on some zones of the picture, so that components add themselves and represent differents parts of the face.

```
1 % Params
2 K = 16;
3 eps = 1e-1;
4 kmax = 1000;
5 rhoU = 1e-5;
6 rhoV = 1e-5;
7
8 % Init
9 U = zeros(n,K);
10 V = rand(p,K);
11 J = Inf;
12 Js = [];
13
14 % Iterate
15 k = 1;
16 while (J > eps && k < kmax)
17     VOld = V;
18
19     for i = 1:n
20         U(i,:) = max(0, U(i,:) - rhoU * V'*(V*U(i,:) - X(i,:)));
21     end
22
23     for j = 1:p
24         V(j,:) = (V(j,:) - rhoV * (U'*(U*V(j,:) - X(:,j))))';
25     end
26
27     k = k + 1;
28     J = norm(VOld - V);
29     Js(end+1) = J;
30     %
31 end
32
33 % estimate
34 Xhat = U*V';
35
36 % plot
37 i = 1;
38 for k = randperm(n, 5)
39     subplot(2,5,i)
40     imshow(reshape(X(k,:), 50, 50), [])
41     title(['Original n°', int2str(k)])
42     subplot(2,5,5+i)
43     imshow(reshape(Xhat(k,:), 50, 50), [])
44     title(['Estimé n°', int2str(k) ' (K = ', int2str(K) ')'])
45     i = i + 1;
46 end
47
48 figure
49 for i = 1:K
50     subplot(4, ceil(K/4), i)
51     imshow(reshape(V(:,i), 50, 50), [])
52     title(['Composante V_', int2str(i) ])
53 end
54
55 figure
56 plot(Js)
57 title('Evolution of V matrix (norm of the variation)');
58 xlabel('Iteration')
59 ylabel('Variation norm')
```




Figure 3 :  $V$  matrix of the NNMF.



Figure 4 : Results of the estimate.




Figure 5 : Evolution of the  $V$  matrix.