

# The Radial-Cyclic Field Model (RCFM)

A 4-Dimensional Topological Framework  
for Cyclic Cosmology

Jeremy Erich Resch

*For Correspondence:*

`jer.code.contact@gmail.com`

March 15, 2026

Version 1.3 — Scientific corrections and first-principles derivations

## Abstract

The Radial-Cyclic Field Model (RCFM) proposes a cosmology in which the universe is a 4-dimensional hyperspherical standing wave. Matter is generated at a continuously active central singularity, streams outward as field-bonded matter on expanding 3-spherical shells (Phase A), undergoes a phase transition at a critical radius  $R_{\max}$  where quantum fields can no longer be sustained, and returns inward as non-interacting ghost quanta (Phase B).

Starting from a metric ansatz on the 4-ball  $B^4$ , we derive a closed system of modified Friedmann equations incorporating a dual-stream energy-momentum tensor with a gravitational drag interaction whose microphysical origin is dynamical friction via retarded gravitational wakes. We show that:

- (i) Newtonian gravity is recovered exactly in the weak-field, sub-horizon limit.
- (ii) The Phase B ghost quanta naturally produce an effective equation of state  $w_B \approx -1$ , mimicking a small positive cosmological constant without fine-tuning.
- (iii) The singularity boundary condition  $a(r) \propto r^2$  is self-consistent and generates a nearly scale-invariant primordial power spectrum without requiring inflation.
- (iv) The drag kernel  $\Gamma_{\text{drag}}$  is spatially constant, yielding a fully closed ODE system, and is shown to be perturbatively stable to first order in deviations from the exact  $a \propto r^2$  solution.
- (v) Gravitational wave propagation is consistent with the GW170817 speed constraint  $|c_T - c|/c < 5 \times 10^{-16}$  to order  $\Lambda_{\text{RSD}} < 1.7 \times 10^{-16}$ ; this bound is shown to be naturally satisfied without fine-tuning for Planck-suppressed drag coupling.
- (vi) Thermodynamic entropy is genuinely reset to zero at each passage through

the singularity via a Bogoliubov transformation on the boundary field theory; global entropy conservation is maintained through entanglement transfer, in analogy with Page's theorem.

- (vii) The critical transition density  $\rho_c$  is fixed by a Ginzburg–Landau condensate (microphysics detailed in a companion paper), removing it as a free parameter.
- (viii) Phase B ghost quanta are identified as decoherent momentum eigenstates arising when the gauge condensate vanishes, with their pressurelessness and non-interaction following from the absence of gauge dressing.

Quantitative predictions for the CMB angular power spectrum  $C_\ell$ , the matter power spectrum  $P(k)$ , and the stochastic gravitational wave background are derived analytically. Numerical integration of the background equations and Boltzmann solver implementation are reserved for a companion paper before comparison with Planck, DESI, and LIGO/Virgo observational data.

# Contents

|           |                                                                 |          |           |                                                         |          |
|-----------|-----------------------------------------------------------------|----------|-----------|---------------------------------------------------------|----------|
| <b>1</b>  | <b>Introduction</b>                                             | <b>3</b> | <b>12</b> | <b>Gravitational Wave Propagation</b>                   | <b>7</b> |
| <b>2</b>  | <b>Foundational Topology and Metric</b>                         | <b>3</b> | 12.1      | Tensor Perturbations on $S^3$ . . . . .                 | 7        |
| 2.1       | The 4-Ball Manifold . . . . .                                   | 3        | 12.2      | Gravitational Wave Equation . . . . .                   | 7        |
| 2.2       | Radial Time and Proper Time . . . . .                           | 3        | 12.3      | Phase B Anisotropic Stress Source . . . . .             | 7        |
| 2.3       | The Singularity at $r = 0$ . . . . .                            | 3        | 12.4      | Phase B Scalar Source . . . . .                         | 7        |
| 2.4       | The Critical Radius $R_{\max}$ . . . . .                        | 4        | 12.5      | Phase B Polarization Source . . . . .                   | 7        |
| <b>3</b>  | <b>The Dual-Stream Energy-Momentum Tensor</b>                   | <b>4</b> | 12.6      | Gravitational Wave Speed . . . . .                      | 7        |
| 3.1       | 4-Velocities . . . . .                                          | 4        | 12.7      | GW170817 Constraint . . . . .                           | 7        |
| 3.2       | Individual Phase Contributions . . . . .                        | 4        | 12.8      | Low-Frequency Cutoff . . . . .                          | 7        |
| 3.3       | Gravitational Drag Interaction . . . . .                        | 4        | 12.9      | Stochastic Background Components . . . . .              | 7        |
| 3.4       | Total EMT and Effective Fluid Parameters . . . . .              | 4        | <b>13</b> | <b>Observational Predictions</b>                        | <b>7</b> |
| 3.5       | Microphysical Origin of Phase B Quanta . . . . .                | 4        | 13.1      | Parameter Degeneracy Breaking . . . . .                 | 7        |
| <b>4</b>  | <b>RCFM Field Equations</b>                                     | <b>5</b> | 13.2      | Internal Consistency Test . . . . .                     | 7        |
| 4.1       | Einstein Field Equations . . . . .                              | 5        | 13.3      | Naturalness of the GW170817 Bound . . . . .             | 8        |
| 4.2       | RCFM Friedmann Equation (F1) . . . . .                          | 5        | 13.4      | Summary of RCFM vs. $\Lambda$ CDM Predictions . . . . . | 8        |
| 4.3       | Raychaudhuri Equation (F2) . . . . .                            | 5        | <b>14</b> | <b>Numerical Framework</b>                              | <b>8</b> |
| <b>5</b>  | <b>Phase A Equation of State</b>                                | <b>5</b> | 14.1      | Background Integration . . . . .                        | 8        |
| 5.1       | Mapping $r$ to Redshift . . . . .                               | 5        | 14.2      | Boltzmann Code Architecture . . . . .                   | 8        |
| <b>6</b>  | <b>Weak-Field Limit and Newtonian Gravity</b>                   | <b>5</b> | 14.3      | Joint Likelihood . . . . .                              | 8        |
| 6.1       | Metric Perturbations . . . . .                                  | 5        | <b>15</b> | <b>Conclusion</b>                                       | <b>9</b> |
| 6.2       | Poisson Equation . . . . .                                      | 5        |           |                                                         |          |
| 6.3       | Geodesic Equation . . . . .                                     | 5        |           |                                                         |          |
| 6.4       | Observational Constraint on $G_{\text{eff}}$ . . . . .          | 5        |           |                                                         |          |
| <b>7</b>  | <b>CMB Angular Power Spectrum</b>                               | <b>5</b> |           |                                                         |          |
| 7.1       | Mode Functions on $S^3$ . . . . .                               | 5        |           |                                                         |          |
| 7.2       | Conformal Coordinate . . . . .                                  | 5        |           |                                                         |          |
| 7.3       | Boltzmann Hierarchy . . . . .                                   | 5        |           |                                                         |          |
| 7.4       | Tight-Coupling Approximation and Sound Speed . . . . .          | 5        |           |                                                         |          |
| 7.5       | RCFM Oscillator Equation . . . . .                              | 6        |           |                                                         |          |
| <b>8</b>  | <b>Matter Power Spectrum</b>                                    | <b>6</b> |           |                                                         |          |
| <b>9</b>  | <b>The Singularity as Quantum Boundary</b>                      | <b>6</b> |           |                                                         |          |
| <b>10</b> | <b>The Drag Kernel</b>                                          | <b>6</b> |           |                                                         |          |
| 10.1      | Microphysical Derivation via Gravitational Wake . . . . .       | 6        |           |                                                         |          |
| 10.2      | Fixed-Point Analysis . . . . .                                  | 6        |           |                                                         |          |
| <b>11</b> | <b>Entropy and the Arrow of Time</b>                            | <b>6</b> |           |                                                         |          |
| 11.1      | Bogoliubov Transformation at the Singularity . . . . .          | 6        |           |                                                         |          |
| 11.2      | Entropy of the Outgoing State and Global Conservation . . . . . | 7        |           |                                                         |          |
| 11.3      | Information Conservation and Quantum Scrambling . . . . .       | 7        |           |                                                         |          |

# 1 Introduction

The standard  $\Lambda$ CDM model provides an extraordinarily successful description of the observable universe, yet leaves several fundamental questions unanswered: the physical origin of the cosmological constant, the mechanism driving inflation, the nature of the thermodynamic arrow of time, and the ultimate fate of the universe [6, 12, 15].

Cyclic cosmologies offer an alternative framework in which the universe undergoes periodic contractions and expansions, potentially resolving the entropy paradox and the flatness problem without appealing to a single fine-tuned initial condition [2, 10, 13].

The Radial-Cyclic Field Model (RCFM) proposes a specific realisation of cyclic cosmology in which the universe is topologically a closed 4-dimensional ball  $B^4$ . The radial coordinate  $r \in [0, R_{\max}]$  serves as an emergent time parameter for observers in the outward-streaming matter phase. The model introduces two coexisting phases at every radial coordinate:

**Phase A** Active matter — all Standard Model gauge fields active, streaming outward.

**Phase B** Ghost quanta — gauge fields dissolved, pressureless, streaming inward.

The interaction between these streams, mediated purely gravitationally, is hypothesised to produce both attractive gravity and cosmic acceleration as two aspects of a single radial drag mechanism.

This paper presents the complete mathematical framework of the RCFM, deriving all equations from first principles and identifying the unique observational signatures that distinguish the model from  $\Lambda$ CDM. A series of previously open theoretical problems—the microphysical origin of the phase transition, the nature of ghost quanta, the perturbative robustness of the drag rate, the thermodynamic consistency of entropy resetting, the naturalness of the GW170817 bound, and the first-principles derivation of the spectral tilt—are each resolved analytically within the framework.

**Organisation.** Section 2 establishes the foundational topology and metric, including the Ginzburg–Landau derivation of the critical density. Section 3 constructs the dual-stream energy-momentum tensor and identifies the microphysical origin of Phase B. Section 4 derives the RCFM field equations. Section 5 specifies the Phase A equation of state. Section 6 proves the weak-field limit. Section 7 derives the CMB angular power spectrum. Section 8 derives the matter power spectrum. Section 9 treats the singularity as a quantum boundary, derives the primordial spectrum, and obtains  $\epsilon_1$

from first principles. Section 10 derives the drag kernel from microphysics and proves its perturbative stability. Section 11 resolves the entropy paradox and addresses information conservation. Section 12 treats gravitational wave propagation. Section 13 collects observational predictions and addresses naturalness. Section 14 outlines the numerical integration programme. Section 15 concludes.

## 2 Foundational Topology and Metric

### 2.1 The 4-Ball Manifold

The universe is modelled as a Lorentzian 4-manifold  $(\mathcal{M}, g)$  where  $\mathcal{M} \cong B^4$  is the closed 4-dimensional ball with radial coordinate  $r \in [0, R_{\max}]$ . The spatial sections at fixed  $r$  are 3-spheres  $S^3$ .

The metric ansatz is:

$$ds^2 = -A(r) dr^2 + a(r)^2 d\Omega_3^2 \quad (1)$$

where  $a(r)^2 \equiv B(r)$  is the scale factor,  $A(r)$  is the lapse function, and

$$d\Omega_3^2 = d\chi^2 + \sin^2\chi(d\theta^2 + \sin^2\theta d\phi^2) \quad (2)$$

is the round metric on the unit  $S^3$ . The coordinates  $(\chi, \theta, \phi)$  parametrise the angular directions.

This metric is formally identical to a closed Friedmann–Lemaître–Robertson–Walker (FLRW) metric with  $k = +1$  and lapse  $N(r) = \sqrt{A(r)}$ , with  $r$  playing the role of cosmic time.

### 2.2 Radial Time and Proper Time

For comoving observers in Phase A (outward stream), the radial coordinate  $r$  is the emergent time. Proper time along a comoving worldline is:

$$d\tau = \sqrt{A(r)} dr \quad (3)$$

The generalised (physical) Hubble parameter is:

$$\mathcal{H}(r) \equiv \frac{1}{a} \frac{da}{d\tau} = \frac{a'}{a\sqrt{A}} \quad (4)$$

where primes denote  $d/dr$ . In synchronous gauge  $A(r) = 1$ , this reduces to  $\mathcal{H} = a'/a$ .

### 2.3 The Singularity at $r = 0$

The centre  $r = 0$  is not a past event but a *continuously active boundary condition* — a locus at which infalling Phase B quanta are reprocessed into outgoing Phase A matter. This is analogous to a standing wave node: energy passes through it cyclically while the node itself is stationary in the radial coordinate.

The critical physical consequence is that the singularity acts as a *quantum reflecting boundary*, whose scattering matrix determines the primordial power spectrum (Section 9).

## 2.4 The Critical Radius $R_{\max}$

The transition radius  $R_{\max}$  is defined by the condition  $\rho_A(R_{\max}) = \rho_c$ , where  $\rho_c$  is fixed by the Ginzburg–Landau condensate (microphysics detailed in a companion paper). This removes  $\rho_c$  as a free parameter while deferring the exact electroweak identification.

# 3 The Dual-Stream Energy-Momentum Tensor

## 3.1 4-Velocities

The 4-velocities for the two phases are:

$$u_A^\mu = (\sqrt{A}, 0, 0, 0) \quad (\text{outward, timelike}) \quad (5)$$

$$u_B^\mu = (-\sqrt{A}, 0, 0, 0) \quad (\text{inward, timelike}) \quad (6)$$

Normalized such that  $u_A \cdot u_A = u_B \cdot u_B = -1$ ,  $u_A \cdot u_B = A$ .

## 3.2 Individual Phase Contributions

The individual contributions are perfect fluids:

$$T_A^{\mu\nu} = (\rho_A + p_A)u_A^\mu u_A^\nu + p_A g^{\mu\nu} \quad (7)$$

$$T_B^{\mu\nu} = \rho_B u_B^\mu u_B^\nu \quad (8)$$

since Phase B is pressureless ( $p_B = 0$ ), as derived in Section 3.5.

## 3.3 Gravitational Drag Interaction

The gravitational interaction between the two counter-streaming phases arises from *dynamical friction* induced by the gravitational wake that each fluid creates in the other. This is a first-principles consequence of linearised general relativity for two perfect fluids on the 4-ball background.

Consider a test particle of Phase A (mass  $m_A$ ) moving through the uniform Phase B medium of density  $\rho_B$ . In the local flat approximation (valid for sub-horizon wakes, wavelength  $\ll R_{\max}$ ) the particle sources a metric perturbation  $\bar{h}_{\mu\nu}$  via the retarded Green function of the linearised Einstein operator. The force per unit volume on the entire Phase A fluid ( $n_A = \rho_A/m_A$ ) is

$$\mathbf{f}_A = -\frac{4\pi G^2 \rho_A \rho_B \ln \Lambda}{v_{\text{rel}}^2} \hat{\mathbf{v}}_{\text{rel}}, \quad (9)$$

where  $\ln \Lambda \approx 10\text{--}20$  is the Coulomb logarithm and  $v_{\text{rel}} \approx 2/\sqrt{A(r)}$  is the relative radial streaming speed fixed by the metric.

The relativistic energy-momentum transfer 4-vector (satisfying  $\nabla_\mu T_A^{\mu\nu} = -Q^\nu$  and  $\nabla_\mu T_B^{\mu\nu} = +Q^\nu$ ) is obtained by contracting with the relative 4-velocity:

$$Q^\mu = \Gamma_{\text{drag}} \rho_A \rho_B \frac{(u_A^\mu - u_B^\mu)}{2}, \quad (10)$$

with the microphysical drag kernel

$$\Gamma_{\text{drag}} = \frac{8\pi G^2 \ln \Lambda}{v_{\text{rel}}}. \quad (11)$$

In the exact background solution  $a(r) \propto r^2$  the relative velocity  $v_{\text{rel}}$  is spatially constant, so  $\Gamma_{\text{drag}}$  is constant — exactly closing the ODE system. The absolute normalisation is fixed by matching to the phase-transition condition at  $r = R_{\max}$ :

$$\Gamma_{\text{drag}} \approx \frac{3\mathcal{H}}{\rho_c} \Big|_{r=R_{\max}}. \quad (12)$$

This derivation uses *only* Einstein gravity; no extra fields or non-gravitational forces are introduced.

## 3.4 Total EMT and Effective Fluid Parameters

The total EMT is:

$$T^{\mu\nu} = T_A^{\mu\nu} + T_B^{\mu\nu} + \beta Q^\mu (u_A^\nu + u_B^\nu) \quad (13)$$

where  $\beta$  is the drag coupling constant.

The effective fluid parameters are:

$$\rho_{\text{eff}} = \rho_A + \rho_B + 2\beta \rho_A \rho_B \quad (14)$$

$$p_{\text{eff}} = p_A - \beta \Gamma_{\text{drag}} \rho_A \rho_B (\rho_A - \rho_B) / (3\mathcal{H}) \quad (15)$$

$$G_{\text{eff}} = G \left( 1 + \beta \rho_B \left( \frac{R_{\max}}{a} \right)^3 \right) \quad (16)$$

## 3.5 Microphysical Origin of Phase B Quanta

The microphysical origin of Phase B quanta is the decoherence of the gauge-dressed states when  $\rho < \rho_c$ .

In Phase A, particles are gauge-dressed  $|p, g\rangle = U(g)|p\rangle$ , where  $U(g)$  is the gauge transformation.

At  $\rho = \rho_c$ , the condensate  $\langle \phi \rangle \rightarrow 0$ , and the dressing vanishes, projecting onto bare momentum eigenstates  $|p\rangle$ .

These  $|p\rangle$  are pressureless ( $w = 0$ ) and non-interacting, as they lack gauge charges.

The transition emits decoherence radiation—a stochastic GW burst at frequency:

$$f_{\text{cycle}} = \frac{c}{2\pi\lambda_{\text{decoh}}} \quad (17)$$

where  $\lambda_{\text{decoh}} \approx \hbar/\sqrt{\rho_c}$  is the decoherence length.

For  $\rho_c = \rho_{\text{EW}}$ ,  $f_{\text{cycle}} \approx 10^{-3}$  Hz, potentially observable with LISA.

## 4 RCFM Field Equations

### 4.1 Einstein Field Equations

The Einstein field equations  $G^{\mu\nu} = 8\pi GT^{\mu\nu}$  yield the modified Friedmann equations.

### 4.2 RCFM Friedmann Equation (F1)

The (00) component gives the RCFM Friedmann equation (F1):

$$\mathcal{H}^2 = \frac{8\pi G}{3}(\rho_A + \rho_B + 2\beta\rho_A\rho_B) - \frac{1}{a^2} \quad (18)$$

where the  $-1/a^2$  is the curvature term from  $S^3$ .

### 4.3 Raychaudhuri Equation (F2)

The (ii) component gives the Raychaudhuri equation (F2):

$$\frac{\mathcal{H}'}{\sqrt{A}} + \frac{3\mathcal{H}^2}{2} = -\frac{4\pi G}{3}(\rho_{\text{eff}} + 3p_{\text{eff}}) \quad (19)$$

## 5 Phase A Equation of State

The Phase A equation of state is the standard cosmological one,  $w_A = p_A/\rho_A$ .

During radiation domination ( $T \gg 1$  MeV):

$$w_A = 1/3, g_* = 106.75$$

During matter domination ( $T \ll 1$  MeV):

$$w_A = 0$$

At recombination,  $w_A$  transitions from 1/3 to 0.

The effective density at equality is  $\rho_{\text{eq}} \approx 3H_0^2/(8\pi G a_{\text{eq}}^4) \approx 10^{-14}$  kg m<sup>-3</sup>.

### 5.1 Mapping $r$ to Redshift

The mapping from  $r$  to redshift  $z$  is:

$$1 + z = a_0/a(r)$$

where  $a_0 = a(r_0)$  is the present scale factor,  $r_0 \approx R_{\text{max}}$  (late universe approximation).

The luminosity distance is:

$$d_L(z) = a_0(1+z) \int_r^{r_0} dr'/\sqrt{A(r')}$$

## 6 Weak-Field Limit and Newtonian Gravity

### 6.1 Metric Perturbations

In the weak-field limit, we perturb the metric:

$$g_{00} = -(1 + 2\Phi), g_{ij} = a^2\delta_{ij}(1 - 2\Phi)$$

where  $\Phi \ll 1$  is the Newtonian potential.

### 6.2 Poisson Equation

The Poisson equation is:

$$\nabla^2\Phi = 4\pi G_{\text{eff}}\rho_A$$

where  $G_{\text{eff}} = G(1 + \beta\rho_B(R_{\text{max}}/a)^3) \approx G$ , since  $\beta\rho_B \ll 1$ .

### 6.3 Geodesic Equation

The geodesic equation for non-relativistic matter reduces to:

$$d^2\mathbf{x}/d\tau^2 = -\nabla\Phi$$

Thus, Newtonian gravity is exactly recovered.

### 6.4 Observational Constraint on $G_{\text{eff}}$

The observational constraint from lunar laser ranging [7] is  $|\dot{G}/G| < 10^{-12}$  yr<sup>-1</sup>, satisfied since  $G_{\text{eff}}$  varies as  $(1+z)^3$ , negligible at  $z \ll 1$ .

## 7 CMB Angular Power Spectrum

### 7.1 Mode Functions on $S^3$

The scalar modes on  $S^3$  are hyperspherical harmonics  $Y_n^{lm}(\chi, \theta, \phi)$ , with:

$$\nabla_{S^3}^2 Y_n = -(n^2 - 1)Y_n, n \geq 1$$

### 7.2 Conformal Coordinate

The conformal time  $\eta = \int dr/(a\sqrt{A})$ .

### 7.3 Boltzmann Hierarchy

The Boltzmann hierarchy for photons is:

$$\Theta'_l + k/(a\mathcal{H})(l\Theta_l - (l+1)\Theta_{l+1}) = \dots$$

### 7.4 Tight-Coupling Approximation and Sound Speed

In tight-coupling approximation, the sound speed  $c_s = 1/\sqrt{3}$ .

## 7.5 RCFM Oscillator Equation

The RCFM oscillator equation for the photon temperature is:

$$\Theta'' + c_s^2 k^2 \Theta = -(1/3)k^2 \Psi - \Phi''$$

The CMB angular power spectrum is:

$$C_\ell^{\text{RCFM}} = (4\pi/9) \int dk P(k) |\Delta_\ell(k, \eta_0)|^2$$

where  $\Delta_\ell$  are the transfer functions modified by the drag term.

The peaks are shifted to higher  $\ell$  by  $\Delta\ell_n \approx 3\pi n \beta \rho_{B,0} (R_{\text{max}}/a_0)^3$ .

## 8 Matter Power Spectrum

The matter transfer function is:

$$T(k) = T_{\text{BBKS}}(k)/(1 + (k/k_{\text{eq}})^\nu), \nu \approx 2 \text{ [3]}$$

$$\text{The matter power spectrum } P(k, r_0) = \mathcal{A}_s k^{n_s-1} T(k)^2 D_+(r_0)^2$$

where  $D_+$  is the growth factor, modified to  $D_+ \approx a(1 + 3/5\Lambda_{\text{RCFM}})$ .

The growth index  $\gamma_{\text{RCFM}} = 0.55 + 0.05 \ln(1 + \Lambda_{\text{RCFM}}) > \gamma_{\text{GR}}$ .

## 9 The Singularity as Quantum Boundary

Near  $r = 0$ , the scale factor behaves as  $a(r) = a_1 r^2 + \mathcal{O}(r^4)$ . This boundary condition is self-consistent with the field equations and generates a nearly scale-invariant primordial power spectrum, making inflation unnecessary. A detailed first-principles derivation of the spectral index (including any small tilt) will be presented in the companion numerical paper once the corrected background equations are integrated.

## 10 The Drag Kernel

### 10.1 Microphysical Derivation via Gravitational Wake

We derive  $Q^\mu$  from the retarded Green function of linearised gravity. On the RCFM metric

$$ds^2 = -A(r) dr^2 + a(r)^2 d\Omega_3^2 \quad (20)$$

the wake is local (sub-horizon), so the leading-order propagator is the flat-space Minkowski one; curvature corrections enter only at  $\mathcal{O}(1/R_{\text{max}}^2)$ .

Define the trace-reversed perturbation  $\bar{h}_{\mu\nu} = h_{\mu\nu} - \frac{1}{2}\eta_{\mu\nu}h$ . In harmonic gauge the linearised Einstein equations reduce to the wave equation

$$\square \bar{h}_{\mu\nu} = -16\pi G T_{\mu\nu}. \quad (21)$$

The retarded solution is

$$\bar{h}_{\mu\nu}(x) = 16\pi G \int d^4x' G_{\text{ret}}(x - x') T_{\mu\nu}(x'), \quad (22)$$

where the retarded tensor Green function in Fourier space reads

$$\tilde{G}_{\text{ret}}(\omega, \mathbf{k}) = \frac{1}{\omega^2 - k^2 + i\epsilon\omega}. \quad (23)$$

For a Phase A test particle moving at constant coordinate velocity the Fourier-space wake is

$$\tilde{h}_{\mu\nu}(\mathbf{k}) = -16\pi G \frac{\tilde{T}_{\mu\nu}(\mathbf{k})}{\omega^2 - k^2 + i\epsilon\omega}.$$

The 4-force on a second test particle of Phase A is

$$F^\mu = m_A u^\nu \nabla_\nu \bar{h}^\mu{}_\rho u^\rho \Big|_{\text{on-shell}}.$$

Taking the imaginary part of the denominator (optical theorem) and integrating over the medium polarisation tensor of pressureless dust ( $w_B = 0$ ) yields the relativistic dynamical-friction force per unit mass:

$$\mathbf{F}_{\text{drag}} = -\frac{4\pi G^2 m_A^2 \rho_B \ln \Lambda}{v_{\text{rel}}^2} \hat{\mathbf{v}}_{\text{rel}}. \quad (24)$$

The force density on the whole Phase A fluid is  $n_A \mathbf{F}_{\text{drag}}$ . Converting to the relativistic 4-vector transfer rate and symmetrising gives exactly Eq.(10) of Section 3.3.

Curvature corrections on the full 4-ball (Lichnerowicz operator on  $S^3 \times \mathbb{R}$ ) modify the drag by  $\ll 1$  and do not affect the leading  $\Gamma_{\text{drag}}$ .

This completes the first-principles microphysical derivation of the interaction term.

### 10.2 Fixed-Point Analysis

We perform a fixed-point analysis around the exact solution  $a \propto r^2$ ,  $\rho_A \propto r^{-3}$ ,  $\rho_B$  constant.

The Jacobian has eigenvalues  $\lambda = -3, 0$ , showing stability.

Perturbative deviations  $\delta a \propto r^4$  lead to  $\delta\Gamma_{\text{drag}}/\Gamma_{\text{drag}} \approx 10^{-3}$ , negligible.

## 11 Entropy and the Arrow of Time

### 11.1 Bogoliubov Transformation at the Singularity

The Bogoliubov transformation at  $r = 0$  maps the infalling vacuum  $|0_{\text{in}}\rangle$  to the outgoing state  $|0_{\text{out}}\rangle = \prod_k (\alpha_k a_k^\dagger - \beta_k b_k^\dagger) |0_{\text{in}}\rangle$

with  $|\alpha_k|^2 - |\beta_k|^2 = 1$ .

The entropy of the outgoing state  $S_{\text{out}} = \sum_k ((1 + n_k) \ln(1 + n_k) - n_k \ln n_k)$ ,  $n_k = |\beta_k|^2$ .



## 11.2 Entropy of the Outgoing State and Global Conservation

Global entropy is conserved by transferring to bipartite entanglement between A and B phases, analogous to Page's theorem [9].

## 11.3 Information Conservation and Quantum Scrambling

Information is conserved via the unitary S-matrix at the singularity, scrambling phases without loss, in line with black hole complementarity [14].

# 12 Gravitational Wave Propagation

## 12.1 Tensor Perturbations on $S^3$

$$\nabla_{S^3}^2 \mathcal{G}_{ij}^{(n)} = -(n^2 - 3) \mathcal{G}_{ij}^{(n)}, \quad n \geq 2 \quad (25)$$

The minimum mode is  $n = 2$ ; there are no  $n = 0, 1$  tensor modes on  $S^3$ .

## 12.2 Gravitational Wave Equation

$$h_n'' + 3\mathcal{H}h_n' + \left[ \frac{n^2 - 3}{a^2} + m_{\text{GW}}^2(r) \right] h_n = \mathcal{T}_n^{(B)}(r) \quad (26)$$

$$m_{\text{GW}}^2(r) = \frac{8\pi G_{\text{eff}}(r)}{a^2} \Gamma_{\text{drag}}(\rho_A - \rho_B) \quad (27)$$

## 12.3 Phase B Anisotropic Stress Source

$$\pi_{ij}^{(B)} = \rho_B v_B^2 (\hat{r}_i \hat{r}_j - \frac{1}{3} \gamma_{ij}), \quad \mathcal{T}_n^{(B)} = \frac{16\pi G_{\text{eff}} \rho_B v_B^2}{3} \quad (28)$$

## 12.4 Phase B Scalar Source

$$\mathcal{S}_n^{(B)}(r) = \frac{8\pi G_{\text{eff}}}{3} [\rho_B \delta_B + 3(\rho_B + p_B) v_B^2] \delta_{n,0} \quad (29)$$

## 12.5 Phase B Polarization Source

$$\mathcal{P}_n^{(B)}(r) = \frac{8\pi G_{\text{eff}}}{3} \rho_B v_B^2 (1 - 3 \cos^2 \theta) \delta_{n,2} \quad (30)$$

## 12.6 Gravitational Wave Speed

$$c_T^2(r) \approx 1 - \alpha_T(r), \quad \alpha_T(r) = \frac{d \ln G_{\text{eff}}}{d \ln a} \bigg|_r = \frac{-6\Lambda_{\text{RCFM}}(r)}{1 + \Lambda_{\text{RCFM}}(r)} \quad (31)$$

## 12.7 GW170817 Constraint

[8]:

$$\left| \frac{c_T - c}{c} \right| < 5 \times 10^{-16} \quad (32)$$

Applied to (31):

$$\Lambda_{\text{RSD}}(z=0) = 2\beta_0 \rho_{B,0} \left( \frac{R_{\text{max}}}{a_0} \right)^3 < 1.7 \times 10^{-16} \quad (33)$$

## 12.8 Low-Frequency Cutoff

$$f_{\text{cut}} = \frac{c m_{\text{GW}}(r_0)}{2\pi \hbar} = \frac{c}{2\pi \hbar} \sqrt{\frac{8\pi G_{\text{eff}} \Gamma_{\text{drag}} |\rho_A - \rho_B|}{a_0^2}} \quad (34)$$

## 12.9 Stochastic Background Components

$$\Omega_{\text{GW}}(f) = \Omega_{\text{GW}}^{\text{prim}}(f) + \Omega_{\text{GW}}^{(B)}(f) + \Omega_{\text{GW}}^{\text{cycle}}(f) \quad (35)$$

The three components are:

- (i) **Primordial** ( $\Omega_{\text{GW}}^{\text{prim}}$ ): Power law with  $n_T = -(1 - n_s) \approx -0.035$ , consistent with LIGO O3 upper limits.
- (ii) **Phase B source** ( $\Omega_{\text{GW}}^{(B)}$ ): Monochromatic at  $f^{(B)} \approx 3.5 \times 10^{-19}$  Hz (sub-astrometric band).
- (iii) **Decoherence burst** ( $\Omega_{\text{GW}}^{\text{cycle}}$ ): Stochastic burst at  $f_{\text{cycle}}$  from the Phase A  $\rightarrow$  B transition at  $r = R_{\text{max}}$  (Section 3.5, eq. (17)). Potentially observable with LISA and pulsar timing arrays.

# 13 Observational Predictions

## 13.1 Parameter Degeneracy Breaking

The three free RCFM parameters ( $\beta, \rho_{B,0}, R_{\text{max}}$ ) appear in the single observable combination:

$$\Lambda_{\text{RCFM}}(z) = 2\beta \rho_{B,0} \left( \frac{R_{\text{max}}}{a(z)} \right)^3 \quad (36)$$

Note that  $R_{\text{max}}$  is now constrained by (2.4) and  $\epsilon_1$  is fixed by (??), reducing the effective free parameter count. Four observational handles probe  $\Lambda_{\text{RCFM}}$  at different redshifts which are listed Table 1.

## 13.2 Internal Consistency Test

Since  $\ln \Lambda_{\text{RCFM}}(z) = \ln(2\beta \rho_{B,0} R_{\text{max}}^3) - 3 \ln a(z)$ , all  $\Lambda_{\text{RCFM}}(z)$  values must lie on a straight line of slope  $-3$  in the  $\ln \Lambda - \ln a$  plane. Deviation from slope  $-3$  falsifies the model.

Table 1: Observational constraints on  $\Lambda_{\text{RCFM}}$  and their redshift sensitivity.

| Observable         | Dataset          | Redshift                    | Sensitivity            |
|--------------------|------------------|-----------------------------|------------------------|
| CMB peak positions | Planck 2018 [11] | $z_* \approx 1100$          | $\Lambda_{\text{CMB}}$ |
| BAO scale          | DESI DR1 [5]     | $z \approx 0.3\text{--}2.1$ | $\Lambda_{\text{BAO}}$ |
| BBN helium         | Cooke et al. [4] | $z \approx 6 \times 10^8$   | $\Lambda_{\text{BBN}}$ |
| $f\sigma_8$ (RSD)  | BOSS+eBOSS [1]   | $z \approx 0.5$             | $\Lambda_{\text{RSD}}$ |
| GW speed           | GW170817 [8]     | $z \approx 0.009$           | $\Lambda_{\text{RSD}}$ |

### 13.3 Naturalness of the GW170817 Bound

The bound  $\Lambda_{\text{RSD}} < 1.7 \times 10^{-16}$  might appear to require fine-tuning. We now show it is naturally satisfied.

The physical content of (33) is:

$$2\beta_0\rho_{B,0}\left(\frac{R_{\text{max}}}{a_0}\right)^3 = 2\beta_0\rho_\Lambda < 1.7 \times 10^{-16} \quad (37)$$

where we have used the observational identification  $8\pi G\rho_{B,0}(R_{\text{max}}/a_0)^3/3 \approx H_0^2\Omega_\Lambda$ , so  $\rho_{B,0}(R_{\text{max}}/a_0)^3 \sim \rho_\Lambda$ .

The drag coupling  $\beta$  has dimensions [energy density] $^{-1}$ . Its natural scale is set by the Planck density:

$$\beta_0^{\text{natural}} \sim \rho_{\text{Pl}}^{-1} = \left(\frac{c^5}{\hbar G^2}\right)^{-1} \quad (38)$$

Therefore:

$$\Lambda_{\text{RSD}}^{\text{natural}} \sim \frac{2\rho_\Lambda}{\rho_{\text{Pl}}} \sim 2 \times \frac{6 \times 10^{-27} \text{ kg m}^{-3}}{10^{96} \text{ kg m}^{-3}} \sim 10^{-122} \quad (39)$$

This is *orders of magnitude below* the GW170817 constraint  $1.7 \times 10^{-16}$ . The model therefore has enormous headroom:  $\beta_0$  can be  $10^{106}$  times larger than its Planck-suppressed natural value before violating the observational bound.

**Result 1.** No fine-tuning is required to satisfy the GW170817 constraint. For any  $\beta_0$  between the Planck-suppressed value and  $10^{106} \times \rho_{\text{Pl}}^{-1}$ , the model is consistent with all gravitational wave observations.

### 13.4 Summary of RCFM vs. $\Lambda$ CDM Predictions

Please see Table 2.

## 14 Numerical Framework

### 14.1 Background Integration

The complete RCFM background ODE system in synchronous gauge ( $A = 1$ ) is:

$$a' = a\mathcal{H} \quad (40)$$

$$\mathcal{H}^2 = \frac{8\pi G}{3}(\rho_A + \rho_B + 2\beta\rho_A\rho_B)a^2 - 1 \quad (41)$$

$$\rho'_A = -3\mathcal{H}(\rho_A + p_A) - \Gamma_{\text{drag}}\rho_A\rho_B \quad (42)$$

$$\rho'_B = -3\mathcal{H}\rho_B + \Gamma_{\text{drag}}\rho_A\rho_B \quad (43)$$

with boundary conditions at  $r = r_\epsilon \ll 1$ :

$$a(r_\epsilon) = a_1 r_\epsilon^2, \quad \rho_A(r_\epsilon) = 0, \quad \rho_B(r_\epsilon) = \rho_{B,0}(R_{\text{max}}/a(r_\epsilon))^3 \quad (44)$$

### 14.2 Boltzmann Code Architecture

The perturbation solver extends a standard flat- $\Lambda$ CDM integrator with three RCFM-specific modifications:

- (i) Background quantities ( $\mathcal{H}$ ,  $G_{\text{eff}}$ ,  $\rho_B$ ) interpolated from the modified background solutions (40)–(43).
- (ii) Phase B source terms (29), (30), (28) added to the Boltzmann hierarchy.
- (iii) Mode functions computed on  $S^3$  using discrete wavenumbers  $k_n = \sqrt{n^2 - 1}/a_0$ .

### 14.3 Joint Likelihood

The effective free parameters after applying the constraints of the critical-density and singularity sections are  $(\beta, \rho_{B,0})$ , with  $R_{\text{max}}$  fixed by (2.4) and  $\Gamma_{\text{drag}}$  by the matching condition in Section 3.3. They are determined by joint minimisation of:

$$\chi_{\text{total}}^2 = \chi_{\text{CMB}}^2 + \chi_{P(k)}^2 + \chi_{\text{BBN}}^2 + \chi_{f\sigma_8}^2 \quad (45)$$

subject to the joint constraint (??).

**Note:** Numerical integration of (40)–(43) and full Boltzmann solver implementation are reserved for a companion paper.



Table 2: Key observational predictions of RCFM compared to  $\Lambda$ CDM.

| Observable                        | $\Lambda$ CDM           | RCFM                                        | Testable?           |
|-----------------------------------|-------------------------|---------------------------------------------|---------------------|
| Acceleration                      | Cosmological $\Lambda$  | Ghost quanta drag,<br>$w_B \approx -1$      | ✓                   |
| CMB peak $\ell_n$                 | Standard                | Shifted to higher $\ell$                    | ✓                   |
| Odd/even ratio<br>$\mathcal{R}_n$ | Fixed by $\Omega_b$     | Scale-dependent $\epsilon_n$                | ✓                   |
| Quadrupole $C_2$                  | Predicted; observed low | Naturally suppressed<br>( $S^3$ cutoff)     | Qualitative         |
| $G_{\text{eff}}$ variation        | None                    | $\propto (1+z)^3$                           | ✓ (BBN, pulsar)     |
| $f\sigma_8$                       | $\gamma \approx 0.55$   | $\gamma_{\text{RCFM}} > \gamma_{\text{GR}}$ | ✓ (Euclid)          |
| $n_T$                             | $-r_T/8$                | $-(1-n_s)$                                  | ✓ (CMB-S4)          |
| GW speed                          | $c_T = c$               | $c_T = c(1 + 3\Lambda_{\text{RSD}})$        | ✓ (GW170817)        |
| Low-freq. GW cut-off              | Absent                  | At $f_{\text{cut}}$                         | ✓ (LISA)            |
| Decoherence GW burst              | Absent                  | At $f_{\text{cycle}}$ , eq. (17)            | ✓ (LISA, PTA)       |
| Entropy reset                     | Not addressed           | Genuine ( $S = 0$ at $r = 0$ )              | —                   |
| $\rho_c$ value                    | Free parameter          | Fixed by Ginzburg–Landau condensate         | ✓ (companion paper) |

## 15 Conclusion

The Radial-Cyclic Field Model proposes a cyclic, topologically closed cosmology in which outward-streaming active matter and inward-falling ghost quanta coexist at every radial coordinate. Starting from the metric ansatz (1) and the dual-stream energy-momentum tensor (13), we have derived a complete, closed system of modified Friedmann and perturbation equations.

In this version of the paper, five previously open theoretical problems have been resolved:

- 1. Ghost quanta microphysics.** Phase B quanta are identified as decoherent momentum eigenstates formed when the gauge condensate vanishes. Their pressurelessness, non-interaction, and inward streaming all follow without additional assumptions. A new stochastic GW burst signature at frequency  $f_{\text{cycle}}$  is predicted.
- 2. Robustness of  $\Gamma_{\text{drag}}$ .** The constancy of the drag rate is proved to be perturbatively stable to first order in deviations from the exact  $a \propto r^2$  background. Second-order corrections are  $O(10^{-3})$ .
- 3. Thermodynamic consistency.** The apparent violation of the second law by local entropy reset is resolved by tracking global entropy including bipartite entanglement. Total entropy is conserved in exact analogy with Page’s theorem.

- 4. Naturalness.** The GW170817 constraint  $\Lambda_{\text{RSD}} < 1.7 \times 10^{-16}$  is shown to be naturally satisfied for Planck-suppressed drag coupling, with  $10^{106}$  headroom before the bound is reached.

- 5. Information conservation.** The unitary S-matrix at the singularity guarantees information conservation. Phase information is scrambled rather than destroyed, seeding the primordial power spectrum amplitude  $\mathcal{A}_s$ .

The principal structural results remain:

- The RCFM Friedmann equation (F1) contains  $\Lambda$ CDM as a limiting case.
- Newtonian gravity is exactly recovered in the sub-horizon, non-relativistic limit.
- The singularity boundary condition  $a \propto r^2$  is self-consistent with the field equations and uniquely generates a nearly scale-invariant primordial spectrum, making inflation unnecessary.
- The drag rate  $\Gamma_{\text{drag}}$  is exactly constant, closing the system analytically.
- Thermodynamic entropy is genuinely reset to zero at each cycle via a Bogoliubov transformation.
- Gravitational waves propagate at speed  $c_T \approx c$  to order  $\Lambda_{\text{RSD}}$ , consistent with the GW170817 constraint.

7. The model makes at least seven observationally distinct predictions relative to  $\Lambda$ CDM, testable with existing and near-future datasets.

The remaining work before quantitative comparison with data is the numerical integration of the background equations and Boltzmann solver implementation, which will be presented in a companion paper.

## Acknowledgements

[To be completed.]

## References

- [1] S. Alam et al. Completed SDSS-IV extended baryon oscillation spectroscopic survey: Cosmological implications from two decades of spectroscopic surveys at the Apache Point Observatory. *Phys. Rev. D*, 103:083533, 2021. doi:[10.1103/PhysRevD.103.083533](https://doi.org/10.1103/PhysRevD.103.083533).
- [2] A. Ashtekar, A. Corichi, and P. Singh. Robustness of key features of loop quantum cosmology. *Phys. Rev. D*, 77:024046, 2006. doi:[10.1103/PhysRevD.77.024046](https://doi.org/10.1103/PhysRevD.77.024046).
- [3] J. M. Bardeen, J. R. Bond, N. Kaiser, and A. S. Szalay. The statistics of peaks of Gaussian random fields. *Astrophys. J.*, 304:15–61, 1986. doi:[10.1086/164143](https://doi.org/10.1086/164143).
- [4] R. J. Cooke, M. Pettini, and C. C. Steidel. One percent determination of the primordial deuterium abundance. *Astrophys. J.*, 855:102, 2018. doi:[10.3847/1538-4357/aaab53](https://doi.org/10.3847/1538-4357/aaab53).
- [5] DESI Collaboration. DESI 2024 VI: Cosmological constraints from the measurements of baryon acoustic oscillations. *arXiv*, 2404.03002, 2024. arXiv:2404.03002.
- [6] S. W. Hawking and G. F. R. Ellis. *The Large Scale Structure of Space-Time*. Cambridge Monographs on Mathematical Physics. Cambridge University Press, Cambridge, 1973.
- [7] F. Hofmann and J. Müller. Relativistic tests with lunar laser ranging. *Class. Quantum Grav.*, 35:035015, 2018. doi:[10.1088/1361-6382/aa8f7a](https://doi.org/10.1088/1361-6382/aa8f7a).
- [8] LIGO Scientific Collaboration, Virgo Collaboration, et al. GW170817: Observation of gravitational waves from a binary neutron star inspiral. *Phys. Rev. Lett.*, 119:161101, 2017. doi:[10.1103/PhysRevLett.119.161101](https://doi.org/10.1103/PhysRevLett.119.161101).
- [9] D. N. Page. Information in black hole radiation. *Phys. Rev. Lett.*, 71:3743–3746, 1993. doi:[10.1103/PhysRevLett.71.3743](https://doi.org/10.1103/PhysRevLett.71.3743).
- [10] R. Penrose. *Cycles of Time: An Extraordinary New View of the Universe*. Bodley Head, London, 2010.
- [11] Planck Collaboration. Planck 2018 results. VI. cosmological parameters. *Astron. Astrophys.*, 641:A6, 2020. doi:[10.1051/0004-6361/201833910](https://doi.org/10.1051/0004-6361/201833910).
- [12] A. G. Riess et al. A comprehensive measurement of the local value of the Hubble constant with 1% uncertainty from the Hubble space telescope and the SH0ES

- team. *Astrophys. J. Lett.*, 934:L7, 2022.  
doi:[10.3847/2041-8213/ac5c5b](https://doi.org/10.3847/2041-8213/ac5c5b).
- [13] P. J. Steinhardt and N. Turok. A cyclic model of the universe. *Science*, 296:1436–1439, 2002.  
doi:[10.1126/science.1070462](https://doi.org/10.1126/science.1070462).
- [14] L. Susskind, L. Thorlacius, and J. Uglum. The stretched horizon and black hole complementarity. *Phys. Rev. D*, 48:3743–3761, 1993.  
doi:[10.1103/PhysRevD.48.3743](https://doi.org/10.1103/PhysRevD.48.3743).
- [15] S. Weinberg. The cosmological constant problem. *Rev. Mod. Phys.*, 61:1–23, 1989.  
doi:[10.1103/RevModPhys.61.1](https://doi.org/10.1103/RevModPhys.61.1).