

[返回到第 2 周](#)
[X课程](#)
[上一个](#)
[下一个](#)


## Gradient Descent in Practice II - Learning Rate

**Note:** [5:20 - the x-axis label in the right graph should be  $\theta$  rather than No. of iterations]

**Debugging gradient descent.** Make a plot with *number of iterations* on the x-axis. Now plot the cost function,  $J(\theta)$  over the number of iterations of gradient descent. If  $J(\theta)$  ever increases, then you probably need to decrease  $\alpha$ .


**Automatic convergence test.** Declare convergence if  $J(\theta)$  decreases by less than  $E$  in one iteration, where  $E$  is some small value such as  $10^{-3}$ . However in practice it's difficult to choose this threshold value.

### Making sure gradient descent is working correctly.



It has been proven that if learning rate  $\alpha$  is sufficiently small, then  $J(\theta)$  will decrease on every iteration.

### Making sure gradient descent is working correctly.



- For sufficiently small  $\alpha$ ,  $J(\theta)$  should decrease on every iteration.
- But if  $\alpha$  is too small, gradient descent can be slow to converge.

To summarize:

If  $\alpha$  is too small: slow convergence.

If  $\alpha$  is too large: may not decrease on every iteration and thus may not converge.

✓ 完成

