

Part A

TIME: 15 minutes

- Look at the four texts, A–D, in the separate Text Booklet.
- For each question, 1–20, look through the texts, A–D, to find the relevant information.
- Write your answers on the spaces provided in this Question Paper.
- Answer all the questions within the 15-minute time limit.
- Your answers should be correctly spelt.

Diarrhea and Dehydration in Children: Questions

Questions 1–7

For each question, 1–7, decide which text (A, B, C or D) the information comes from. You may use any letter more than once.

In which text can you find information about

1. the classification of diarrhea according to how long the symptoms last?
2. a structured, step-by-step approach to managing a child with diarrhea?
3. altered consciousness as a sign requiring intravenous fluid resuscitation?
4. rotavirus vaccination as a strategy to prevent diarrheal illness?
5. the recommendation that feeding should not be stopped during a diarrheal illness?
6. poor sanitation and lack of breastfeeding as environmental risk factors?
7. dehydration being classified as mild, moderate, or severe based on clinical signs?

Questions 8–14

Answer each of the questions, 8–14, with a word or short phrase from one of the texts. Each answer may include words, numbers or both.

8. Diarrhea is defined as the passage of _____ or more loose or watery stools within a 24-hour period.
9. The most common viral cause of diarrhea in children globally is _____.
10. Children under _____ years of age are identified as being at greatest risk of diarrheal illness.
11. Zinc supplementation alongside ORS has been shown to reduce the _____ of acute diarrheal episodes.
12. Diarrhea lasting 14 days or more but fewer than 30 days is classified as _____.
13. Children with severe dehydration require prompt _____ fluid resuscitation.
14. Continued feeding during diarrhea helps maintain _____ and supports recovery.

Questions 15–20

Complete each of the questions, 15–20, with a word or short phrase from one of the texts. Each answer may include words, numbers or both.

15. What sign of dehydration involves the eyes appearing hollow or sunken?

16. What oral solution is used as the first-line treatment for mild to moderate dehydration?

17. What life-threatening condition can result from untreated severe dehydration?

18. What feeding practice in the first six months of life significantly reduces diarrheal morbidity?

19. What assessment is recommended for children with persistent diarrhea exceeding 14 days?

20. What should parents and caregivers be educated on before a child is discharged after a diarrheal illness?

END OF PART A

THIS QUESTION PAPER WILL BE COLLECTED