Pedo Extractions

URL: `/dental-care/kids-dentistry/pedo-extractions`

Gentle Tooth Removal for Children

Pedo extractions, also known as paediatric tooth extractions, involve removing a child's tooth when it cannot be preserved or when removal is necessary for their oral health. Although baby teeth naturally fall out, some may need to be extracted because of extensive decay, infection, dental trauma or problems with the development of permanent teeth.

At Jerush Dentofacial & Cosmetic Laser Centre, the decision to remove a child's tooth is based on its condition, the child's stage of dental development, and the potential effect on the surrounding teeth. Whenever possible, preserving a baby tooth is considered to support normal chewing and maintain space for the permanent tooth.

"C:\Jerush-dentoface-new-site-react\public\images\medical-services\dental-care\kids-dentistry-pedodontics\pedo-extractions\Gentle-pediatric-tooth-extraction.webp"

Overview:

Children sometimes need a tooth removed to prevent discomfort, manage dental infections, or make way for healthy dental development. Pedo extractions are carried out specifically for children, with attention to their comfort, safety, and individual dental needs. Depending on the situation, the dentist may recommend removing a badly damaged baby tooth, a loose tooth that is not shedding properly or a tooth affecting the eruption of permanent teeth.

"C:\Jerush-dentoface-new-site-react\public\images\medical-services\dental-care\kids-dentistry-pedodontics\pedo-extractions\overview.webp"

When Might a Child Need a Tooth Extraction?

A paediatric dentist may recommend extraction in situations such as:

- Severe tooth decay: When the tooth is too damaged to be restored predictably.

"C:\Jerush-dentoface-new-site-react\public\images\medical-services\dental-care\kids-dentistry-pedodontics\pedo-extractions\severe-tooth-decay.webp"

- Dental infection: When infection or damage makes removal the appropriate treatment.

  "C:\Jerush-dentoface-new-site-react\public\images\medical-services\dental-care\kids-dentistry-pedodontics\pedo-extractions\dental-infection.webp"

- Dental trauma: When an injury has severely damaged a baby tooth.

  "C:\Jerush-dentoface-new-site-react\public\images\medical-services\dental-care\kids-dentistry-pedodontics\pedo-extractions\dental-trauma.webp"

- Delayed shedding: When a baby tooth remains in place and interferes with a permanent tooth's eruption.

  "C:\Jerush-dentoface-new-site-react\public\images\medical-services\dental-care\kids-dentistry-pedodontics\pedo-extractions\delayed-shedding.webp"

- Orthodontic reasons: In selected cases, removal may be part of a planned orthodontic treatment.

  "C:\Jerush-dentoface-new-site-react\public\images\medical-services\dental-care\kids-dentistry-pedodontics\pedo-extractions\orthodontic-reasons.webp"

- Problematic teeth: When a tooth's position or condition affects normal dental development.

  "C:\Jerush-dentoface-new-site-react\public\images\medical-services\dental-care\kids-dentistry-pedodontics\pedo-extractions\problematic-teeth.webp"

Not every loose, decayed or delayed baby tooth needs extraction. The dentist examines the tooth and may recommend restorative treatment or monitoring when appropriate.

How Is a Paediatric Tooth Extraction Performed?

The procedure is adapted to the child's age, comfort level, medical history, and the tooth being treated.

1. Dental examination: The dentist checks the tooth, surrounding gums and stage of tooth development. An X-ray may be recommended when necessary.

   "C:\Jerush-dentoface-new-site-react\public\images\medical-services\dental-care\kids-dentistry-pedodontics\pedo-extractions\dental-examination.webp"

2. Preparing the child: The procedure is explained in age-appropriate language to help the child feel more comfortable.

   "C:\Jerush-dentoface-new-site-react\public\images\medical-services\dental-care\kids-dentistry-pedodontics\pedo-extractions\preparing-the-child.webp"

3. Local anaesthesia: Appropriate anaesthesia is used to numb the area when indicated.

   "C:\Jerush-dentoface-new-site-react\public\images\medical-services\dental-care\kids-dentistry-pedodontics\pedo-extractions\local-anaesthesia.webp"

4. Tooth removal: The dentist gently removes the tooth using suitable instruments.

   "C:\Jerush-dentoface-new-site-react\public\images\medical-services\dental-care\kids-dentistry-pedodontics\pedo-extractions\tooth-removal.webp"

5. Bleeding control and aftercare: Gauze pressure is used as needed and the parent receives instructions for recovery.

   "C:\Jerush-dentoface-new-site-react\public\images\medical-services\dental-care\kids-dentistry-pedodontics\pedo-extractions\bleeding-control-after-care.webp"

Some children may require additional behaviour guidance or other anxiety-management options, depending on their needs and the complexity of treatment.

Caring for Your Child After Extraction

Most children recover well after a straightforward extraction. Following the dentist's instructions helps protect the extraction site while it heals.


- Apply gentle pressure with the gauze for the time the dentist advises.
- Offer soft, cool or lukewarm foods when your child is ready.
- Avoid vigorous rinsing, spitting, or disturbing the extraction site during the initial healing period.
- Avoid hard, crunchy foods that may irritate the area.
- Give medicines only as directed by the dentist or healthcare professional.
- Continue oral hygiene carefully, avoiding direct disturbance of the extraction site at first.

Contact the dental team if bleeding does not settle, pain becomes severe or worsens, or swelling or fever develops. Significant facial swelling or difficulty swallowing or breathing requires urgent medical attention.

"C:\Jerush-dentoface-new-site-react\public\images\medical-services\dental-care\kids-dentistry-pedodontics\pedo-extractions\caring-for-your-child.webp"

Will Removing a Baby Tooth Affect Permanent Teeth?

Baby teeth help children chew, speak, and maintain space for permanent teeth. When a baby tooth is removed earlier than expected, the dentist considers whether a space maintainer may be needed to reduce the risk of neighbouring teeth drifting into the space.



The need for a space maintainer depends on which tooth was removed, the child's age, the development of the permanent tooth, and the available space in the dental arch. Not every extraction requires one, so follow-up assessment is important.

Frequently Asked Questions

1. Do all baby teeth need to be extracted when they become loose?

No. Most baby teeth fall out naturally. Extraction is considered when there is a clinical reason, such as infection, significant damage, or interference with normal tooth development.

2. Is tooth extraction painful for children?

Local anaesthesia helps numb the area during the procedure. The child may experience some tenderness afterward, which can usually be managed with the dentist's aftercare advice.

3. How long does a paediatric tooth extraction take?

A simple extraction is generally brief, but the appointment may take longer if the child needs additional preparation, imaging, or more complex treatment.

4. Will my child need an X-ray before extraction?

Not always. The dentist recommends an X-ray when it is needed to evaluate the tooth roots, surrounding structures, or developing permanent teeth.

5. Can a badly decayed baby tooth be saved?

Sometimes. Depending on the extent of decay and the tooth's condition, a filling, pulp treatment, or crown may be possible. Extraction may be recommended when the tooth cannot be predictably restored.

6. Does every child need a space maintainer after extraction?

No. It depends on the tooth removed, the child's dental development, and the risk of losing space for the permanent tooth.

7. When can my child return to normal activities?

Many children can return to usual activities after a simple extraction once they feel comfortable. Follow the dentist's instructions regarding eating, physical activity, and oral hygiene.

Gentle Paediatric Dental Care at Jerush

When a tooth needs to be removed, the priority is to protect your child's comfort and support healthy dental development. Jerush Dentofacial & Cosmetic Laser Centre provides individualised dental care based on the child's needs and clinical findings.

CTA: Book a paediatric dental consultation to discuss your child's tooth pain, damaged baby tooth, or delayed tooth shedding.

Add "C:\Jerush-dentoface-new-site-react\public\images\doctors\priya2.webp"

SEO Details

- SEO title: Pedo Extractions for Children | Jerush Dental Care
- Meta description: Learn when children may need tooth extractions, how paediatric extractions are performed, recovery tips, and when a space maintainer may be recommended.
- Primary keyword: Pedo extractions

- Related keywords: Paediatric tooth extraction, baby tooth removal, children's dental extraction, primary tooth extraction, kids dentistry
- Image alt text: Paediatric dentist preparing for a child's tooth extraction in a child-friendly dental clinic.