

# Perfect Reconstruction of Discrete Framelet Transforms

## Chapter 1.1.2 Summary

Based on *Framelets and Wavelets*  
Bin Han

August 6, 2025

# Sequence Spaces

- $\ell(\mathbb{Z}) = \{v = \{v(k)\}_{k \in \mathbb{Z}} \mid v(k) \in \mathbb{C}\}$
- $\ell_0(\mathbb{Z}) = \{u \in \ell(\mathbb{Z}) \mid \text{only finitely many } u(k) \neq 0\}$
- filter bank:  $\{u_0, \dots, u_s\} \subset \ell_0(\mathbb{Z})$

- **Subdivision:**

$$[S_u v](n) = 2 \sum_{k \in \mathbb{Z}} v(k) u(n - 2k)$$

- **Transition:**

$$[T_u v](n) = 2 \sum_{k \in \mathbb{Z}} v(k) \overline{u(k - 2n)}$$

- **Discrete Framelet Transform (DFrT):**

$$\frac{1}{2} \sum_{\ell=0}^s S_{u_\ell} T_{\tilde{u}_\ell} v$$

# Perfect Reconstruction Condition

- the transform satisfies perfect reconstruction if

$$\frac{1}{2} \sum_{\ell=0}^s S_{u_\ell} T_{\tilde{u}_\ell} v = v \quad \forall v \in \ell(\mathbb{Z}).$$

# Main Theorem – Four Equivalent Statements

The following are equivalent:

- ① The filter bank  $(\{\tilde{u}_\ell\}, \{u_\ell\})$  has PR.
- ②  $\frac{1}{2} \sum_{\ell=0}^s S_{u_\ell} T_{\tilde{u}_\ell} v = v$  for all  $v \in \ell(\mathbb{Z})$ .
- ③ The identity holds for  $v = \delta$  and  $v = \delta(\cdot - 1)$ .
- ④ Frequency-domain identities for all  $\omega \in \mathbb{R}$ :

$$\sum_{\ell=0}^s \overline{\hat{\tilde{u}}_\ell(\omega)} \hat{u}_\ell(\omega) = 1,$$

$$\sum_{\ell=0}^s \overline{\hat{\tilde{u}}_\ell(\omega)} \hat{u}_\ell(\omega + \pi) = 0.$$

$$(i) \Rightarrow (ii) \Rightarrow (iii)$$

Trivial inclusions:

$$\ell_0(\mathbb{Z}) \subseteq \ell(\mathbb{Z}) \quad \text{and} \quad \delta, \delta(\cdot - 1) \in \ell_0(\mathbb{Z}).$$

(iii)  $\Rightarrow$  (iv)

- ① **Frequency Domain Conversion:** Using Fourier transforms of  $T_{\tilde{u}_\ell}$  and  $S_{u_\ell}$ :

$$\widehat{T_{\tilde{u}_\ell} v}(\omega) = \hat{v}(\omega/2) \overline{\hat{u}_\ell(\omega/2)} + \hat{v}(\omega/2 + \pi) \overline{\hat{u}_\ell(\omega/2 + \pi)},$$

$$\widehat{S_{u_\ell} w}(\omega) = \hat{w}(2\omega) \hat{u}_\ell(\omega).$$

- ② **Substitute Basis Signals:** Plugging  $v = \delta$  ( $\hat{v} = 1$ ) and  $v = \delta(\cdot - 1)$  ( $\hat{v} = e^{-i\omega}$ ) into the PR condition yields the system:

$$\sum_{\ell=0}^s \overline{\hat{u}_\ell(\omega)} \hat{u}_\ell(\omega) + \sum_{\ell=0}^s \overline{\hat{u}_\ell(\omega + \pi)} \hat{u}_\ell(\omega + \pi) = 2,$$

$$\sum_{\ell=0}^s \overline{\hat{u}_\ell(\omega)} \hat{u}_\ell(\omega) - \sum_{\ell=0}^s \overline{\hat{u}_\ell(\omega + \pi)} \hat{u}_\ell(\omega + \pi) = 0.$$

- ③ **Solve System:** Adding and subtracting these equations gives condition (iv).

(iv)  $\Rightarrow$  (ii) and (ii)  $\Rightarrow$  (i)

- (iv)  $\Rightarrow$  (ii): Condition (iv) implies the frequency identity holds for all  $v \in \ell_0(\mathbb{Z})$  by linearity of Fourier transform.
- (ii)  $\Rightarrow$  (i): Localization Argument: For any  $v \in \ell(\mathbb{Z})$ , truncate to local signal  $v_n \in \ell_0(\mathbb{Z})$  using finite support of filters. Apply (ii) to show:

$$v(n) = \frac{1}{2} \sum_{\ell=0}^s [S_{u_\ell} T_{\tilde{u}_\ell} v](n).$$