D	٨	D	D

## Operational symmetries of entangled states

To cite this article: Ilan Tzitrin et al 2020 J. Phys. A: Math. Theor. 53 095304

ซ้ายเพื่อใช้ <u>article on libite</u> for optiates ชองเรียะเรียะกับอะที่กิจาริรโร (To c artconts.) was downloaded from IP address Tf9.97.131 44lin 10/02/0

## Operational symmetries of entangled states

Ilan Tzitrin<sup>1</sup>, Aaron Z Goldberg<sup>1</sup> and Jesse C Cresswell<sup>1</sup>

1

/ , 1, so V  $U^T$  shares the unitarity of U. Unitarity for V, on the other hand, requ0res  $V^{\dagger}V$  1 for all V  $U^T$   $U^T$ 

By convention — is a  $d_A \times d_B$  diagonal matrix with the Schmidt coef cients — i listed in decreasing order.

If /  $_{AB}$  is fully entangled, then, by de nition, its Schmidt rank is maximal, viz. rank  $d_A$ . This implies that there exists a right inverse  $_R^{-1}$  such that



In this language, the Reeh Schlieder theorem says that the vacuum state of a QFT on Minkowski spacetime is cyclic and separating with respect to any local algebra  $A_{\rm b}$  of eld operators supported in an open neighbourhood V

7

that any local operation on  ${\cal A}$ 

This is the condition that the set

The von Neumann operator trace inequality tells us that /Tr  $X / \leq$  Tr /X/, and / $V U^* / U^*$  / is independent of V

for unitaries U and states / whose Schmidt matrices are not necessarily full rank. Arranging the bases such that the nonzero Schmidt coef cients are in the rst  $r \times r$  blocks of their corresponding

$$dU/U_{11}/ \qquad \qquad d \qquad 2^{2d-2} \quad 2d-2 \quad ^{-1} \\ 2 \quad d \quad \frac{1}{2} \qquad 2d-1 \quad d-1 \qquad (B.12)$$

If we try to deviate slightly from a separable state, with  $$diag$ 1-\ , \ \ ,0,\cdots$ 

- [51] Spreeuw R J C 1998 Found. Phys. 2 361
  [52] T ppel F, Aiello A, Marquardt C, Giacobino E and Leuchs G 2014 New J. Phys. 1 073019
  [53] Karimi E and Boyd R W 2015 Science 3 0 1172
  [54] Eberly J2 gs 6h, Qian.8 (J2 gs 6X-lNe)1, Qas[(Karimy )3A.8 (J2 gs 6A)80 436J2 gs 6A