
patient
case
time
treatment
found
used
group
normal
p less
associated
performed
compared
obtained
control
showed
observed
result
increase
seen
presented
number
women
CONCLUSION
evaluated
reduced
technique
pattern
prior
type
hypertension
mm Hg
due
children
mild
measured
similar
following
response
examined
multiple
problem
operation
operating room
value
either
disease
one patient
equal
using
data
therapy
decrease
occurred
low
subject
incidence
including
risk factor
death
week
period
two patient
primary
difference
within
development
removed
n
conclude
rate
three
first
left ventricular
old
year
history
coronary artery
four
demonstrated
induced
tissue
five
area
long term
described
incision
draped
procedure
one
lesion
year
initial
percent
heart rate
high
hospital
hour
report
negative
minute
acute
mean
month
determined
surgery
effect
symptom
possible
given
will
studied
secondary
evidence
cause
less
site
risk
myocardial infarction
whereas
suggest
human
reviewed
appear
greater
stroke
point
treated
well
finding
done
discussed
severe
chronic
respectively
condition
revealed
mm
second
present
evaluation
left
need
rat
new
change
activity
may
tumor
closed
studies
method
management
applied
placed
review
positive
determine
decreased
addition
skin
complication
without
diagnosis
reported
presence
age
pain
wound
related
role
taken
early
clinical
total
use
level
two
made
developed
Thus
concentration
blood pressure
considered
follow
reduction
abnormal
test
identified
infection
significant
increased
region
Although
detected