

# Prediction

Wenda Zhou

June 15, 2017

# What is prediction?

Attempt to predict a random outcome with potentially unknown mechanism (may be viewed as random) from covariates.

# Prediction in statistics

- ▶ One of fastest growing fields in statistics
- ▶ Driven by big data applications
- ▶ More used in non-scientific contexts

# Prediction vs Estimation

## Estimation

- ▶ Understand “true state of nature”.
- ▶ Attempt to understand latent parameter

## Prediction

- ▶ Attempt to guess another random quantity
- ▶ Prediction outcome is visible and observable

# Prediction vs Estimation

## Example

### Clinical trial: estimation

Given a person with some characteristics, obtain the probability of success of the treatment on that person.

# Prediction vs Estimation

## Example

### Clinical trial: estimation

Given a person with some characteristics, obtain the probability of success of the treatment on that person.

### Clinical trial: prediction

For the next person that the doctor sees, predict whether the treatment will succeed.

# Prediction vs Estimation

## Example

### Clinical trial: estimation

Given a person with some characteristics, obtain the probability of success of the treatment on that person.

### Clinical trial: prediction

For the next person that the doctor sees, predict whether the treatment will succeed.

Two problems closely related.

# What is a good prediction?

Define what a good prediction is.

## Loss

Loss defines how bad a prediction was compared to the truth.



# What is a good prediction?

Define what a good prediction is.

## Loss

Loss defines how bad a prediction was compared to the truth.

## Square loss

Suppose we predict  $\hat{y}$ , but the true value was  $y$ . The square loss is defined as:

$$L(y, \hat{y}) = (y - \hat{y})^2 \quad (1)$$

# What is a good prediction?

Define what a good prediction is.

## Loss

Loss defines how bad a prediction was compared to the truth.

## Square loss

Suppose we predict  $\hat{y}$ , but the true value was  $y$ . The square loss is defined as:

$$L(y, \hat{y}) = (y - \hat{y})^2 \quad (1)$$

## 0 – 1 loss

For binary outcomes: 0 loss if we are right, 1 loss if we are wrong.

## First example: prediction using linear models

Suppose that we have an ordinary linear model.

Then can predict a new outcome  $\hat{y}$  with covariates  $x$  by putting:

$$\hat{y} = \hat{\alpha} + \hat{\beta}x \quad (2)$$

## First example: prediction using linear models

Suppose that we have an ordinary linear model.

Then can predict a new outcome  $\hat{y}$  with covariates  $x$  by putting:

$$\hat{y} = \hat{\alpha} + \hat{\beta}x \quad (2)$$

Unlike estimation, prediction is less worried about confounding.

# Evaluating our prediction model

Easier to evaluate prediction models: can check whether we are right.

However, need care to separate the following quantities:

## Training error

Loss incurred on training dataset.

## Testing error

Loss incurred on new dataset.

# Evaluating our prediction model

## Warning

It is always true that:

$$\text{Training error} < \text{Testing error} \quad (3)$$

# Evaluating our prediction model

## Warning

It is always true that:

$$\text{Training error} < \text{Testing error} \quad (3)$$

In fact, training error is not a good measure of testing error.

# Overfitting

Will often want to evaluate our model to select between multiple models.

Picking the model with lowest training error may not always be the best. May have overfitting.



# Overfitting

Will often want to evaluate our model to select between multiple models.

Picking the model with lowest training error may not always be the best. May have overfitting.

## Bias-variance and prediction

Although the bias-variance decomposition applies to estimation, prediction faces a very similar problem in overfitting vs. underfitting.

# Classification

Take a binary (yes/no) decision – or discrete decision (which group is the subject in).

Very common case, so has specific vocabulary.

# Classification

Can separate the performance of classification in two parts:

**Precision** The precision  $p$  is the proportion of correct results among all predicted yes.

**Recall** The recall  $r$  is the proportion of correct results among all true yes.

# Classification

Can separate the performance of classification in two parts:

**Precision** The precision  $p$  is the proportion of correct results among all predicted yes.

**Recall** The recall  $r$  is the proportion of correct results among all true yes.

Combine these measures into the  $F1$  score:

$$F_1 = 2 \frac{pr}{p + r} \quad (4)$$

# Linear models for prediction

If have linear model:

$$y = \alpha + \beta x \quad (5)$$

Can generate predictions for new individual with covariates  $x_{\text{new}}$ :

$$\hat{y} = \hat{\alpha} + \hat{\beta} x_{\text{new}} \quad (6)$$

# Linear models for prediction

Prediction is less sensitive to confounding:  $\hat{\beta}$  can be completely wrong and still helpful!

# Linear models for prediction

Prediction is less sensitive to confounding:  $\hat{\beta}$  can be completely wrong and still helpful!

## Example: diamonds

Saw that diamonds with worse color rating were more expensive.  
Incorrect inference, but still helpful for prediction.

# Uncertainty in prediction

Wish to quantify uncertainty of our prediction.

## Prediction intervals

Interval likely to contain the true value of the **outcome**.



# Uncertainty in prediction

Wish to quantify uncertainty of our prediction.

## Prediction intervals

Interval likely to contain the true value of the **outcome**.

Unlike in estimation, even with infinite amounts of data, still uncertainty in prediction. Aggregate uncertainty in estimation and natural randomness of the outcome.