

# Uncertainty in estimation

Wenda Zhou

June 2, 2017

# Uncertainty in estimation

We have seen how to compute a single estimate  $\hat{\theta}$  of a parameter.  
Can we also quantify the uncertainty in the estimate?

# Uncertainty in estimation

Example: coin toss

Suppose  $X$  follows a binomial distribution with parameters  $n$  and  $p$ .  
We have seen the m.l.e. estimator for  $p$  given by:

$$\hat{p} = \frac{1}{n}X \quad (1)$$

# Uncertainty in estimation

Example: coin toss

Suppose  $X$  follows a binomial distribution with parameters  $n$  and  $p$ .  
We have seen the m.l.e. estimator for  $p$  given by:

$$\hat{p} = \frac{1}{n}X \quad (1)$$

Intuition: estimator should be better for  $n = 100$  than  $n = 10$ .

# Uncertainty in estimation

Example: coin toss

Recall: for unbiased estimators, lower variance is better.

Let us compute:

$$\text{Var}_p \hat{p} = \frac{1}{n} p(1 - p) \quad (2)$$

# Uncertainty in estimation

Example: coin toss

Recall: for unbiased estimators, lower variance is better.

Let us compute:

$$\text{Var}_p \hat{p} = \frac{1}{n} p(1 - p) \quad (2)$$

Larger  $n$  leads to lower variance and better estimation. Hence may also wish to report standard error.

# Confidence intervals

Instead of giving a single point as estimate: give range of values.

## Confidence interval

A confidence interval is an interval that contains the true parameter with high probability.

# Confidence intervals

Instead of giving a single point as estimate: give range of values.

## Confidence interval

A confidence interval is an interval that contains the true parameter with high probability.

Note: the interval depends on the data, thus is random. The parameter is fixed (although unknown)



# Confidence intervals

Example: toin coss

Suppose that we toss a coin 100 times, and observe exactly 50 heads.

The computed 95% confidence interval is:  $[0.398, 0.602]$ .

# Confidence intervals

Example: coin toss

Suppose that we toss a coin 100 times, and observe exactly 50 heads.

The computed 95% confidence interval is:  $[0.398, 0.602]$ .

Suppose instead that we toss a coin 1000 times, and observe exactly 500 heads.

The computed 95% confidence interval is:  $[0.469, 0.531]$ .

# Confidence intervals

Example: toin coss

Suppose that we toss a coin 100 times, and observe exactly 50 heads.

The computed 95% confidence interval is:  $[0.398, 0.602]$ .

Suppose instead that we toss a coin 1000 times, and observe exactly 500 heads.

The computed 95% confidence interval is:  $[0.469, 0.531]$ .

The second confidence interval is narrower, reflecting our increased certainty.

# Confidence intervals

## Interpreting the 95%

Let us write the confidence interval as  $[a(X), b(X)]$ .

Then,  $[a(X), b(X)]$  is a 95% confidence interval if:

$$P_p(a(X) \leq p \leq b(X)) = 0.95 \quad (3)$$

# Confidence intervals

## Interpreting the 95%

Let us write the confidence interval as  $[a(X), b(X)]$ .

Then,  $[a(X), b(X)]$  is a 95% confidence interval if:

$$P_p(a(X) \leq p \leq b(X)) = 0.95 \quad (3)$$

If we repeat the experiment and compute a confidence interval in the same manner, it will cover the true value 95% of the time on average.

# Computing a confidence interval

## Normal distribution

Suppose that we observe a single observation:

$$\hat{\theta} \sim \mathcal{N}(\theta, \sigma_{\theta}^2) \quad (4)$$

$\sigma_{\theta}$  is the standard error (known).

Let us compute a confidence interval for  $\theta$ .

# Computing a confidence interval

## Normal distribution

A  $(1 - \alpha)$  confidence interval is given by:

$$[\hat{\theta} - \sigma_{\theta} z_{1-\alpha/2}, \hat{\theta} + \sigma_{\theta} z_{1-\alpha/2}] \quad (5)$$

Where  $z_{\beta}$  is such that  $P(Z \leq z_{\beta}) = \beta$  (quantile).

# Computing a confidence interval

## Example

Suppose that after running an experiment, we obtain an estimate of  $\hat{\theta} = 4.5$  and standard error 1.2. Compute a 95% estimate for  $\theta$ .

We have that:  $\alpha = 0.025$ , hence  $z_{1-\alpha/2} = 1.96$ .

Hence we have that a 95% confidence interval for this value is:

$$[4.5 - 1.96 \times 1.2, 4.5 + 1.96 \times 1.2] = [2.15, 6.85] \quad (6)$$