

Indefinite Integration

1 Fundamental Definition of Indefinite Integration

If f and F are functions such that $\frac{d}{dx}(F(x)) = f(x)$ then F is anti-derivative of f w.r.t. x symbolically,

$$\int f(x) dx = F(x) + C$$

where C is the constant of Integration

2 Anti-Derivatives of Some Standard Functions

i. $\int k \cdot f(x) dx = k \cdot \int f(x) dx$

ii. $\int [f_1(x) \pm f_2(x) \pm f_3(x) \pm \dots \pm f_n(x)] dx = \int f_1(x) dx \pm \int f_2(x) dx \pm \int f_3(x) dx \pm \dots \pm \int f_n(x) dx$

iii. $\int x^n dx = \frac{x^{n+1}}{n+1} + C$ $n \neq -1$

iv. $\int \frac{1}{x} dx = \ln |x| + C$

v. $\int e^x dx = e^x + C$

vi. $\int a^x dx = \frac{a^x}{\ln a} + C$

vii. $\int \sin x dx = -\cos x + C$

viii. $\int \cos x dx = \sin x + C$

$$\text{ix. } \int \sec^2 x \, dx = \tan x + C$$

$$\text{x. } \int \csc^2 x \, dx = -\cot x + C$$

$$\text{xi. } \int \sec x \tan x \, dx = \sec x + C$$

$$\text{xii. } \int \csc x \cot x \, dx = -\csc x + C$$

$$\text{xiii. } \int \cot x \, dx = \ln |\sin x| + C$$

$$\text{xiv. } \int \tan x \, dx = -\ln |\cos x| + C$$

$$\text{xv. } \int \sec x \, dx = \ln |\sec x + \tan x| + C$$

$$\text{xvi. } \int \csc x \, dx = \ln |\csc x - \cot x| + C$$

$$\text{xvii. } \int \frac{1}{\sqrt{a^2 - x^2}} \, dx = \sin^{-1} \left(\frac{x}{a} \right) + C$$

$$\text{xviii. } \int \frac{-1}{\sqrt{a^2 - x^2}} \, dx = \cos^{-1} \left(\frac{x}{a} \right) + C$$

$$\text{xix. } \int \frac{1}{a^2 + x^2} \, dx = \frac{1}{a} \tan^{-1} \left(\frac{x}{a} \right) + C$$

$$\text{xx. } \int \frac{-1}{a^2 + x^2} \, dx = \frac{1}{a} \cot^{-1} \left(\frac{x}{a} \right) + C$$

$$\text{xxi. } \int \frac{1}{x\sqrt{x^2 - a^2}} \, dx = \frac{1}{a} \sec^{-1} \left(\frac{x}{a} \right) + C$$

$$\text{xxii. } \int \frac{-1}{x\sqrt{x^2 - a^2}} \, dx = \frac{1}{a} \csc^{-1} \left(\frac{x}{a} \right) + C$$

$$\text{xxiii. } \int \sqrt{x} \, dx = \frac{2x\sqrt{x}}{3} + C$$

$$\text{xxiv. } \int \frac{dx}{x^2 - 1} = \ln \left| \frac{x-1}{x+1} \right| + C$$

$$\text{xxv. } \int \frac{dx}{\sqrt{1+x^2}} = \ln \left| x + \sqrt{x^2 + 1} \right| + C$$

$$\text{xxvi. } \int \frac{dx}{\sqrt{x^2-1}} = \ln \left| x + \sqrt{x^2-1} \right| + C$$

$$\text{xxvii. } \int \sqrt{1-x^2} dx = \frac{1}{2}x\sqrt{1-x^2} + \frac{1}{2}\sin^{-1}x + C$$

$$\text{xxviii. } \int \sqrt{x^2-1} dx = \frac{1}{2}x\sqrt{x^2-1} - \frac{1}{2}\ln \left| x + \sqrt{x^2-1} \right| + C$$

Important Results

1. If $F_1(x)$ and $F_2(x)$ are two anti-derivatives of a function $f(x)$, then $F_1(x)$ and $F_2(x)$ only differ by a constant, i.e.

$$F_1(x) - F_2(x) = C$$

where, C is a \mathbb{R} constant.

2. If $f(x)$ is continuous $\forall x \in D_f$ and,

$$\int f(x) dx = F(x) + C,$$
then $F(x)$ always exists and is continuous.
3. If $f(x)$ is discontinuous at $x = x_1$, then its anti-derivative can be continuous at $x = x_1$.
4. Anti-derivative of a periodic function may not be periodic.

3 Methods of Integration

3.1 u substitution

Integrals of form,

$$I = \int f(g(x)) \cdot g'(x) dx$$

Can be solved by, the substitution,

$$u = g(x)$$

Differentiating both sides w.r.t. x ,

$$du = g'(x)dx$$

Now,

$$I = \int f(u) du$$

3.2 Integrals of form

$$\int \frac{dx}{ax^2 + bx + c}, \int \frac{dx}{\sqrt{ax^2 + bx + c}}, \int \sqrt{ax^2 + bx + c} dx$$

Using completing the square,

$$ax^2 + bx + c = a \left[\left(x + \frac{b}{2a} \right)^2 + \frac{c}{a} - \frac{b^2}{4a} \right]$$

Now, using u sub, let

$$u = x + \frac{b}{2a}$$

The transformed integral can be integrated using previous methods.

3.3 Integrals of form

$$\int \frac{px + q}{ax^2 + bx + c} dx, \int \frac{px + q}{\sqrt{ax^2 + bx + c}} dx, \int (px + q) \sqrt{ax^2 + bx + c} dx$$

$$px + q = \lambda \frac{d}{dx} (ax^2 + bx + c) + \mu$$

Now, after finding λ, μ ,

For the 1st part, use u sub,

Let

$$u = ax^2 + bx + c$$

2nd part of the integral can be integrated using previous methods.

3.4 Integrals of form

$$\int \frac{K(x)}{ax^2 + bx + c} dx$$

where $\deg(K(x)) \geq 2$

By polynomial long division

$$\frac{K(x)}{ax^2 + bx + c} = Q(x) + \frac{R(x)}{ax^2 + bx + c}$$

Now, $\deg(R(x)) \leq 1$, Now, the integral

$$\int \frac{R(x)}{ax^2 + bx + c} dx$$

can be integrated using previous methods.

3.5 Integrals of form

$$\int \frac{ax^2 + bx + c}{px^2 + qx + r} dx, \int \frac{ax^2 + bx + c}{\sqrt{px^2 + qx + r}} dx, \int (ax^2 + bx + c) \sqrt{px^2 + qx + r} dx$$

$$ax^2 + bx + c = \lambda (px^2 + qx + r) + \mu \frac{d}{dx}(px^2 + qx + r) + \gamma$$

3.6 Trig. Integrals

3.6.1 Integrals of form

$$\int \frac{dx}{a \cos^2 x + b \sin^2 x}, \int \frac{dx}{a + b \sin^2 x}, \int \frac{dx}{a + b \cos^2 x}$$

$$\int \frac{dx}{(a \sin x + b \cos x)^2}, \int \frac{dx}{a + b \sin^2 x + c \cos^2 x}$$

Steps -

1. Multiply numerator and denominator by $\sec^2 x$
2. Replace $\sec^2 x$ (if any) by $1 + \tan^2 x$ except the one multiplied in step 1.
3. Let $u = \tan x$, then $du = \sec^2 x dx$

Now, the transformed integral can be integrated using previous methods.

3.6.2 Integrals of form

$$\int \frac{dx}{a \sin x + b \cos x}, \int \frac{dx}{a + b \sin x}, \int \frac{dx}{a + b \cos x}, \int \frac{dx}{a \sin x + b \cos x + c}$$

Steps -

1. Replace $\sin x = \frac{2 \tan x/2}{1 + \tan^2 x/2}$ and $\cos x = \frac{1 - \tan^2 x/2}{1 + \tan^2 x/2}$
2. Let $u = \tan x/2$, $du = \frac{1}{2} \sec^2 x/2 dx$ is already present in the numerator.

Now, the transformed integral can be integrated using previous methods.

3.6.2.1 Alternative Method to Integrate

$$I = \int \frac{dx}{a \sin x + b \cos x}$$

$$a \sin x + b \cos x = \sqrt{a^2 + b^2} \sin \left(x + \tan^{-1} \left(\frac{b}{a} \right) \right)$$

$$I = \frac{1}{\sqrt{a^2 + b^2}} \int \csc \left(x + \tan^{-1} \left(\frac{b}{a} \right) \right) dx$$

$$I = \frac{1}{\sqrt{a^2 + b^2}} \ln \left| \tan \left(\frac{x}{2} + \frac{1}{2} \tan^{-1} \left(\frac{b}{a} \right) \right) \right| + C$$

3.6.3 Integrals of form

$$\int \frac{p \cos x + q \sin x + r}{a \cos x + b \sin x + c} dx, \int \frac{p \cos x + q \sin x}{a \cos x + b \sin x} dx$$

Steps for (i),

1. Express $Numerator = \lambda Denominator + \mu Derivative of denominator + \gamma$

Now, the transformed integral can be integrated using previous methods.

Steps for (ii),

1. Express $Numerator = \lambda Denominator + \mu Derivative of Denominator$

Now, the transformed integral can be integrated using previous methods.

3.7 Integration by parts