

# Coalgebraic Modal Logic over Set

2026 Dutch Winter School on Logic and Verification

---

Helle Hvid Hansen

Bernoulli Institute for Maths, CS and AI, University of Groningen, NL



# Introduction

---

Previous lectures:

**Universal Coalgebra: a theory of systems**  $X \rightarrow F(X)$

- developed uniformly, parametric in functor  $F$  (system type);
- captures observable system behaviour;
- canonical notion of behavioural equivalence and bisimulation.

**Coalgebraic modal logic: framework for specifying coalgebra behaviours**

**Modal logics are coalgebraic:**  $\frac{\text{Equational Logic}}{\text{Algebra}} = \frac{\text{Coalgebraic Modal Logic}}{\text{Coalgebra}}$

## At the level of universal coalgebra:

- What is a modality?
- Which properties do we want to express? Which ones can be expressed?
- Which concrete results (e.g. Hennessy-Milner theorem) hold more generally?
- ... and under which conditions?

This lecture: **Set**-based coalgebras  $X \rightarrow F(X)$  (i.e.,  $X$  is a set,  $F: \mathbf{Set} \rightarrow \mathbf{Set}$ ).  
(intuitions, covers many examples relevant for applications)

Next lecture: Beyond **Set** (more abstract setting)

## Overview of lecture

- Modal semantics as predicate liftings
- Invariance and expressiveness results
- (Modal semantics via relation lifting)
- Concluding remarks

Focus on definitions, results  
examples.

## After this lecture...

- You can explain the basic concepts of coalgebraic modal logic;
- You can explain some basic invariance and expressiveness results of coalgebraic modal logic;
- You can apply the basic theory of coalgebraic modal logic to concrete system types.

## Modalities as Predicate Liftings

---

## Labelled transition systems are $\mathcal{P}^A$ -Coalgebras

Covariant powerset  $\mathcal{P}: \mathbf{Set} \rightarrow \mathbf{Set}$

$$\begin{aligned} \mathcal{P}(X) &= \{U \mid U \subseteq X\} \\ \mathcal{P}(f: X \rightarrow Y) &= f[-]: \mathcal{P}(X) \rightarrow \mathcal{P}(Y) \\ &\quad \text{(direct image)} \end{aligned}$$

Exponentiation  $(-)^A: \mathbf{Set} \rightarrow \mathbf{Set}$

$$\begin{aligned} X^A &= \{g: X \rightarrow A\} \\ (f: X \rightarrow Y)^A &= f \circ (-): X^A \rightarrow Y^A \\ &\quad \text{(pre-composition)} \end{aligned}$$

LTS is coalgebra  $R: X \rightarrow \mathcal{P}(X)^A$ :  $R(x)(a) = \{y \in X \mid x \xrightarrow{a} y\} =: R_a(x)$

## Semantics of modal formulas:

Let  $\llbracket \varphi \rrbracket = \{x \in X \mid \mathbb{M}, x \models \varphi\}$  (truth-set of  $\varphi$ ).

$$\begin{aligned} (X, R), x \models [a]\varphi &\quad \text{iff} \quad R_a(x) \subseteq \llbracket \varphi \rrbracket &\quad \text{iff} \quad R(x) \in \{g \in \mathcal{P}(X)^A \mid g(a) \subseteq \llbracket \varphi \rrbracket\} \\ (X, R), x \models \langle a \rangle \varphi &\quad \text{iff} \quad R_a(x) \cap \llbracket \varphi \rrbracket \neq \emptyset &\quad \text{iff} \quad R(x) \in \{g \in \mathcal{P}(X)^A \mid g(a) \cap \llbracket \varphi \rrbracket \neq \emptyset\} \end{aligned}$$

## Markov chains are $\mathcal{D}_\omega$ -Coalgebras

Finitary distribution functor  $\mathcal{D}_\omega: \mathbf{Set} \rightarrow \mathbf{Set}$

$$\mathcal{D}_\omega(X) = \{\mu \mid X \rightarrow [0, 1] \mid \text{supp}(\mu) \text{ finite}, \sum_{x \in X} \mu(x) = 1\}$$

$$\mathcal{D}_\omega(f: X \rightarrow Y) = \mathcal{D}_\omega(f): \mathcal{D}_\omega(X) \rightarrow \mathcal{D}_\omega(Y)$$

$$\mathcal{D}_\omega(f)(\mu)(y) = \sum_{x \in X, f(x)=y} \mu(x)$$

Markov chain is coalgebra  $\gamma: X \rightarrow \mathcal{D}_\omega(X)$ :  $\gamma(x)(x') = p \iff x \xrightarrow{p} x'$

## Semantics of modal formulas:

Let  $\llbracket \varphi \rrbracket = \{x \in X \mid (X, \gamma), x \models \varphi\}$  (truth-set of  $\varphi$ ).

Let  $r \in [0, 1]$ . Consider modality  $L_r$  ("probability at least  $r$ ")

$$(X, \gamma), x \models L_r \varphi \iff \sum_{x \in \llbracket \varphi \rrbracket} \gamma(x) \geq r \iff \gamma(x) \in \{\mu \in \mathcal{D}_\omega(X) \mid \sum_{x \in \llbracket \varphi \rrbracket} \mu(x) \geq r\}$$



# Neighbourhood Structures are Coalgebras

- Contravariant powerset functor  $\mathcal{Q}: \mathbf{Set}^{\text{op}} \rightarrow \mathbf{Set}$

$$\mathcal{Q}(X) = \mathcal{P}(X) (= 2^X)$$

$$\mathcal{Q}(f) = f^{-1}[-]: \mathcal{Q}(Y) \rightarrow \mathcal{Q}(X) \quad (\text{inverse image})$$

- Neighbourhood frames are  $\mathcal{N}$ -coalgebras  $X \rightarrow \mathcal{N}(X)$  where

$$\mathcal{N}(X) = (\mathcal{Q} \circ \mathcal{Q}^{\text{op}})(X)$$

$$\mathcal{N}(f) = (f^{-1})^{-1}[-]: \mathcal{N}(X) \rightarrow \mathcal{N}(Y) \quad (\text{double inverse image})$$

$$U \in \mathcal{N}(f)(H) \text{ iff } f^{-1}[U] \in H$$

- Monotonic neighbourhood frames are coalgebras  $X \rightarrow \mathcal{M}(X)$  for

$$\mathcal{M}(X) = \{H \in \mathcal{N}(X) \mid H \text{ closed under supersets}\}$$

$$\mathcal{M}(f) = (f^{-1})^{-1}[-]: \mathcal{M}(X) \rightarrow \mathcal{M}(Y) \quad (\text{double inverse image})$$

Remark:  $\mathcal{N}$  and  $\mathcal{M}$  do not preserve weak pullbacks, hence  $\text{behav.equiv.} \not\Rightarrow$  bisimilarity.

In neighbourhood model  $\mathbb{M} = (X, N: X \rightarrow \mathcal{N}(X), v: X \rightarrow \mathcal{P}(\mathbf{Prop}))$ :

$$\mathbb{M}, x \models \Box\varphi \quad \text{iff} \quad \llbracket \varphi \rrbracket \in N(x) \quad \text{iff} \quad N(x) \in \{H \in \mathcal{N}(X) \mid \llbracket \varphi \rrbracket \in H\}$$

If  $\mathbb{M} = (X, N: X \rightarrow \mathcal{M}(X), v: X \rightarrow \mathcal{P}(\mathbf{Prop}))$  monotonic, then equivalent with:

$$\begin{aligned} \mathbb{M}, x \models \Box\varphi \quad &\text{iff} \quad \exists U \in N(x) : U \subseteq \llbracket \varphi \rrbracket \\ &\text{iff} \quad N(x) \in \{H \in \mathcal{M}(X) \mid \exists U \in N(x) : U \subseteq \llbracket \varphi \rrbracket\} \end{aligned}$$

## General picture:

In coalgebraic model  $\mathbb{X} = (X, \gamma: X \rightarrow F(X), v: X \rightarrow \mathcal{P}(\mathbf{Prop}))$

$$\mathbb{X}, x \models \heartsuit\varphi \quad \text{iff} \quad \gamma(x) \in \{f \in F(X) \mid \text{condition involving } \llbracket \varphi \rrbracket\}$$

Predicate  $\llbracket \varphi \rrbracket$  over state space  $X$  is lifted to predicate  $F(X)$ .

A  $n$ -ary predicate lifting for  $F$  is a natural transformation  $\lambda: \mathcal{Q}^n \Rightarrow \mathcal{Q} \circ F$  (recall  $\mathcal{Q}$  is contravariant powerset), that is,  $\lambda$  is a family of set-indexed functions  $\lambda_X: (2^X)^n \rightarrow 2^{F(X)}$  such that:

$$\begin{array}{ccccc} X & & (2^X)^n & \xrightarrow{\lambda_X} & 2^{F(X)} \\ \forall f \downarrow & & \uparrow (f^{-1})^n & & \uparrow F(f)^{-1} \\ Y & & (2^Y)^n & \xrightarrow{\lambda_Y} & 2^{F(Y)} \end{array}$$

# Coalgebraic Modal Logic via Predicate Liftings

Coalgebraic modal logic means coalgebraic semantics of modal languages.

## Syntax

Given a modal signature  $\Lambda$  (i.e., collection of modal operators with arities), and a set  $\mathbf{Prop}$  of propositional variables, the set  $\mathcal{L}(\Lambda, \mathbf{Prop})$  of formulas over  $\Lambda$  and  $\mathbf{Prop}$  is:

$$\mathcal{L}(\Lambda, \mathbf{Prop}) \ni \varphi ::= p \in \mathbf{Prop} \mid \top \mid \neg\varphi \mid \varphi \wedge \varphi \mid \heartsuit(\varphi_1, \dots, \varphi_n), \quad \heartsuit \in \Lambda, \text{ } n\text{-ary}$$

**Coalgebraic semantics:** We want to interpret formulas in  $F$ -coalgebra model

$$\mathbb{X} = (X, \gamma: X \rightarrow F(X), v: \mathbf{Prop} \rightarrow \mathcal{P}(X))$$

which corresponds to  $F \times \mathcal{P}(\mathbf{Prop})$ -coalgebra  $\langle \gamma: X \rightarrow F(X), \hat{v}: X \rightarrow \mathcal{P}(\mathbf{Prop}) \rangle$ .

(We can take atomic props to be part of the functor.)

# Coalgebraic Modal Logic via Predicate Liftings

Coalgebraic semantics of  $\mathcal{L}(\Lambda, \mathbf{Prop})$  consists of:

- a functor  $F: \mathbf{Set} \rightarrow \mathbf{Set}$
- for every  $n$ -ary modal operator  $\heartsuit \in \Lambda$ , a natural transformation

$$\lambda^{\heartsuit} : Q^n \Rightarrow QF.$$

**Truth in  $F$ -model**  $\mathbb{X} = (X, \gamma : X \rightarrow F(X), v : X \rightarrow \mathcal{P}(\mathbf{Prop}))$

$$\mathbb{X}, x \models p \quad \text{iff} \quad x \in v(p) \quad \text{for } p \in \mathbf{Prop}$$

$$\vdots$$

$$\mathbb{X}, x \models \heartsuit(\varphi_1, \dots, \varphi_n) \quad \text{iff} \quad \gamma(x) \in \lambda_X^{\heartsuit}(\llbracket \varphi_1 \rrbracket, \dots, \llbracket \varphi_n \rrbracket)$$

# Predicate Lifting Examples

HML box:

$$\begin{aligned}\lambda_X^{[a]}(U) &= \{g \in \mathcal{P}(X)^A \mid g(a) \subseteq U\} \\ (X, R), x \models [a]\varphi &\iff R(x) \in \lambda_X^{[a]}(\llbracket \varphi \rrbracket)\end{aligned}$$

HML diamond:

$$\begin{aligned}\lambda_X^{\langle a \rangle}(U) &= \{g \in \mathcal{P}(X)^A \mid g(a) \cap U \neq \emptyset\} \\ (X, R), x \models \langle a \rangle \varphi &\iff R(x) \in \lambda_X^{\langle a \rangle}(\llbracket \varphi \rrbracket)\end{aligned}$$

Probabilistic modality:

$$\begin{aligned}\lambda_X^{L_r}(U) &= \{\mu \in \mathcal{D}_\omega(X) \mid \sum_{x \in U} \mu(x) \geq r\} \\ (X, \gamma), x \models L_r \varphi &\iff \gamma(x) \in \lambda_X^{L_r}(\llbracket \varphi \rrbracket)\end{aligned}$$

Neighbourhood modality:

$$\begin{aligned}\lambda_X^\square(U) &= \{H \in \mathcal{N}(X) \mid U \in H\} \\ (X, N), x \models \square \varphi &\iff N(x) \in \lambda_X^\square(\llbracket \varphi \rrbracket)\end{aligned}$$

Exercise: Check naturality condition for the above.

# Modalities as "Allowed Patterns", cf. [Schröder'08],[Gumm]

Let  $C_{\heartsuit} \subseteq F(2^n)$  specify which 0-1 patterns of  $F$ -structures are "allowed" by  $\heartsuit$ .

(For  $U \subseteq X$ , characteristic function  $\chi_U: X \rightarrow 2$ )

$$\begin{array}{ccc} X & \xrightarrow{\langle \chi_{\llbracket \varphi_1 \rrbracket}, \dots, \chi_{\llbracket \varphi_n \rrbracket} \rangle} & 2^n \\ \gamma \downarrow & & \\ F(X) & \xrightarrow{F\langle \chi_{\llbracket \varphi_1 \rrbracket}, \dots, \chi_{\llbracket \varphi_n \rrbracket} \rangle} & F(2^n) \xrightarrow{\chi_{C_{\heartsuit}}} 2 \end{array}$$

Define:  $(X, \gamma), x \models \heartsuit(\varphi_1, \dots, \varphi_n) \iff F(\chi_{\llbracket \varphi_1 \rrbracket}, \dots, \chi_{\llbracket \varphi_n \rrbracket})(\gamma(x)) \in C_{\heartsuit}$

**Exercise:** Let  $F(X) = X \times \{a, b\} \times X$  (labelled binary trees).

What are the possible unary patterns ( $n = 1$ )?

Which  $C \subseteq F(2)$  corresponds to:

$\heartsuit\varphi$ : "the label is  $a$  and  $\varphi$  is true at some child";

$C_{\heartsuit} = \{(1, a, 0), (0, a, 1), (1, a, 1)\}$

$\clubsuit\varphi$ : "if the label is  $b$  then  $\varphi$  is true at both children";

$C_{\clubsuit} = \{(1, b, 1), (0, a, 0), (1, a, 0), (0, a, 1), (1, a, 1)\}$

## Exercise:

For  $\mathcal{P}$ , the unary 0-1 patterns are subsets  $C \subseteq \mathcal{P}(2) = \mathcal{P}(\{0, 1\})$ .

The "singleton patterns" (i.e.,  $c \in \mathcal{P}(2)$ ) are:

- $\emptyset$       No successors
- $\{0\}$      Some successors and  $\varphi$  is false at all
- $\{1\}$      Some successors and  $\varphi$  is true at all
- $\{0, 1\}$    Some successors where  $\varphi$  is false, and some where  $\varphi$  is true

Describe informally the modality represented by each singleton pattern.

Which subsets  $C \subseteq \mathcal{P}(2)$  correspond to  $\Box\varphi$  and  $\Diamond\varphi$ ?



# Correspondence and Counting Modalities

Via Yoneda Lemma, 1-1 correspondence:

$$\frac{\text{predicate liftings } \lambda^\heartsuit : (2^-)^n \Rightarrow 2^{F-}}{\text{subsets } C_\heartsuit \in 2^{F(2^n)}}$$

It also tells us **how many predicate liftings**, there are.

E.g. Number of distinct unary modalities/predicate liftings for  $\mathcal{P}$  is  $2^{\mathcal{P}(2)} = 16$

Exercise. The correspondence can be proven directly.

- For  $C \subseteq F(2^n)$ , let  $\lambda_X^C(U_1, \dots, U_n) := F(\chi_{U_1}, \dots, \chi_{U_n})^{-1}[C]$ .
- For  $n$ -ary  $\lambda$ , let  $C_\lambda := \lambda_{2^n}(\pi_1^{-1}[\{1\}], \dots, \pi_n^{-1}[\{1\}])$  where  $\pi_i: 2^n \rightarrow 2$  is  $i$ -th projection.

Verify the correspondence.

**QUESTIONS?**

# Invariance and Expressiveness

---

## Theorem (Invariance for morphisms)

For all  $F$ -coalgebra morphisms  $f: (X, \gamma) \rightarrow (Y, \delta)$  and all  $\varphi: x \models \varphi$  iff  $f(x) \models \varphi$ .  
 (Equivalently:  $\llbracket \varphi \rrbracket_X = f^{-1}[\llbracket \varphi \rrbracket_Y]$ ).

**Proof** By structural induction on  $\varphi$ . Induction step, modal case, use that (writing  $2^X$  for  $\mathcal{Q}X$ ):

$$\begin{array}{ccc}
 2^X & \xleftarrow{2^f} & 2^Y \\
 2^\gamma \uparrow & & \uparrow 2^\delta \\
 2^{FX} & \xleftarrow{2^{Ff}} & 2^{FY} \\
 \lambda_X^\heartsuit \uparrow & & \uparrow \lambda_Y^\heartsuit \\
 (2^X)^n & \xleftarrow{(2^f)^n} & (2^Y)^n
 \end{array}$$

which says:

for all  $x \in X$ , and all  
 $\langle U_1, \dots, U_n \rangle \in (2^Y)^n$ :

$$\begin{aligned}
 \gamma(x) \in \lambda_X^\heartsuit(f^{-1}[U_1], \dots, f^{-1}[U_n]) \\
 \text{iff} \\
 \delta(f(x)) \in \lambda_Y^\heartsuit(U_1, \dots, U_n)
 \end{aligned}$$

**Corollary:** Truth is invariant for bisimulations and behavioural equivalence:

**Def.** A logic  $\mathcal{L}(\Lambda)$  is **expressive** if  $\mathbb{X}, x \equiv \mathbb{Y}, y$  implies  $\mathbb{X}, x \sim \mathbb{Y}, y$ .

**Def.** A collection  $[\Lambda] = \{\lambda^\heartsuit\}_{\heartsuit \in \Lambda}$  of predicate liftings for  $F$  is **separating** if for all  $t_1 \neq t_2$  in  $F(X)$  there is a  $\heartsuit \in \Lambda$  ( $n$ -ary) and  $(A_1, \dots, A_n) \in (\mathcal{Q}X)^n$  such that  $t_1 \in \lambda_X^\heartsuit(A_1, \dots, A_n)$  and  $t_2 \notin \lambda_X^\heartsuit(A_1, \dots, A_n)$ , or vice versa. [Pattinson'04]

Informally: points in  $F(X)$  can be distinguished with lifted predicates.

**Exercise:** Show that

- $\Lambda = \{\lambda^\square\}$  is separating for  $\mathcal{P}$  where  $\lambda^\square(U)_X = \{V \in \mathcal{P}(X) \mid V \subseteq U\}$ .
- $\Lambda = \{\lambda^{L_q} \mid q \in [0, 1] \cap \mathbb{Q}\}$  is separating for  $\mathcal{D}_\omega$  where  $\lambda_X^{L_q}(U) = \{\mu \in \mathcal{D}_\omega(X) \mid \sum_{x \in U} \mu(x) \geq q\}$ .

# Finitary Functors

Recall: Over *image-finite* LTSs, Hennessy-Milner Logic is expressive.  
(But not over arbitrary LTSs.)

A functor  $F: \mathbf{Set} \rightarrow \mathbf{Set}$  is **finitary** if

$$\text{for all sets } X: \quad F(X) = \bigcup \{(Fi)[F(A)] \mid A \subseteq_{\omega} X, i: A \hookrightarrow X\}$$

If  $F$  preserves inclusions, i.e.,  $X \subseteq Y \Rightarrow F(X) \subseteq F(Y)$ , then  $F$  is finitary if

$$\text{for all sets } X: \quad F(X) = \bigcup \{F(A) \mid A \subseteq_{\omega} X\}$$

(Every  $\mathbf{Set}$ -functor can be modified in a non-essential way to preserve inclusions.)

**Examples:**  $\mathcal{P}_{\omega}$ ,  $\mathcal{D}_{\omega}$ ,  $(-)^A$  for finite  $A$ , bag functor, integer-bag, ...

Finitary functors are closed under composition.

**Theorem 1** If  $F$  is finitary and  $\llbracket \Lambda \rrbracket$  is separating, then  $\mathcal{L}(\Lambda)$  is expressive.

[Pattinson'04][Schröder'08]

## Do Separating $[\![\Lambda]\!]$ Exist?

Yes, but not necessarily of unary predicate liftings.

### Definition

Let  $\alpha \in \mathbb{N} \cup \{\omega\}$ . We call  $\Lambda$   **$\alpha$ -bounded** if for all  $\lambda \in \Lambda$ ,  $\lambda$  has arity  $k < \alpha$ .

### Lemma

There exists an  $\alpha$ -bounded separating set of predicate liftings for  $F$

$\iff \{F(f): F(X) \rightarrow F(2^k) \mid f: X \rightarrow 2^k, k < \alpha\}$  is jointly injective for all  $X$ .

**Exercise:** Prove the Lemma. Hint: Use Yoneda correspondence.

### Theorem 2 [Schröder'08]

If  $F$  is finitary, then there is a separating set of finite-arity predicate liftings for  $F$  (and hence an expressive modal logic).

**Exercise:** Show that  $\mathcal{P}_\omega \circ \mathcal{P}_\omega$  is finitary, but not  $n$ -separating for any  $n < \omega$ . Hence, modalities with unbounded arity needed. Cf. Instantial neighbourhood logic:

$\Box(\varphi_1, \dots, \varphi_n; \psi)$  for all  $n < \omega$ .

**QUESTIONS?**



# Coalgebraic Modal Logic via Relation Lifting

---

# Coalgebraic Modal Logic via Relation Lifting

Introduced by [Moss'00]. Basic idea:

- Language has one “canonical” modality  $\nabla$  that takes elements from  $F(\mathcal{L})$  as argument (where  $\mathcal{L}$  is the set of formulas).
- Semantics of  $\nabla$  via lifting of satisfaction relation  $\models \subseteq X \times \mathcal{L}$ : For  $\alpha \in F(\mathcal{L})$ ,

$$(X, \gamma), x \models \nabla \alpha \quad \text{iff} \quad (\gamma(x), \alpha) \in \overline{F}(\models)$$

where  $\overline{F}$  is the so-called *Barr lifting*:

$$\overline{F}(R) = \{ \langle F(\pi_1)(u), F(\pi_2)(u) \rangle \mid u \in F(R) \} \subseteq F(X_1) \times F(X_2)$$

Remarks:

- To show adequacy,  $F$  needs to preserve weak pullbacks.
- $\nabla$ -logic is always expressive.
- Canonical language, but non-standard.

## Example: $\nabla$ for $\mathcal{P}$ -coalgebras

**Example:** For  $F = \mathcal{P}$ ,  $\overline{\mathcal{P}}$  is also known as the **Egli-Milner lifting**

$$\begin{aligned}\overline{\mathcal{P}}(R) = & \{(U, V) \in \mathcal{P}X \times \mathcal{P}Y \mid \forall u \in U \exists v \in V : (u, v) \in R\} \cap \\ & \{(U, V) \in \mathcal{P}X \times \mathcal{P}Y \mid \forall v \in V \exists u \in U : (u, v) \in R\}\end{aligned}$$

That means, for a set  $\Phi \in \mathcal{P}(\mathcal{L})$  of formulas

$x \models \nabla\Phi$  iff

- all  $R$ -successors of  $x$  satisfy some  $\varphi \in \Phi$ , and
- all  $\varphi \in \Phi$  are satisfied by some  $R$ -successor of  $x$ .

In other words,

$$\nabla\Phi \quad \text{equiv. with} \quad \Box \bigvee_{\varphi \in \Phi} \varphi \wedge \bigwedge_{\varphi \in \Phi} \Diamond\varphi$$

In general,  $\nabla$  can be expressed by predicate liftings and vice versa. [Leal & Kurz]

## Concluding

---

## Coalgebraic modal logic

- uniform framework for studying modal logics for coalgebras.
- encompasses many known logics: basic ML, HML, probabilistic ML, graded ML, ...
- is adequate for reasoning about coalgebraic behaviours: behavioural equivalence implies logical equivalence.
- finitary functors  $F$ , expressive logics exist. (Note: property of functor!)

Framework identifies abstract conditions (e.g. natural transf., separating, finitary,...) that suffice for certain results to hold. Avoids reproving from scratch.

Framework allows exploring limits for logics for coalgebras.

Extensions of basic coalgebraic modal logic:

- with fixpoints: coalgebraic  $\mu$ -calculus (both  $\nabla$  and pred.lifts) [Venema, Kupke, Fontaine, Enqvist, Seifan,...]
- with temporal operators [Cirstea]
- coalgebraic dynamic logic (PDL) [H, Kupke]
- coalgebraic predicate logic [Litak, Sano, Pattinson, Schröder]

# (Uniform) Coalgebraic Theorems

Some coalgebraic generalisations of classic results

- Hennessy-Milner thm (Schröder),
- Van Benthem Characterisation thm:  $\text{CML} = \text{FOL} / \sim$   
(Pattinson, Schröder, Litak, Sano)
- Janin-Walukiewics thm:  $\mu\text{-CML} = \text{MSOL} / \sim$   
(Enqvist, Seifan, Venema)
- Goldblatt-Thomason thm: modal analogue of Birkhoff Variety thm. (Kurz, Rosický)
- Completeness
  - coalgebraic canonical model construction (Pattinson, Schröder),
  - $\nabla$ -logic (Kupke, Kurz, Venema),
  - coalgebraic dynamic logics (H, Kupke)
- Decidability in PSPACE (Schröder, Pattinson)
- Uniform Interpolation (Marti, Enqvist, Seifan, Venema)

# Modal Logic via Dual Adjunctions

Stone-type duality:

$$\begin{array}{ccc}
 \text{Coalg}(F) & \xrightleftharpoons[\text{Uf}]{\overline{Q}} & \text{Alg}(L)^{\text{op}} \\
 \downarrow & & \downarrow \\
 F \curvearrowright \text{Set} & \xrightleftharpoons[\text{Uf}]{Q, \perp} & \text{BA}^{\text{op}} \curvearrowright L^{\text{op}}
 \end{array}$$

Generalise to non-classical base logic and other base categories

$$\begin{array}{ccc}
 \text{Coalg}(F) & \xrightleftharpoons[\overline{S}]{\overline{P}} & \text{Alg}(L)^{\text{op}} \\
 \downarrow & & \downarrow \\
 F \curvearrowright \mathbf{C} & \xrightleftharpoons[S]{P, \perp} & \mathbf{D}^{\text{op}} \curvearrowright L^{\text{op}}
 \end{array}$$

Functor  $L$  specifies type of "modal algebras". Modal semantics given by natural transformation  $\delta: LP \Rightarrow PF$ .

cf. [Kupke et al'04], [Bonsangue & Kurz'05], [Klin'07], [Jacobs & Sokolova'10], [Klin & Rot'16], [de Groot et al.'20] m.m.



**QUESTIONS?**

## References

---

# References

- A. Ballester-Bolinches, M. M. Bonsangue, E. Cosme-Llópez, J. J. M. M. Rutten. The dual equivalence of equations and coequations for automata. *Inf. Comp.* 244 (2015).
- N. Bezhanishvili, M. Bonsangue, H.H. Hansen, D. Kozen, C. Kupke, P. Panangaden, A. Silva. Minimisation in Logical Form. *Samson Abramsky on Logic and Structure in Computer Science and Beyond, Outstanding Contributions to Logic* 25. Springer, 2023.
- N. Bezhanishvili, S. Enqvist, J. De Groot. Duality for Instantial Neighbourhood Logic via Coalgebra. *Proceedings CMCS* 2020.
- M. Bonsangue, A.Kurz, Duality for logics for transition systems, *Proc. FoSSaCS 2005*, *Lect. Notes in Comp. Sci.* 3441 (2005)
- C. Cirstea. A Coalgebraic Approach to Linear-Time Logics, *Proceedings FoSSaCS* (2014).
- C. Cirstea, A. Kurz, D. Pattinson, L. Schröder, Y. Venema. Modal logics are coalgebraic, *Visions of Computer Science*, British Computer Society (2008).
- F. Dahlqvist, T. Schmid. How to write a coequation. *Proceedings of CALCO 2021*, *LiPiCS*.
- S. Enqvist, F. Seifan, Y. Venema. Completeness for coalgebraic fixpoint logic, *Proceedings of Computer Science Logic* (2016)

# References

- S. Enqvist, F. Seifan, Y. Venema. Monadic second-order logic and bisimulation invariance for coalgebras. Proceedings of LICS (2015)
- G. Fontaine, R. Leal, Y. Venema, Automata for coalgebras: An approach using predicate liftings, Proceedings ICALP 2010.
- J. de Groot, H.H. Hansen, A Kurz. Logic-Induced Bisimulations. Advances in Modal Logic (AiML 2020).
- H.P. Gumm, T. Schröder. Types and coalgebraic structure. Algebra Universalis 53 (2005).
- H.H. Hansen, C. Kupke. Weak completeness of coalgebraic dynamic logics. Proceedings of Fixed Points in Computer Science (FICS 2015)
- H.H. Hansen, C. Kupke, E. Pacuit. Neighbourhood structures: Bisimilarity and basic model theory. LMCS 5, 2009.
- H.H. Hansen, W. Poiger. Safety and Strong Completeness via Reducibility for Many-Valued Coalgebraic Dynamic Logics. Proceedings of CALCO 2025.
- B. Jacobs. Introduction to Coalgebra: Towards Mathematics of States and Observation, CUP (2016).
- B. Jacobs, J. Rutten. A tutorial on (co)algebras and (co)induction. EATCS Bulletin 62 (1997).
- B. Jacobs, A. Sokolova. Exemplaric Expressivity of Modal Logics. J. Log. Comp. 20(5), 2010.

# References

- B. Klin. Coalgebraic modal logic beyond sets, Proceedings of Mathematical Foundations of Programming Semantics (MFPS XXIII), ENTCS 173 (2007).
- R. Kojima, C. Cîrstea, K. Muroya, I. Hasuo. Coalgebraic CTL: Fixpoint Characterization and Polynomial-Time Model Checking. Proceedings of CMCS 2024.
- C. Kupke, D. Pattinson, Coalgebraic semantics of modal logics: An overview, TCS 412 (2011).
- C. Kupke, A. Kurz, Y. Venema, Stone coalgebras, TCS 327 (2004).
- C. Kupke, Y. Venema, Coalgebraic automata theory: Basic results, LMCS 4 (2008).
- A.Kurz, J.Rosický. The Goldblatt–Thomason theorem for coalgebras, Proceedings of CALCO 2007.
- R. Leal, A. Kurz. Modalities in the Stone age: A comparison of coalgebraic logics TCS 430 (2012).
- T. Litak, D. Pattinson, K. Sano, L. Schröder. Model Theory and Proof Theory of Coalgebraic Predicate Logic. LMCS 14 (2018)
- J. Marti, Y. Venema. Lax Extensions of Coalgebra Functors. Proceedings CMCS 2012.
- L.S. Moss, Coalgebraic logic. Annals of Pure and Applied Logic 96 (1999)

# References

- D. Pattinson, Coalgebraic modal logic: Soundness, completeness and decidability of local consequence. TCS 309 (2003)
- D. Pattinson, Expressive logics for coalgebras via terminal sequence induction, Notre Dame Journal of Formal Logic 45 (2004).
- J. Rutten, Universal coalgebra: a theory of systems, TCS 249 (1) (2000).
- J. Salamanca, A. Ballester-Bolinches, M. M. Bonsangue, E. Cosme-Llópez, J. J. M. M. Rutten. Regular Varieties of Automata and Coequations. Mathematics of Program Construction 2015.
- J. Salamanca. Coequations and Eilenberg-type Correspondences. PhD thesis. Radboud University, 2018.
- E. Schoen. Relation Lifting and Coalgebraic Logic. Master of Logic Thesis. University of Amsterdam, 2021.
- L. Schröder, Expressivity of coalgebraic modal logic: the limits and beyond, TCS 390 (2008).