


THE BEAUTIFUL GEOMETRY OF APERIODIC TESSELLATIONS

Jesse Bettencourt

April 15, 2015

Supervised by Miroslav Lovric

WHAT IS A TESSELLATION?



Checkerboard Tessellation

WHAT IS A TESSELLATION?


Definition:

A **tessellation** \mathcal{T} of the space \mathbb{E}^n is a countable family of closed sets, T , called tiles:

$$\mathcal{T} = \{T_1, T_2, \dots\}$$

such that

1. \mathcal{T} has **no overlaps**: $\overset{\circ}{T}_i \cap \overset{\circ}{T}_j = \emptyset$ if $i \neq j$
2. \mathcal{T} has **no gaps**: $\bigcup_{i=1}^{\infty} T_i = \mathbb{E}^n$



Definition:

Let $\{T_1, T_2, \dots\}$ be the set of tiles of tessellation \mathcal{T} , partitioned into a set of equivalence classes by **criterion** \mathcal{M} . The set, \mathcal{P} , of representatives of these equivalence classes is called the **protoset** for \mathcal{T} with respect to \mathcal{M} .

Example:

Criterion: $\mathcal{M} = \{\text{Colour of the tile. Only opposite colours may touch.}\}$

Protoset:

$$\mathcal{P} = \left\{ \blacksquare, \square \right\}$$

Definition:


If \mathcal{T} is a tessellation with protoset \mathcal{P} , then we say that \mathcal{P} admits \mathcal{T} .

Example:

We say

$$\mathcal{P} = \{ \blacksquare, \square \}$$


admits




PROTOSETS CAN ADMIT MULTIPLE TESSELLATIONS

$$\mathcal{P} = \{ \blacksquare, \square \}$$

admits both




and




MATCHING RULES CORRESPOND TO DEFORMED PROTOSETS

Edge deformations can force matching rules.

$$\mathcal{P} = \{ \blacksquare, \square \}$$



$$\mathcal{P} = \{ \bullet, \text{deformed square} \}$$



Matching Rules \implies Deformed Protoiset


DESCRIBING TESSELLATIONS

Definition:


A tessellation is said to be **periodic** if it admits translational symmetry in two directions.

Definition:

A tessellation is said to be **non-periodic** if it admits no translational symmetries.



Periodic




Non-Periodic


Checkerboard protoset,

$$\mathcal{P} = \{ \blacksquare, \square \}$$

admits **both** periodic and **non-periodic** tessellations.



Periodic



Non-Periodic

Checkerboard protoset,


$$\mathcal{P} = \{ \blacksquare, \square \}$$

admits **both** periodic and non-periodic tessellations.

Some protosets,

$$\mathcal{P} = \{ \bullet, \diamond \}$$

admit **only** periodic tessellations.



Checkerboard protoset,

$$\mathcal{P} = \{ \blacksquare, \square \}$$

admits **both** periodic and non-periodic tessellations.

Some protosets,

$$\mathcal{P} = \{ \bullet, \square \}$$

admit **only** periodic tessellations.

Are there any protosets,


$$\mathcal{P} = \{ ? \}$$

that admit **only** non-periodic tessellations?

AN APERIODIC TESSELLATION

$$\mathcal{P} = \{ \text{dark blue rhombus}, \text{orange rhombus} \}^*$$

* With matching rules




admits **only non-periodic** tessellations, called Penrose Tessellations.

Cannot be constructed through **local** procedures.

Example:

This arrangement




follows the matching rules.

NON-LOCALITY OF THE PENROSE TESSELLATIONS

Cannot be constructed through **local** procedures.

Example:

This arrangement




*follows the matching rules, but is a **mistake**:*


CONSTRUCTION BY SUBSTITUTION

SUBSTITUTION RULES ON ROBINSON TRIANGLES


First, consider the Rhombs as joined triangles:




SUBSTITUTION RULES ON PENROSE RHOMBS




SUBSTITUTION RULES ON PENROSE RHOMBS




START WITH A SEED: THICK RHOMBUS







APPLY SUBSTITUTION RULES AGAIN




APPLY SUBSTITUTION RULES AGAIN AND AGAIN




APPLY SUBSTITUTION RULES AGAIN AND AGAIN AND AGAIN



APPLY SUBSTITUTION RULES AGAIN AND AGAIN AND AGAIN AND AGAIN



APPLY SUBSTITUTION RULES 10 TIMES




APPLY SUBSTITUTION RULES 11 TIMES




PENROSE TESSELLATION FEATURES

Any pattern with diameter d will begin repeating within $\frac{\sqrt[3]{\phi}}{2}d$ from its perimeter.


DENSITY OF PATTERNS




DENSITY OF PATTERNS



DENSITY OF PATTERNS




DENSITY OF PATTERNS



Penrose Tessellations can admit **five-fold** rotational symmetry.

(Impossible for periodic tessellations)

RATIO OF THICK TO THIN RHOMBS



Questions?



`github.com/jessebett/PenroseTilingThesis`



J H Conway and K M Knowles.
Quasiperiodic tiling in two and three dimensions.
Journal of Physics A: Mathematical and General, 19(17):3645–3653, 1986.



N G De Bruijn.
Algebraic theory of Penrose's non-periodic tilings of the plane. I.
In *Indagationes Mathematicae (Proceedings)*, volume 84, pages 39–52. Elsevier, 1981.



N.G. de Bruijn.
Updown generation of Penrose patterns.
Indagationes Mathematicae, 1(2):201–219, June 1990.



Martin Gardner.
Penrose Tiles to Trapdoor Ciphers: And the Return of Dr Matrix.
Cambridge University Press, 1997.



T. Ogawa and R. Collins.
Chains, Flowers, Rings and Peanuts: Graphical Geodesic Lines and Their Application to Penrose Tiling.
Visual Mathematics, (4), 1999.



R. Penrose.
Pentaplexity A Class of Non-Periodic Tilings of the Plane.
The Mathematical Intelligencer, 2(1):32–37, March 1979.



Marjorie Senechal.
Quasicrystals and geometry.
CUP Archive, 1996.