

	<b>Table 1 Prevalence of Hypertention and Sociodemographic Characteristics and Risk Factors, California, 2007</b>		
	<b>Charaterisitc</b>	<b>Unweighted Sample Size</b>	<b>Prevalence of Bringe Drinking</b>
	Total Adults	51048	
	<b>Age</b>		
	18-25 (ref.)	3181	43.60%
	26-34	4632	41.11%
	35-49	12801	30.60%
	50+	30434	15.35%
	<b>Gender</b>		
	Male (ref.)	20507	31.48%
	Female	30541	17.76%
	<b>Race/ethnicity</b>		
	Non-Hipanic White (ref.)	33329	24.02%
	Hispanic	9077	26.97%
	Non-Hipanic Black	2398	17.47%
	Non-Hipanic Asian	4340	12.53%
	Non-Hipanic American Indians/Alaska	421	25.18%
	Non-Hipanic Other	1483	24.07%
	<b>Education status</b>		
	< High-school	4963	16.78%
	High-school graduate (ref.)	11380	24.81%
	Some college	14456	24.07%

	College or more	20249	23.42%
	<b>Poverty level</b>		
	<100% FPL (ref.)	7109	19.16%
	100%-199% FPL	8351	18.86%
	200%-399% FPL	12698	22.52%
	≥400% FPL	22890	26.57%
	<b>Employment level</b>		
	Full-time (ref.)	25450	31.39%
	Part-time	3951	22.20%
	Employed but not work	169	22.49%
	Unemployed and look for work	1232	30.52%
	Unemployed but not look for work	20246	39.46%
	<b>Marital status</b>		
	Married (ref.)	26192	22.61%
	Never married	17080	15.49%
	Other	7776	42.61%
	<b>Body weight status</b>		
	Underweight	1055	13.36%
	Normal (ref.)	21162	22.53%
	Overweight	17560	25.34%
	Obesity	11271	22.37%
	<b>Smoking Status</b>		
	Ever smoking	22124	29.00%

	No smoking	28924	18.88%
--	------------	-------	--------