

STATISTICS WORKSHEET-3

Q1 to Q9 have only one correct answer. Choose the correct option to answer your question.

1. Which of the following is the correct formula for total variation?
- a) Total Variation = Residual Variation – Regression Variation
 - b) Total Variation = Residual Variation + Regression Variation
 - c) Total Variation = Residual Variation * Regression Variation
 - d) All of the mentioned

Answer: Option B – Total Variation = Residual Variation + Regression Variation

2. Collection of exchangeable binary outcomes for the same covariate data are called_____outcomes.
- a) random
 - b) direct
 - c) binomial
 - d) none of the mentioned

Answer: Option C- binomial

3. How many outcomes are possible with Bernoulli trial?
- a) 2
 - b) 3
 - c) 4
 - d) None of the mentioned

Answer: Option A - 2

4. If H_0 is true and we reject it is called
- a) Type-I error
 - b) Type-II error
 - c) Standard error
 - d) Sampling error

Answer: Option A – Type-I error

5. Level of significance is also called:
- a) Power of the test
 - b) Size of the test
 - c) Level of confidence
 - d) Confidence coefficient

Answer: Option B – Size of the test

6. The chance of rejecting a true hypothesis decreases when sample size is:
- a) Decrease
 - b) Increase
 - c) Both of them
 - d) None

Answer: Option D – None

7. Which of the following testing is concerned with making decisions using data?
- a) Probability
 - b) Hypothesis
 - c) Causal
 - d) None of the mentioned

Answer: Option B - Hypothesis

8. What is the purpose of multiple testing in statistical inference?
- a) Minimize errors
 - b) Minimize false positives
 - c) Minimize false negatives
 - d) All of the mentioned

Answer: Option D – All of the mentioned

9. Normalized data are centred at ____ and have units equal to standard deviations of the original data
- a) 0
 - b) 5
 - c) 1
 - d) 10

Answer: Option A - 0

Q10 and Q15 are subjective answer type questions, Answer them in your own words briefly.

10. What Is Bayes' Theorem?

Answer: In statistics and probability theory, the Bayes' theorem (also known as the Bayes' rule) is a mathematical formula used to determine the conditional probability of events. Essentially, the Bayes' theorem describes the probability of an event based on prior knowledge of the conditions that might be relevant to the event.

11. What is z-score?

Answer: A measure of how many standard deviations below or above the population mean a raw is called z score. It will be positive if the value lies above the mean and negative if it lies below the mean. It is also known as standard score. It indicates how many standard deviations an entity is, from the mean. In order to use a z-score, the mean μ and also the population standard deviation σ should be known. A z score helps to calculate the probability of a score occurring within a standard normal distribution. It also enables us to compare two scores that are from different samples. A table for the values of ϕ , indicating the values of the cumulative distribution function of the normal distribution is termed as a z score table.

12. What is t-test?

Answer: A t-test is a statistical test that is used to compare the means of two groups. It is often used in hypothesis testing to determine whether a process or treatment actually has an effect on the population of interest, or whether two groups are different from one another.

13. What is percentile?


Answer: In statistics, a percentile is a term that describes how a score compares to other scores from the same set. While there is no universal definition of percentile, it is commonly expressed as the percentage of values in a set of data scores that fall below a given value.

14. What is ANOVA?

Answer: Analysis of variance (ANOVA) is an analysis tool used in statistics that splits an observed aggregate variability found inside a data set into two parts: systematic factors and random factors. The systematic factors have a statistical influence on the given data set, while the random factors do not. Analysts use the ANOVA test to determine the influence that independent variables have on the dependent variable in a regression study.

15. How can ANOVA help?

Answer: The one-way ANOVA can help you know whether or not there are significant differences between the means of your independent variables (such as the first example: age, sex, income). When you understand how each independent variable's mean is different from the others, you can begin to understand which of them has a connection to your dependent variable (landing page clicks), and begin to learn what is driving that behavior.



FLIP ROBO