

Properties of Expectations

Multivariate Normal Random Variables [Ross S7.8]

Definition of Multivariate Normal

Let Z_1, Z_2, \dots, Z_n be independent $\sim \mathcal{N}(0, 1)$.

Then, define X_1, X_2, \dots, X_m by

$$\begin{aligned} X_1 &= a_{11}Z_1 + \dots + a_{1n}Z_n + \mu_1 \\ X_2 &= a_{21}Z_1 + \dots + a_{2n}Z_n + \mu_2 \\ &\vdots \qquad \qquad \qquad \vdots \\ X_m &= a_{m1}Z_1 + \dots + a_{mn}Z_n + \mu_m \end{aligned}$$

We say that X_1, \dots, X_m are **multivariate normal (or jointly Gaussian)**.

We can write this in vector form as $\mathbf{X} = \mathbf{AZ} + \boldsymbol{\mu}$:

$$\underbrace{\begin{pmatrix} X_1 \\ X_2 \\ \vdots \\ X_m \end{pmatrix}}_{\mathbf{X}} = \underbrace{\begin{pmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{pmatrix}}_A \underbrace{\begin{pmatrix} Z_1 \\ Z_2 \\ \vdots \\ Z_n \end{pmatrix}}_{\mathbf{Z}} + \underbrace{\begin{pmatrix} \mu_1 \\ \mu_2 \\ \vdots \\ \mu_m \end{pmatrix}}_{\boldsymbol{\mu}}$$

Now, let B be a $k \times m$ matrix, and $\boldsymbol{\nu}$ a column vector of length k . Then

$$\begin{aligned} \mathbf{Y} &= \mathbf{BX} + \boldsymbol{\nu} \\ &= (\mathbf{BA})\mathbf{Z} + (\mathbf{B}\boldsymbol{\mu} + \boldsymbol{\nu}) \end{aligned}$$

So \mathbf{Y} is multivariate Gaussian too: an affine transformation of a multivariate Gaussian is again multivariate Gaussian!

Marginal Distribution of X_i

Since X_i is a sum of independent Gaussian random variables

$\rightarrow X_i$ is Gaussian [Proposition 26.1 in Notes #26]

Also:

$$\begin{aligned} E[X_i] &= E[a_{i1}Z_1 + \dots + a_{in}Z_n + \mu_i] \\ &= a_{i1}E[Z_1] + \dots + a_{in}E[Z_n] + \mu_i \\ &= \mu_i \end{aligned}$$

$$\begin{aligned} Var[X_i] &= Var[a_{i1}Z_1 + \dots + a_{in}Z_n + \mu_i] \\ &= Var[a_{i1}Z_1 + \dots + a_{in}Z_n] \\ &= a_{i1}^2 Var[Z_1] + \dots + a_{in}^2 Var[Z_n] \\ &= a_{i1}^2 + \dots + a_{in}^2 \end{aligned}$$

A single Gaussian random variable U is uniquely specified by:

- its mean $E[U]$
- and its variance $Var[U]$.

Similarly:

The joint distribution of a multivariate Gaussian (normal) depends only on:

- the means $E[X_i]$ for $i = 1, \dots, m$
- and the co-variances $Cov[X_i, X_j]$ for $i = 1, \dots, m$ and $j = 1, \dots, m$

What happened to $Var[X_1], Var[X_2]$, etc?

$Var[X_1] = Cov[X_1, X_1]$, so these are in the second bullet.

=====

Common Notation

For random variables X_1, \dots, X_m , it is common to define:

$$\mathbf{X} = \begin{pmatrix} X_1 \\ X_2 \\ \vdots \\ X_m \end{pmatrix} \qquad \text{[random vector]}$$

$$\boldsymbol{\mu} = E[\mathbf{X}] = \begin{pmatrix} E[X_1] \\ E[X_2] \\ \vdots \\ E[X_m] \end{pmatrix} \qquad \text{[mean vector]}$$

$$\begin{aligned} \Sigma &= E[(\mathbf{X} - \boldsymbol{\mu})(\mathbf{X} - \boldsymbol{\mu})^T] \qquad \text{[covariance matrix]} \\ &= E\left[\begin{pmatrix} (X_1 - \mu_1)(X_1 - \mu_1) & (X_1 - \mu_1)(X_2 - \mu_2) & \cdots & (X_1 - \mu_1)(X_m - \mu_m) \\ (X_2 - \mu_2)(X_1 - \mu_1) & (X_2 - \mu_2)(X_2 - \mu_2) & \cdots & (X_2 - \mu_2)(X_m - \mu_m) \\ \vdots & \vdots & \ddots & \vdots \\ (X_m - \mu_m)(X_1 - \mu_1) & (X_m - \mu_m)(X_2 - \mu_2) & \cdots & (X_m - \mu_m)(X_m - \mu_m) \end{pmatrix}\right] \\ &= \begin{pmatrix} Cov[X_1, X_1] & Cov[X_1, X_2] & \cdots & Cov[X_1, X_m] \\ Cov[X_2, X_1] & Cov[X_2, X_2] & \cdots & Cov[X_2, X_m] \\ \vdots & \vdots & \ddots & \vdots \\ Cov[X_m, X_1] & Cov[X_m, X_2] & \cdots & Cov[X_m, X_m] \end{pmatrix} \end{aligned}$$

Also, note that

$$\begin{aligned} \Sigma &= E[(\mathbf{X} - \boldsymbol{\mu})(\mathbf{X} - \boldsymbol{\mu})^T] \\ &= E[\mathbf{X}\mathbf{X}^T - \boldsymbol{\mu}\mathbf{X}^T - \mathbf{X}\boldsymbol{\mu}^T + \boldsymbol{\mu}\boldsymbol{\mu}^T] \\ &= E[\mathbf{X}\mathbf{X}^T] - E[\boldsymbol{\mu}\mathbf{X}^T] - E[\mathbf{X}\boldsymbol{\mu}^T] + E[\boldsymbol{\mu}\boldsymbol{\mu}^T] \\ &= E[\mathbf{X}\mathbf{X}^T] - \boldsymbol{\mu}E[\mathbf{X}^T] - E[\mathbf{X}]\boldsymbol{\mu}^T + \boldsymbol{\mu}\boldsymbol{\mu}^T \\ &= E[\mathbf{X}\mathbf{X}^T] - \boldsymbol{\mu}\boldsymbol{\mu}^T - \boldsymbol{\mu}\boldsymbol{\mu}^T + \boldsymbol{\mu}\boldsymbol{\mu}^T \\ &= E[\mathbf{X}\mathbf{X}^T] - \boldsymbol{\mu}\boldsymbol{\mu}^T \end{aligned}$$

If X_1, \dots, X_m are jointly Gaussian with $\boldsymbol{\mu}$ and Σ , we write $\mathbf{X} \sim \mathcal{N}(\boldsymbol{\mu}, \Sigma)$.

It can be shown that if Σ is invertible, then

$$f_{\mathbf{X}}(\mathbf{x}) = \frac{1}{\sqrt{(2\pi)^m |\Sigma|}} e^{-\frac{1}{2}(\mathbf{x} - \boldsymbol{\mu})\Sigma^{-1}(\mathbf{x} - \boldsymbol{\mu})}$$

Note: as expected, this depends only on $\boldsymbol{\mu}$ and Σ .

=====

Covariance Matrix

Say Z_1, \dots, Z_n are independent $\sim \mathcal{N}(0, 1)$. Then

$$\begin{aligned} \boldsymbol{\mu}_Z &= E[\mathbf{Z}] = \mathbf{0} \\ \Sigma_Z &= \begin{pmatrix} Cov[Z_1, Z_1] & Cov[Z_1, Z_2] & \cdots & Cov[Z_1, Z_n] \\ Cov[Z_2, Z_1] & Cov[Z_2, Z_2] & \cdots & Cov[Z_2, Z_n] \\ \vdots & \vdots & \ddots & \vdots \\ Cov[Z_n, Z_1] & Cov[Z_n, Z_2] & \cdots & Cov[Z_n, Z_n] \end{pmatrix} \\ &= \begin{pmatrix} 1 & 0 & \cdots & 0 \\ 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 1 \end{pmatrix} = I \end{aligned}$$

Effect of Affine transformation on Covariance Matrix

Let \mathbf{X} have mean $\boldsymbol{\mu}_X$ and co-variance matrix Σ_X .

Let B be a matrix, and $\boldsymbol{\nu}$ a column vector.

Let $\mathbf{Y} = \mathbf{BX} + \boldsymbol{\nu}$. Then

$$\boldsymbol{\mu}_Y = E[\mathbf{Y}] = E[\mathbf{BX} + \boldsymbol{\nu}] = BE[\mathbf{X}] + \boldsymbol{\nu} = B\boldsymbol{\mu}_X + \boldsymbol{\nu}$$

$$\begin{aligned} \Sigma_Y &= E[\mathbf{Y}\mathbf{Y}^T] - \boldsymbol{\mu}_Y\boldsymbol{\mu}_Y^T \\ &= E[(\mathbf{BX} + \boldsymbol{\nu})(\mathbf{BX} + \boldsymbol{\nu})^T] - \boldsymbol{\mu}_Y\boldsymbol{\mu}_Y^T \\ &= E[\mathbf{B}\mathbf{X}\mathbf{X}^T\mathbf{B}^T + \mathbf{B}\mathbf{X}\boldsymbol{\nu}^T + \boldsymbol{\nu}\mathbf{X}^T\mathbf{B}^T + \boldsymbol{\nu}\boldsymbol{\nu}^T] - \boldsymbol{\mu}_Y\boldsymbol{\mu}_Y^T \\ &= BE[\mathbf{X}\mathbf{X}^T]\mathbf{B}^T + BE[\mathbf{X}]\boldsymbol{\nu}^T + \boldsymbol{\nu}E[\mathbf{X}^T]\mathbf{B}^T + \boldsymbol{\nu}\boldsymbol{\nu}^T - \boldsymbol{\mu}_Y\boldsymbol{\mu}_Y^T \\ &= BE[\mathbf{X}\mathbf{X}^T]\mathbf{B}^T + B\boldsymbol{\mu}_X\boldsymbol{\nu}^T + \boldsymbol{\nu}\boldsymbol{\mu}_X^T\mathbf{B}^T + \boldsymbol{\nu}\boldsymbol{\nu}^T - \boldsymbol{\mu}_Y\boldsymbol{\mu}_Y^T \\ &= BE[\mathbf{X}\mathbf{X}^T]\mathbf{B}^T + B\boldsymbol{\mu}_X\boldsymbol{\nu}^T + \boldsymbol{\nu}\boldsymbol{\mu}_X^T\mathbf{B}^T + \boldsymbol{\nu}\boldsymbol{\nu}^T - (B\boldsymbol{\mu}_X + \boldsymbol{\nu})(B\boldsymbol{\mu}_X + \boldsymbol{\nu})^T \\ &= BE[\mathbf{X}\mathbf{X}^T]\mathbf{B}^T + B\boldsymbol{\mu}_X\boldsymbol{\nu}^T + \boldsymbol{\nu}\boldsymbol{\mu}_X^T\mathbf{B}^T + \boldsymbol{\nu}\boldsymbol{\nu}^T \\ &\quad - (B\boldsymbol{\mu}_X\boldsymbol{\mu}_X^T\mathbf{B}^T + B\boldsymbol{\mu}_X\boldsymbol{\nu}^T + \boldsymbol{\nu}\boldsymbol{\mu}_X^T\mathbf{B}^T + \boldsymbol{\nu}\boldsymbol{\nu}^T) \\ &= BE[\mathbf{X}\mathbf{X}^T]\mathbf{B}^T - B\boldsymbol{\mu}_X\boldsymbol{\mu}_X^T\mathbf{B}^T \\ &= B(E[\mathbf{X}\mathbf{X}^T] - \boldsymbol{\mu}_X\boldsymbol{\mu}_X^T)\mathbf{B}^T \\ &= B\Sigma_X\mathbf{B}^T \end{aligned}$$

=====

Not all square matrices can be covariance matrices.

Below, is a general condition.

Proposition 37.1 *a) A covariance matrix Σ is i) symmetric and ii) positive semi-definite.
b) Any matrix Σ that is symmetric and positive semi-definite is the covariance matrix of $\mathbf{X} = \mathbf{AZ} + \boldsymbol{\mu}$ for some choice of matrix A .*

Why?

$$\begin{aligned} i) \quad \Sigma^T &= E[(\mathbf{X} - \boldsymbol{\mu})(\mathbf{X} - \boldsymbol{\mu})^T]^T \\ &= E[(\mathbf{X} - \boldsymbol{\mu})(\mathbf{X} - \boldsymbol{\mu})^T]^T \\ &= E[(\mathbf{X} - \boldsymbol{\mu}^T)^T(\mathbf{X} - \boldsymbol{\mu})^T] \\ &= E[(\mathbf{X} - \boldsymbol{\mu})(\mathbf{X} - \boldsymbol{\mu})^T] \\ &= \Sigma \\ ii) \quad \mathbf{v}^T\Sigma\mathbf{v} &= \mathbf{v}^TE[(\mathbf{X} - \boldsymbol{\mu})(\mathbf{X} - \boldsymbol{\mu})^T]\mathbf{v} \\ &= E[\mathbf{v}^T(\mathbf{X} - \boldsymbol{\mu})(\mathbf{X} - \boldsymbol{\mu})^T\mathbf{v}] \\ &= E[|(\mathbf{X} - \boldsymbol{\mu})^T\mathbf{v}|^2] \\ &\geq 0 \end{aligned}$$

b) Since Σ is symmetric, it can be diagonalized as $\Sigma = \mathbf{UDU}^T$ where D is diagonal.

The diagonal entries of D are ≥ 0 since it Σ is positive semi-definite.

Then $\Sigma = \mathbf{UD}^{1/2}\mathbf{D}^{1/2}\mathbf{U}^T$.

Let $A = \mathbf{UD}^{1/2}$.

Then

$$\begin{aligned} \Sigma_X &= A\Sigma_ZA^T \\ &= AA^T \\ &= \mathbf{UD}^{1/2}(\mathbf{UD}^{1/2})^T \\ &= \mathbf{UD}^{1/2}(\mathbf{D}^{1/2})^T\mathbf{U}^T \\ &= \mathbf{UD}^{1/2}\mathbf{D}^{1/2}\mathbf{U}^T \\ &= \Sigma \end{aligned}$$