

Nest detection and habitat mapping of burrow-nesting seabirds using aerial drone based imagery

Jiménez López, Jesús[1]; García Perez, Agapito[2]¹

¹Department of Arcane Biology, University of Miskatonic

²Department of Astrobiology, University of Lebrija

May 17, 2022

Abstract

See https://github.com/jesusjl/proyecto_final

In the last decade, drones (also known as unmanned aerial systems, remotely piloted aircraft systems, RPAS, UAS, UAV) have been the subject of a growing interest in both the civilian and scientific sphere. Drones offer a relatively risk-free and low-cost manner to rapidly and systematically observe natural phenomena at high spatio-temporal resolution. In this study, a drone was used to capture aerial footage to count North Atlantic White-faced Storm-petrel ground nests and describe habitat-specific environmental conditions.

Keywords: drones, UAV, remote sensing, wildlife monitoring, conservation, island ecology, *Pelagodroma marina*, burrow nesting seabirds, Selvagem Grande; Macaronesia, Pitingo

1 Introduction

The North Atlantic White-faced Storm-petrel (*Pelagodroma marina hypoleuca*) digs burrow nests in extremely frangible sandy soils, typically forming dense

colonies that are often confined to remote and inhabited islands and islets. Conservation and management efforts require regular monitoring of their nests to assess their breeding status and success, as well as possible impacts of anthropogenic pressures. However, fieldwork and censuses are infrequent and with considerable risk of nest trampling. In this context, drone-based aerial imagery surveys can be a non-intrusive, reliable, and effective method of counting nests where close inspection is impractical or constitutes an unacceptable risk of nest disruption and habitat disturbance [3].

In this study, a drone was used to collect aerial photographs to assess the feasibility of detecting and counting nests in the imagery and describe habitat-specific environmental conditions. An ortho-photomosaic of the target nesting area was generated, confirming that in stark contrast to sandy soils and surrounding patches of vegetation, burrows were explicitly visible. Hence, burrows were exhaustively counted by visual inspection and compared with semi-automatic detection methods. In addition, object-based image analysis (OBIA) was applied to the ortho-photomosaic to produce a detailed map of vegetation and landforms. Ecological and environmental conditions driving the spatial distribution of the nests were explored to unravel apparent clustering patterns. We discuss the potential and limitations in the use of drones to detect burrowing nests and mapping breeding areas in remote unpopulated islands, and provide guidelines for improved data acquisition to monitor Important Bird and Biodiversity Area (IBA) over time.

Our study case illustrates that regular scientific ecological information from conspicuous burrow-nesting seabirds can be acquired using very high-resolution aerial images and different processing analytical tools, with minimal disturbance and reduced risks to the nests integrity, contributing therefore to the conservation of North Atlantic White-faced Storm-petrel.

2 Materials and Methods

Located in the North Atlantic Ocean, the Portuguese Archipelago of Selvagens ($30^{\circ}8'46.68''\text{N}$ $15^{\circ}51'51.8''\text{W}$) are the smaller group of islands in the Macaronesian region, 280 kilometres south of Madeira, and 165 kilometres north of the Canary Islands. The archipelago includes two major cluster of islands and islets (Selvagem Grande and Selvagem Pequena) where the White-faced Storm-petrel are known to burrow their nests. In July 2019, we conducted a drone survey in the northernmost colony of the two main breed-

ing areas in Selvagem Grande. Since flights were performed in the daylight, bird activity was minimum and no disturbance from drone was observed.

Burrow nests spots on the orthophoto were systematically digitized as spatial points shapefiles with the assistance of a vector grid of equal squared cells (QGIS). Following, we compare the manual counted nest dataset with semi-automatic detection using object based image analysis (OBIA) techniques (eCognition Essential). For this purpose, the vector shapefile was split in training and validation sets and respectively used for the image segmentation / classification and accuracy assessment phases.

Similar, an habitat map of the study area was produced using OBIA. The very high resolution mosaic permit to clearly differentiate main land cover classes. For this reason, training data was also

The habitat map was used to generate landscape metrics. We used R

References

- [1] Ana Ferreira, Sónia Seixas, and João Carlos Marques. “bottom-up management approach to coastal marine protected areas in portugal”. 118:275–281.
- [2] Miguel Pessanha Pais, Sofia Henriques, Marisa Isabel Batista, Carlos Rius Errando, Henrique Nogueira Cabral, Paula Chaínho, Maria Ana Dionísio, Inês Afonso, Frederico Almada, Pedro Duarte-Coelho, Inês Carneiro, Cristina Lima, Sara Francisco, Ana Pereira, and Joana Robalo. AquaSig 3: Recolha de informação georreferenciada sobre a diversidade e abundância de macrofauna e flora no espaço marítimo de cascais.
- [3] Daniele Ventura, Andrea Bonifazi, Maria Flavia Gravina, Andrea Beluscio, and Giandomenico Ardizzone. Mapping and classification of ecologically sensitive marine habitats using unmanned aerial vehicle (UAV) imagery and object-based image analysis (OBIA). 10(9):1331.
- [4] M.J. Westoby, J. Brasington, N.F. Glasser, M.J. Hambrey, and J.M. Reynolds. ‘structure-from-motion’ photogrammetry: A low-cost, effective tool for geoscience applications. 179:300–314.