

# Mathematical Statistics I

## Chapter 3: Joint Distributions

---

Jesse Wheeler

1. Introduction
2. Discrete Random Variables
3. Continuous Random Variables
4. Independent Random Variables
5. Conditional Distributions
  - Discrete random variables
  - Continuous Random Variables
6. Functions of Jointly Distributed Random Variables

# Introduction

---

# Introduction

- This material is based on the textbook by Rice (2007, Chapter 3).
- Our goal is to better understand the joint probability structure of more than one random variable, defined on the same sample space.
- One reason that studying joint probabilities is an important topic is that it enables us to use what we know about one variable to study another.

- Just like the univariate case, the joint behavior of two random variables,  $X$  and  $Y$ , is determined by the cumulative distribution function

$$F(x, y) = P(X \leq x, Y \leq y).$$

- This is true for both discrete and continuous random variables.
- The any set  $A \subset \mathbb{R}^2$ , the joint cdf can give  $P((X, Y) \in A)$ .

## Joint cdf II

- For example, let  $A$  be the rectangle defined by  $x_1 < X < x_2$ , and  $y_1 < Y < y_2$ . (It helps to draw a picture...)
- $F(x_2, y_2)$  gives  $P(X < x_2, Y < y_2)$ , an area that is too big, so we subtract off pieces
  - $F(x_2, y_1) = P(X < x_2, Y < y_1)$  (we already have the area  $X < x_2$ , but now subtract away the area  $Y < y_1$ ).
  - $F(x_1, y_2) = P(X < x_1, Y < y_2)$  (Now subtracting the area  $X < x_1$ )
  - We have “double subtracted” the area  $\{X < x_1, Y < y_1\}$ , so we add it back.

$$P((X, Y) \in A) = F(x_2, y_2) - F(x_2, y_1) - F(x_1, y_2) + F(x_1, y_1).$$

## Joint cdf III

- The definition also applies to more than two random variables.
- Let  $X_1, \dots, X_n$  be jointly distributed random variables defined on the same sample space. Then

$$F(x_1, x_2, \dots, x_n) = P(X_1 \leq x_1, X_2 \leq x_2, \dots, X_n \leq x_n).$$

- Like the univariate case, we can also define the pmf and pdf of jointly distributed random variables as well.

# Discrete Random Variables

---



# Discrete Random Variables

## Definition: Joint pmf

Let  $X$  and  $Y$  be discrete random variables defined on the same sample space, and take on values  $x_1, x_2, \dots$  and  $y_1, y_2, \dots$ , respectively. The **joint pmf** (or joint frequency function), is

$$p(x_i, y_j) = P(X = x_i, Y = y_j).$$

- For discrete RVs, it's often useful to describe the joint pmf as a frequency table.

## Discrete Random Variables II

- Suppose a fair coin is tossed 3 times. Let  $X$  denote the number of heads on the first toss, and  $Y$  the total number of heads.
- The sample space is

$$\Omega = \{hhh, hht, hth, thh, htt, tht, tth, ttt\}.$$

- The joint pmf can be expressed as the frequency table below (Table 1).

## Discrete Random Variables III

|     | $y$           |               |               |               |
|-----|---------------|---------------|---------------|---------------|
| $x$ | 0             | 1             | 2             | 3             |
| 0   | $\frac{1}{8}$ | $\frac{2}{8}$ | $\frac{1}{8}$ | 0             |
| 1   | 0             | $\frac{1}{8}$ | $\frac{2}{8}$ | $\frac{1}{8}$ |

**Table 1:** Frequency table for  $X$  and  $Y$ , flipping a fair coin three times.

- Note that the probabilities in Table 1 sum to one.
- Using the probability laws we have already learned, we can calculate **marginal** probabilities.

$$\begin{aligned}p_Y(0) &= P(Y = 0) \\&= P(Y = 0, X = 0) + P(Y = 0, X = 1) \\&= \frac{1}{8} + 0 = \frac{1}{8}\end{aligned}$$

$$\begin{aligned}p_Y(1) &= P(Y = 1) \\&= P(Y = 1, X = 0) + P(Y = 1, X = 1) \\&= \frac{2}{8} + \frac{1}{8} = \frac{3}{8}.\end{aligned}$$

# Discrete Random Variables V

- In general, to find the frequency function for  $Y$  and  $X$ , we just need to sum the appropriate columns or rows, respectively.
- $p_X(x) = \sum_i P(x, y_i)$  and  $p_Y(y) = \sum_j P(x_j, y)$ .
- The case with multiple random variables is similar:

$$p_{X_i}(x_i) = \sum_{x_j: j \neq i} p(x_1, x_2, \dots, x_n).$$

- We can also get marginal frequencies for more than one variable:

$$p_{X_i X_j}(x_i, x_j) = \sum_{x_k: k \notin \{i, j\}} p(x_1, x_2, \dots, x_n).$$

## Example: Multinomial Distribution

- The **multinomial** distribution is a generalization of the binomial distribution.
- Suppose there are  $n$  independent trials, each with  $r$  possible outcomes, with probabilities  $p_1, p_2, \dots, p_r$ , respectively.
- Let  $N_i$  be the total number of outcomes of type  $i$  in the  $n$  trials, with  $i \in \{1, 2, \dots, r\}$ .
- The probability of any particular sequence  $(N_1, N_2, \dots, N_r) = (n_1, n_2, \dots, n_r)$  is

$$p_1^{n_1} p_2^{n_2} \cdots p_r^{n_r}$$

## Example: Multinomial Distribution II

- The total number of ways to do this was an identity from Chapter 1 (Proposition 1.3):

$$\binom{n}{n_1 \cdots n_r}.$$

- Combining this gives us the pmf of the multinomial distribution:

### Multinomial Distribution

Let  $N_1, N_2, \dots, N_r$  be random variables that follow a multinomial distribution with parameters  $N$  and  $(p_1, \dots, p_r)$ . The joint pmf is

$$p(n_1, n_2, \dots, n_r) = \binom{n}{n_1 \cdots n_r} p_1^{n_1} p_2^{n_2} \cdots p_r^{n_r}$$

## Example: Multinomial Distribution III

- The marginal distribution for any  $N_i$  can be found by summing the joint frequency function over the other  $n_j$ .
- While possible, this is a non-trivial algebraic exercise.
- The simple alternative is to reframe the problem: Let  $N_i$  be the number of successes in  $n$  trials, and  $\tilde{N}_i = \sum_{j \neq i} N_j$  be the number of failures. The probability of success is still  $p_i$ , leaving the probability of failure to be  $1 - p_i$ .
- Thus, we see that the marginal distribution for  $N_i$  must follow a binomial distribution:

$$\begin{aligned} p_{N_i}(n_i) &= \sum_{n_j: j \neq i} \binom{n}{n_1 \dots n_r} p_1^{n_1} p_2^{n_2} \dots p_r^{n_r} \\ &= \binom{n}{n_i} p_i^{n_i} (1 - p_i)^{n - n_i} \end{aligned}$$



# Continuous Random Variables

---

# Continuous Random Variables

- Let  $X, Y$  be continuous random variables with joint cdf  $F(x, y)$ .
- Their **joint density function** is a piecewise continuous function of two variables,  $f(x, y)$ .
- A few properties:
  - $f(x, y) \geq 0$  for all  $(x, y) \in \mathbb{R}$  (or the support).
  - $\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f(x, y) dx dy = 1$ .
  - For any “measureable set”  $A \subset \mathbb{R}^2$ ,  
$$P((X, Y) \in A) = \int \int_A f(x, y) dx dy$$
  - In particular,  $F(x, y) = \int_{-\infty}^x \int_{-\infty}^y f(u, v) du dv$ .

## Continuous Random Variables II

- From the fundamental theorem of multivariable calculus, it follows that

$$f(x, y) = \frac{\partial^2}{\partial x \partial y} F(x, y),$$

wherever the derivative is defined.

## Continuous Random Variables III

### Finding joint probabilities

Let  $X, Y$  be jointly defined RVs with pdf

$$f(x, y) = \frac{12}{7}(x^2 + xy), \quad 0 \leq x \leq 1, \quad 0 \leq y \leq 1.$$

Find  $P(X > y)$ .

*Solution:*

## Marginal cdf

The **marginal cdf** of  $X$ , denoted  $F_X$ , is

$$\begin{aligned}F_X(x) &= P(X \leq x) \\&= P(X \leq x \cap Y \in \mathbb{R}) = P(X \leq x \cap Y < \infty) \\&= \lim_{y \rightarrow \infty} F(x, y) \\&= \int_{-\infty}^x \int_{-\infty}^{\infty} f(u, y) dy du.\end{aligned}$$

By taking the derivative of both sides of the equation, we get the **marginal density** of  $X$ :

$$f_X(x) = F'_X(x) = \int_{-\infty}^{\infty} f(x, y) dy.$$

### Calculating Marginal Densities

Using the same joint distribution as the previous example, find the marginal density of  $X$ :

$$\begin{aligned}f_X(x) &= \int_Y f(x, y) dy \\&= \frac{12}{7} \int_0^1 (x^2 + xy) dy \\&= \frac{12}{7} \left( x^2 y + \frac{x}{2} y^2 \right) \Big|_0^1 \\&= \frac{12}{7} \left( x^2 + \frac{x}{2} \right)\end{aligned}$$

## More than two random variables

- For several jointly continuous random variables, we can make the obvious generalizations.
- That is, to find the *marginal* densities, we need to “marginalize-” or “integrate-” out the **nusaince** variables.
- This means integrating out any combination of variables that we want.
- Example: Let  $X$ ,  $Y$ , and  $Z$  be jointly continuous RVs with pdf  $f(x, y, z)$ . Then the two-dimensional marginal distribution of  $X$  and  $Z$  is:

$$f_{XZ}(x, z) = \int_{-\infty}^{\infty} f(x, y, z) dy.$$

## Example: constructing bivariate cdfs

- Suppose that  $F(x)$  and  $G(y)$  are cdfs for random variables  $X$  and  $Y$ , resp.
- It can be shown that the following function,  $H(x, y)$ , is always a bivariate cdf for all  $-1 \leq \alpha \leq 1$ :

$$H(x, y) = F(x)G(y)\left(1 + \alpha(1 - F(x))(1 - G(y))\right).$$

- Because  $\lim_{x \rightarrow \infty} F(x) = \lim_{y \rightarrow \infty} G(y) = 1$ , the marginal distributions are:

$$\lim_{y \rightarrow \infty} H(x, y) = F(x)$$

$$\lim_{x \rightarrow \infty} H(x, y) = G(y)$$



## Example: constructing bivariate cdfs II

- Thus, we can use this approach to build an infinite number of bivariate distributions that have a particular marginal distribution.

## Example: constructing bivariate cdfs III

- One important example is when the marginal distributions are uniformly distributed.
- Let  $F(x) = x, 0 \leq x \leq 1$ , and  $G(y) = y, 0 \leq y \leq 1$ .
- By selecting  $\alpha = -1$ , we have

$$\begin{aligned} H(x, y) &= xy[1 - (1 - x)(1 - y)] \\ &= x^2y + y^2x - x^2y^2, \quad 0 \leq x, y \leq 1. \end{aligned}$$

- The density is

$$\begin{aligned} h(x, y) &= \frac{\partial^2}{\partial x \partial y} H(x, y) \\ &= 2x + 2y - 4xy, \quad 0 \leq x, y \leq 1. \end{aligned}$$

- [Here is a link](#) to a 3D rendering of this function.

## Example: constructing bivariate cdfs IV

- Now, let's select  $\alpha = 1/2$ :

$$\begin{aligned} H(x, y) &= xy \left( 1 + \frac{1}{2} (1 - F(x)) (1 - G(y)) \right) \\ &= \frac{1}{2} x^2 y^2 - \frac{1}{2} x^2 y - \frac{1}{2} x y^2 + \frac{3}{2} xy. \end{aligned}$$

- Taking the derivative, we get:

$$\begin{aligned} h(x, y) &= \frac{\partial^2}{\partial x \partial y} H(x, y) \\ &= 2xy - x - y + \frac{3}{2}, \quad 0 \leq x, y \leq 1. \end{aligned}$$

- [Here is a link](#) to a 3D rendering of this function.

## Example: constructing bivariate cdfs V

- The last two joint cdfs were examples of a **copula**.

### Definition: Copulas

A copula is a joint cdf that has uniform marginal distributions.

- Let  $C(u, v)$  be a copula. One immediate consequence of the definition is that if  $U$  and  $V$  are uniform random variables, then  $P(U \leq u) = C(u, 1) = u$ , and  $P(V \leq v) = C(1, v) = v$ .

## Example: constructing bivariate cdfs VI

- Let  $C(u, v)$  be a copula, we will restrict ourselves to the case where it is twice differentiable, such that
$$c(u, v) = \frac{\partial^2}{\partial u \partial v} C(u, v) \geq 0.$$
- let  $F_X$  and  $F_Y$  be the cdfs of  $X$  and  $Y$ , resp.
- Now define  $U = F_X(X)$ , and  $V = F_Y(Y)$ . From Proposition 2.2,  $U$  and  $V$  are uniformly distributed.
- Now consider the function
$$H(x, y) = C(u, v) = C(F_X(x), F_Y(y)).$$

## Example: constructing bivariate cdfs VII

- Thus, by the property that  $C(u, 1) = u$  and  $C(1, v) = v$ , we have

$$C(F_X(x), 1) = F_X(x)$$

$$C(1, F_Y(y)) = F_Y(y).$$

Therefore by definition,

$$F_{XY}(x, y) = H(x, y) = C((F_X(x), F_Y(y))).$$

- Using the chain rule, we can differentiate to obtain

$$f_{XY}(x, y) = c(F_X(x), F_Y(y)) f_X(x) f_Y(y).$$

## Example: constructing bivariate cdfs VIII

- **Takeaway:** We took arbitrary marginal distributions  $F_X$  and  $F_Y$ , and created a family of joint density functions, defined by *any* copula. Thus: the marginal distributions do not determine the joint distribution.
- There is a Theorem known as Sklar's Theorem (Wikipedia contributors, 2025) that generalizes this statement: All joint distributions can be expressed using a copula and marginal distributions, *and* the representation is unique.
- That is, the copula can be thought of as a way to describe the dependence between the variables in any joint distribution.

## Uniform on specific region

- So far when we have talked about *uniform distributions*, we think about being uniform over  $[0, 1]$ , or a higher dimensional box:  $[a, b]^d$ .
- It's often useful to have a uniform distribution for other regions of space.
- Let  $R \subset \mathbb{R}^2$  be any region of interest. The two-dimensional uniform distribution over  $R$  is defined by the probability

$$P((X, Y) \in A) = \frac{|A|}{|R|},$$

where  $||$  denotes the measure of the area.



## Uniform on specific region II

- Example: Suppose a point is chosen randomly in a disk of radius 1.
- The area of the disk is  $\pi r^2 = \pi$ , and therefore the joint pdf for the location  $(X, Y)$  is

$$f(x, y) = \begin{cases} \frac{1}{\pi} & x^2 + y^2 \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

- Now let  $R$  be the random variable denoting the distance of the point from the origin.

## Uniform on specific region III

- Note that  $R \leq r$  if and only if the point lies in a disk of radius  $r$ . This disk has area  $\pi r^2$ , and therefore the joint probability is

$$P(R \leq r) = \frac{\pi r^2}{\pi} = r^2, \quad 0 \leq r \leq 1.$$

- Taking a derivative, the corresponding density function is

$$f_R(r) = 2r, \quad 0 \leq r \leq 1.$$

## Uniform on specific region IV

- Now let us compute the marginal density of the  $x$  coordinate:

# Independent Random Variables

---

# Independence

## Definition: Independent Random Variables

Random variables  $X_1, X_2, \dots, X_n$  are said to be **independent** if their joint cdf factors into the product of their marginal cdf's:

$$F(x_1, x_2, \dots, x_n) = F_{X_1}(x_1)F_{X_2}(x_2) \cdots F_{X_n}(x_n)$$

for all  $x_1, x_2, \dots, x_n$ .

- This definition holds for both continuous and discrete random variables.
- For discrete RVs, it is equivalent to state that their joint pmf factors.
- For continuous RVs, it is equivalent to state that their joint pdf factors.

## Independence II

- To see why this is true, consider the case of two RVs,  $X, Y$ .
- From the definition, if they are independent, then
$$F(x, y) = F_X(x)F_Y(y).$$
- Taking the second mixed partial derivative makes it immediately clear that the joint pdf  $f(x, y)$  factors (assuming all densities exist).

## Independence III

- Conversely, suppose that the densities factor. Then by definition:

$$\begin{aligned}F(x, y) &= \int_{-\infty}^x \int_{-\infty}^y f(u, v) dv du \\&= \int_{-\infty}^x \int_{-\infty}^y f_X(u) f_Y(v) dv du \\&= \left( \int_{-\infty}^x f_X(u) du \right) \left( \int_{-\infty}^y f_Y(v) dv \right) \\&= F_X(x) F_Y(y).\end{aligned}$$

- It can also be shown that the definition implies that if  $X$  and  $Y$  are independent, then

$$P(X \in A, Y \in B) = P(X \in A)P(X \in B).$$

## Independence IV

- Furthermore, if  $g$  and  $h$  are functions on  $\mathbb{R}$ , then  $Z = g(X)$  and  $W = h(Y)$  are also independent.



# Conditional Distributions

---

## Conditional distributions: discrete RVs

- If  $X$  and  $Y$  are jointly distributed discrete RVs, the *conditional probability* that  $X = x_i$  given that  $Y = y_i$  is

$$\begin{aligned} P(X = x_i | Y = y_i) &= \frac{P(X = x_i, Y = y_i)}{P(Y = y_i)} \\ &= \frac{p_{XY}(x_i, y_i)}{P_Y(y_i)}, \end{aligned}$$

- If  $p_Y(y_i) = 0$ , the probability above is defined to be zero.
- We denote this conditional probability as  $p_{X|Y}$ .
- It's important to note that the conditional pmf is a genuine pmf, as it is non-negative and sums to one.
- If  $X$  and  $Y$  are independent,  $p_{Y|X}(y|x) = p_Y(y)$ .

## Conditional distributions: discrete RVs II

- Let's return to a previous joint pmf example (Table 2).

|     | $y$           |               |               |               |
|-----|---------------|---------------|---------------|---------------|
| $x$ | 0             | 1             | 2             | 3             |
| 0   | $\frac{1}{8}$ | $\frac{2}{8}$ | $\frac{1}{8}$ | 0             |
| 1   | 0             | $\frac{1}{8}$ | $\frac{2}{8}$ | $\frac{1}{8}$ |

**Table 2:** Frequency table for  $X$  and  $Y$ , flipping a fair coin three times.

- The conditional frequency function of  $X$  given  $Y = 1$  is:

$$p_{X|Y}(0|1) = \frac{2/8}{3/8} = 2/3$$

$$p_{X|Y}(1|1) = \frac{1/8}{3/8} = 1/3$$

## Conditional distributions: discrete RVs III

- The definition of the conditional frequency can be reexpressed as

$$p_{XY}(x, y) = p_{X|Y}(x|y)p_Y(y).$$

- By summing up over all possible values of  $y$ , we have the following

$$p_X(x) = \sum_y p_{X|Y}(x|y)p_Y(y).$$

- Both of these identities resemble what we have already seen previously when talking about probabilities: The multiplication principle and the law of total probability.

## Conditional distributions: discrete RVs IV

### Example: Counting particles

Suppose that a particle counter is imperfect; for each particle, it detects the particle with probability  $0 < p < 1$ . If the number of incoming particles in a unit of time is a Poisson distribution with parameter  $\lambda$ , what is the distribution of the number of counted particles?

## The continuous case

- Although a formal argument is beyond the scope of this course, the definition for conditional density of  $Y|X$  will be analogous to the discrete case.

### Definition: Conditional density

Let  $X, Y$  be jointly continuous random variables with joint density  $f_{XY}(x, y)$  and marginal densities  $f_X(x)$  and  $f_Y(y)$ , respectively. Then the conditional density of  $Y$  given  $X$  is defined to be

$$f_{Y|X}(y|x) = \frac{f_{XY}(x, y)}{f_X(x)},$$

if  $0 < f_X(x) < \infty$ , and 0 otherwise.

## The continuous case II

- A heuristic argument of why this definition makes sense is provided in Rice (2007, Section 3.5.2) using differentials.
- With our definition, we can express the joint density in terms of the marginal and conditional densities:

$$f_{XY}(x, y) = f_{Y|X}(y|x)f_X(x).$$

- We often use this expression to find marginal densities, using principles we have already discussed.

$$f_Y(y) = \int_{\mathbb{R}} f_{XY}(x, y)dx = \int_{\mathbb{R}} f_{Y|X}(y|x)f_X(x)dx.$$

- We can think of the above expression as the law of total probability for the continuous case.

## Example: finding conditional densities

- Let  $X$  and  $Y$  be jointly distributed random variables with joint and marginal densities

$$f_{XY}(x, y) = \lambda^2 e^{-\lambda y}, \quad 0 \leq x \leq y$$

$$f_X(x) = \lambda e^{-\lambda x}, \quad x \geq 0$$

$$f_Y(y) = \lambda^2 y e^{-\lambda y}, \quad y \geq 0$$

- Note that if  $x$  is held constant, the joint density decays exponentially in  $y$  for  $y \geq x$ .
- If  $y$  is held constant, the joint density is constant for  $0 \leq x \leq y$ .



## Example: finding conditional densities II

- Find the conditional densities for  $Y|X$  and  $X|Y$ .

## Example: finding conditional densities III

- Suppose we wanted to generate samples from the joint distribution  $(X, Y)$ ; how can this be done?
- Using what we have found about the conditional distributions, there are two simple ways for this to be done. Recall that the joint density is

$$f_{XY}(x, y) = f_{X|Y}(x|y)f_Y(y) = f_{Y|X}(y|x)f_X(x).$$

1. We could generate  $X$ , which is an exponential random variable ( $f_X(x)$ ). Then, we could generate  $Y$  conditioned on the simulated value of  $X = x$ , which follows an exponential distribution on the interval  $[x, \infty)$ .
2. Similarly, we can note that  $Y$  has a gamma distribution, and therefore generate a  $y$  following a gamma distribution, and then generate a value from  $X|Y = y$ , which is uniform on  $[0, y]$ .

# The rejection method

- We are often interested in generating random variables from a density function.
- If we have a closed form of the inverse cdf, we can use the “inverse cdf method” (Proposition 2.3).
- If a closed-form of the inverse cdf is not available, a commonly used approach is known as **rejection sampling**.

## The rejection method II

- Setup: let  $f$  be a density function we wish to simulate from, that is non-zero on an interval  $[a, b]$ .
- Pick a function  $M(X)$  such that  $M(x) \geq f(x)$  on  $[a, b]$ , and let

$$m(x) = \frac{M(x)}{\int_a^b M(x)dx}.$$

- Note that  $m(x)$ , as defined, is a probability density function. Then, to generate RV with density  $f$ , we can do the following:


Step 1: Generate  $T$  with density  $m$ .

Step 2: Generate  $U \sim U[0, 1]$  independent of  $T$ . If

$M(T) \times U \leq f(T)$ , then we “accept”  $T$  as a sample ( $X = T$ );  
otherwise, we “reject” and go back to Step 1.

# Rejection Method Figure

- A geometric justification is randomly throwing a dart (uniformly) at Figure 1.



**Figure 1:** Illustration of the rejection method, copied from Rice (2007, Figure 3.15).

- If the dart lands below the curve  $f$ , record the  $x$  coordinate; otherwise, reject it. With enough throws, the distribution of  $x$  coordinates will be proportional to the height of the curve.

## Rejection sampling

- A more formal argument using differentials is given in Rice (Example D 2007, Figure 3.15).
- In order for the rejection method to be worth-while (computationally efficient), it is important that the algorithm has high-acceptance (good choice of  $M$ ), otherwise you may need a large number of samples because many are being rejected.

# **Functions of Jointly Distributed Random Variables**

---

# Convolutions

- Suppose that  $X$  and  $Y$  are discrete random variables that take values on the integers and joint pmf  $p(x, y)$ .
- Find the pmf of  $Z = X + Y$ .
- Note that  $Z = z$  only when  $X = x$  and  $Y = z - x$ , whenever  $x$  is an integer.
- Thus, using the law of total probability, we can write

$$p_Z(z) = \sum_{x=-\infty}^{\infty} p(x, z - x).$$

- If  $X$  and  $Y$  are independent, then  $p(x, y) = p_X(x)p_Y(y)$ , and

$$p_Z(z) = \sum_{x=-\infty}^{\infty} p_X(x)p_Y(z - x).$$



## Convolutions II

- This sum is called the **convolution** of the sequences  $p_X$  and  $p_Y$ .
- The continuous case is similar. Let  $X$  and  $Y$  be jointly continuous RVs, and  $Z = X + Y$ .

## Convolutions III

- If we want to find the cdf of  $Z$ , then:

$$\begin{aligned}F_Z(z) &= P(Z \leq z) \\&= P(X + Y \leq z) \\&= \int \int_{\{x+y \leq z\}} f(x, y) dy dx \\&= \int_{-\infty}^{\infty} \int_{-\infty}^{z-x} f(x, y) dy dx \\&= \int_{-\infty}^{\infty} \int_{-\infty}^z f(x, v - x) dv dx \\&= \int_{-\infty}^z \int_{-\infty}^{\infty} f(x, v - x) dx dv\end{aligned}$$

## Convolutions IV

- Differentiating both sides, the fundamental theorem of calculus (with proper assumptions) gives

$$f_Z(z) = \int_{-\infty}^{\infty} f(x, z-x) dx$$

- Like in the discrete case, if  $X$  and  $Y$  are independent, then

$$f_Z(z) = \int_{-\infty}^{\infty} f_X(x) f_Y(z-x) dx$$

- This integral is called the **convolution** of the functions  $f_X$  and  $f_Y$ .

## Example: Sum of Exponential RVs

Suppose that the lifetime of an electrical component is exponentially distributed with rate  $\lambda$ , and that an independent and identical backup is available. If the system operates as long as one of the components is functional, and the components will not be replaced if they fail, what is the distribution of the life of the system?

*Solution:*

## Convolutions VI

- In the previous example, note carefully the change in integration:
  - The exponential density is only positive when  $t > 0$ , and zero every where else.
  - Thus, from  $(-\infty, 0)$ , the integral is zero.
  - Similarly, we evaluate the density at  $x - t$ , and hence when  $t > x$ , the integral is also zero.
- You may notice that the density of  $X = T_1 + T_2$  that we calculated is the same as a gamma distribution with parameters 2 and  $\lambda$ .

## Quotients of random variables

- Let  $X$  and  $Y$  be jointly continuous random variables, and let  $Z = Y/X$ .
- Our derivation for the pdf of  $Z$  is similar as what we did with the sum: find the cdf, then take the derivative.
- $F_Z(z) = P(Z \leq z) = P(Y/X \leq z)$ . Thus, we are interested in the probability of the set  $\{x, y : y/x \leq z\}$ .
- We have to be a little careful about what happens if  $X = 0$ , so we will split it into two parts:
  - If  $x > 0$ , then the set is  $y \leq xz$ .
  - If  $x < 0$ , then the set is  $y \geq xz$ .

## Quotients of random variables II

Thus,

$$F_Z(z) = \int_{-\infty}^0 \int_{xz}^{\infty} f(x, y) dy dx + \int_0^{\infty} \int_{-\infty}^{xz} f(x, y) dy dx.$$

- To remove dependence of the inner integrals on  $x$ , we make the change of variables  $y = xv$ :

$$\begin{aligned} F_Z(z) &= \int_{-\infty}^0 \int_z^{-\infty} x f(x, xv) dv dx + \int_0^{\infty} \int_{-\infty}^z x f(x, xv) dv dx \\ &= \int_{-\infty}^0 \int_{-\infty}^z (-x) f(x, xv) dv dx + \int_0^{\infty} \int_{-\infty}^z x f(x, xv) dv dx \\ &= \int_{-\infty}^z \int_{-\infty}^{\infty} |x| f(x, xv) dx dv \end{aligned}$$

## Quotients of random variables III

- And differentiating both sides, we obtain

$$f_Z(z) = \int_{-\infty}^{\infty} |x| f(x, xz) dx.$$

- If  $X$  and  $Y$  are independent,

$$f_Z(z) = \int_{-\infty}^{\infty} |x| f_X(x) f_Y(xz) dx.$$



## References and Acknowledgements

- Compiled on August 18, 2025 using R version 4.5.1.
- Licensed under the [Creative Commons Attribution-NonCommercial license](#). Please share and remix non-commercially, mentioning its origin.
- We acknowledge [students and instructors for previous versions of this course / slides](#).



## References and Acknowledgements II

Rice JA (2007). *Mathematical statistics and data analysis*, volume 371. 3 edition. Thomson/Brooks/Cole Belmont, CA.

Wikipedia contributors (2025). “Copula (statistics)#Sklar’s Theorem: Wikipedia, The Free Encyclopedia.” [Online; accessed 18-August-2025], URL [https://en.wikipedia.org/w/index.php?title=Copula\\_\(statistics\)&oldid=1303484991](https://en.wikipedia.org/w/index.php?title=Copula_(statistics)&oldid=1303484991).