

Mathematical Statistics II

The Bayesian Approach to Parameter Estimation

Jesse Wheeler

Introduction

Bayesian Estimation

- Much of this work is based on Rice (2007, Section 8.6).
- We have already discussed the philosophy of Bayesian statistics.
- We start with a prior belief about parameter values, and update these beliefs using observed data.
- The resulting **distribution** is called the *posterior*, and it represents our updated belief after observing data.
- This is very natural idea that is closely related to the idea of likelihood: likelihood quantifies some degree of belief about a parameter value.

Review

Some Review

- Before we begin, we will first do a bit of review.
- In the context of Bayesian inference, we treat unknown parameter vectors as random variables, which I will denote Θ .
- Thus, our probability model can be expressed as $f(x|\Theta = \theta)$, which we often shorten to $f(x|\theta)$.

Some Review II

Bayes' Theorem

Let X be the random vector representing observed data, and Θ the random parameter vector, and x^* the observed data. Bayes Theorem states:

$$\begin{aligned}\pi_{\Theta|X}(\theta|x^*) &= \frac{f_{X|\Theta}(x^*|\theta)\pi_{\Theta}(\theta)}{f_X(x^*)} \\ &= \frac{f_{X|\theta}(x^*|\theta)\pi_{\Theta}(\theta)}{\int f_{X|\Theta}(x^*|\tau)\pi_{\Theta}(\tau) d\tau}\end{aligned}$$

- As before, f is taken to be either a pmf or pdf, depending on the problem.

Some Review III

Flipping 10 coins

Our friend hands us a coin from another country, and we want to estimate $\theta = p$, the probability that the coin lands heads.

Suppose we flip a coin 10 times, and see n heads. Find a Bayesian estimate for θ .

Some Review IV

- Even in the simple problem above, we see two of the primary challenges with Bayesian parameter estimation:
 - How do we choose the prior distribution $\pi(\theta)$? A generally safe and accepted approach is a uniform prior. However, this formally only exists if θ is bounded, which is not always the case. Also, it represents a prior belief: given a new coin, do we really think all values of p are equally likely, or maybe values close to $p = 0.5$ are more likely than extreme values $p = 0, 1$? Since the prior represents our beliefs about θ , is a uniform prior actually appropriate? If it isn't appropriate, how exactly should we specify the prior?
 - Even in this very simple model and prior, the denominator $f(x)$ was difficult to compute. What about more complex models and priors? A large amount of Bayesian computation and theory is dedicated to solving this problem.

Proposition: the MAP and MLE

Let θ be a parameter of interest, and x^* the observed data. If our prior distribution is proportional to 1, i.e., $\pi(\theta) \propto 1$ (which is effectively a uniform prior on a bounded interval), then

$$\hat{\theta}_{\text{MAP}} = \hat{\theta}_{\text{MLE}}.$$

- This is true for the Coin-tossing example; look back at the likelihood function and posterior, and use R to plot them both.

Examples

Bayesian point-estimate examples

Poisson model

Suppose we have observations n observations, which we wish to model as IID $\text{Poisson}(\lambda)$. Find a Bayesian estimate of $\Lambda = \lambda$ given the observed data x^* .

Bayesian point-estimate examples II

Poisson posterior, uniform prior

Revisit the $\text{Poisson}(\lambda)$ model, while taking the alternative approach of using a uniform prior.

Real-data example: Poisson Distribution

- Now let's look at a real-data example. These data are the 23 observations from the asbestos-filter problem.

```
x <- c(  
  31, 29, 19, 18, 31, 28, 34, 27, 34, 30, 16, 18,  
  26, 27, 27, 18, 24, 22, 28, 24, 21, 17, 24  
)  
x
```

```
[1] 31 29 19 18 31 28 34 27 34 30 16 18 26 27 27 18 24 22  
[19] 28 24 21 17 24
```

Real-data example: Poisson Distribution II

Comparing Estimates

Using the data above, compare estimates using the MoM, MLE, and the two Bayesian approaches, as well as the corresponding errors related to these estimates.

Posteriors and Likelihood

- In the problem above, we saw that we get very similar estimates using MLE or Bayesian approaches, regardless of which prior we picked.
- We can argue why this will often be the case, especially for IID data.
- Previously, we saw:

$$\text{posterior} \propto \text{likelihood} \times \text{prior}$$

- When n gets large, the likelihood dominates in this equation.
In the IID case:

$$\text{likelihood} = \prod_{i=1}^n f(x_i^* | \theta).$$

Posteriors and Likelihood II

- In particular, each new data point scales the likelihood larger and larger, to the point where the prior has little impact on the posterior distribution.
- See the accompanying Lecture 4 R code for a visual demonstration of this using the Poisson distribution.

Introduction to Numeric Integration

Numeric Integration

- As we saw in the previous example, one of the primary challenges of Bayesian estimation is the integration in the denominator of the posterior.
- Bayesian statistics has really exploded since the late 20th century, largely thanks to improved computational tools that help with the numeric integration.
- For this set of lectures, we only briefly introduce this topic. Depending on time and interest, we can explore this topic more later in the semester.

Choice of Priors

Conjugate priors

- The Poisson(λ) example showed two main approaches:
 - The traditional (subjective) Bayesian, who takes seriously the choice of prior, and chose a Gamma density to aid computations.
 - The utilitarian (objective) Bayesian, who picked an uninformative prior.
- The former approach was aided by what is known as a **conjugate prior**.

Conjugate priors II

Definition: Conjugate priors

Suppose the prior distribution belongs to a family of distributions, G , and the data come from a family of distributions H .

G is said to be conjugate to H if the posterior is in the family G .

- Example: If the data-model is $\text{Poisson}(\lambda)$, then the family H is the family of Poisson distributions. The Gamma family (G) of distributions is conjugate to the Poisson family, because if Gamma is selected as the prior distribution, then the posterior distribution (under data model H) is still Gamma (G), with updated parameters.

Conjugate priors III

- Much of the Bayesian statistics of the 20th century relied on conjugate priors to help with integration, or were confined to models with very few parameters.
- Recent developments in computing, both hardware, software, and theory of Bayesian computing, has enabled fitting much more complex models using arbitrary priors.
- Still, it's worth discussing conjugate priors, and we will provide a few examples.

- TODO

Hierarchical Bayes

Hierarchical Bayes

- The idea behind Hierarchical Bayes is simple: our model f depends on parameters θ .
- We can get a prior for θ , $\pi(\theta)$.
- The prior itself depends on parameters, say $\pi(\theta; \psi)$.
- How do we choose ψ ? Why don't we put a prior on these as well?
- In some way, this allows us to be less-committal about the parameters in the prior model, and instead allow the data to inform our choice of priors (to some degree).
- Philosophically, this situation naturally arises if we want to pick a conjugate prior for Θ , but are not committal about the **hyperparameters** Ψ that define the distribution of Θ . Then, Ψ should also be modeled as a random variable.

Hierarchical Bayes II

- This leads to a hierarchical model:

$$X|\Theta = \theta, \Psi = \psi \sim f(x|\theta, \psi),$$

and

$$\Theta|\Psi = \psi = g(\theta|\psi), \quad \Psi \sim \pi(\psi).$$

- Why would we want to do this? Do we need a prior now hyper-parameters for ψ ?
 1. We'll see that this is sometimes a very useful way of thinking about a problem.
 2. We could, but this is not usually useful.

Hierarchical Bayes III

Trivial case: hierarchical Normal-Normal

Suppose that the data X_i are iid $N(\theta, 1)$. Set a prior for θ as $\Theta|M = \mu \sim N(\mu, 1)$, and $M \sim N(0, 1)$.

Empirical Bayes

TODO

Uncertainty quantification

Uncertainty in Bayes estimates

- TODO

References and Acknowledgements

Rice JA (2007). *Mathematical statistics and data analysis*, volume 371. 3 edition. Thomson/Brooks/Cole Belmont, CA.

- Compiled on January 29, 2026 using R version 4.5.2.
- Licensed under the [Creative Commons Attribution-NonCommercial](#) license. Please share and remix non-commercially, mentioning its origin.
- We acknowledge [students and instructors for previous versions of this course / slides](#).

