Random Effects Models

Jesse Wheeler

Overview

Objectives and overview

- This material is based Chapter 11–12 on the textbook "A First Course in Design and Analysis of Experiments" (Oehlert, 2000).
- Chapter outcomes include:
  - Students will be able to distinguish between a random effect and a fixed effect.
  - Students will be able to identify when random effects are appropriate.
  - Students will know how to implement random effects and mixed effects models.
  - Students will gain some familiarity with the `lme4` package in R.

Introduction

- So far, the treatment effects that we have been working with can be classified as fixed-effects: we wish to estimate the effect of specific (fixed) treatments.
- What if the treatments themselves are a random selection from a larger population?
- **Example:** A farmer has 100 pieces of equipment, and would like to know how the equipment used may impact crop-yield. Rather than testing all 100 pieces of equipment, they *randomly select* g pieces of equipment, and performs a CRD with these equipment.

Introduction II

Random Effects

A random effect is appropriate to model treatments that are a *random sample* from a population of potential treatments.

Random effects are useful if the focus of inference is on the population, not the individual treatment effects.

Models for Random Effects

Example: Carton factory

- We will build intuition of Random Effects models and designs using an example.

Carboard company

Suppose a company makes cardboard cartons for storing goods. They have 50 machines to make cartons, and want to understand the variation in quality for finished products.

They choose 10 machines at random from the 50, and make 40 cartons on each machine. The resulting cartons are tested for strength.

Example: Carton factory II

- This is a completely randomized design (CRD), with $g = 10$ treatments (the different machines), and $n_i = 40$, for a total of $N = 400$ EUs.
- The model we have been using for this type of data is:

$$y_{ij} = \mu + \alpha_i + \epsilon_{ij}, \quad \epsilon_{ij} \overset{iid}{\sim} N(0, \sigma^2).$$

- We are still going to use a very similar model. However, we are not as interested in the exact α_i for the 10 randomly chosen machines. Instead, we want to make inference on the larger population.

Example: Carton factory III

- We could proceed as usual, but that only tells us about the 10 specific machines that were randomly chosen, and we would get an entirely different set of machines if we repeated the experiment.
- Thus, we want to introduce a random effect: let the effect α_i represent a random draw from a population of possible effects:

$$y_{ij} = \mu + \alpha_i + \epsilon_{ij}, \qquad \alpha_i \overset{iid}{\sim} N(0, \sigma_\alpha^2), \quad \epsilon_{ij} \overset{iid}{\sim} N(0, \sigma^2).$$

- We no longer assume $\sum_i \alpha_i = 0$, rather $E[\alpha_i] = 0$.
- We have introduced a new parameter, σ_α^2, which describes the spread of the effects.
- We assume that the two random components are independent. That is, α_i and ϵ_{ij} are independent.

Example: Carton factory IV

- If they are independent, the variance of the observed data is increased:

$$\mathrm{Var}(y_{ij}) = \mathrm{Var}(\mu + \alpha_i + \epsilon_{ij}) = \sigma_\alpha^2 + \sigma^2.$$

- The parameters σ_α^2 and σ^2 are called *variance components*.
- If observations belong to the same treatment group, then they now have non-zero correlation:

$$\mathrm{Cor}(y_{ij}, y_{ik}) = \begin{cases} 1 & j = k \\ \sigma_\alpha^2/(\sigma_\alpha^2 + \sigma^2) & j \neq k \end{cases}$$

- We call this within treatment group correlation the intraclass correlation.

Example: Carton factory V

Key summary

- In a random effects model, the parameters are μ, σ^2, and σ^2_α.
- We are not interested in making inferences about the specific α_i and ϵ_{ij}, which are treated as random variables, not parameters.
- Inference wise, we are now interested in confidence intervals and hypothesis tests regarding μ and σ^2.

Estimation: REML and LRT

Beyond ANOVA: Maximum Likelihood

- Classical random effects estimation relies on Expected Mean Squares (EMS) from ANOVA tables (See Oehlert (2000, Sections 11.3–11.6)).
- These approaches can be computed by hand, and would naturally extend our workflow with the `aov()` function and the `emmeans` package.
- Unfortunately, these approaches have their weaknesses for random (and mixed) effects models.
- Our models will remain the same, but how we analyze them will shift, keeping in-line with modern practices.
- For the type of models considered so far (fixed-effects), the two approaches are the same.

Beyond ANOVA: Maximum Likelihood II

- Modern software relies on Likelihood methods.
- **Maximum Likelihood (ML)** finds the parameter estimates (mean and variances) that make the observed data most probable; confidence intervals are obtained using likelihood ratio tests.

Beyond ANOVA: Maximum Likelihood III

- However, standard ML has a flaw when estimating variance components: it does not account for the degrees of freedom used to estimate the fixed effects (like the grand mean).
- This makes standard ML variance estimates *biased* (they underestimate the true variance).

Restricted Maximum Likelihood (REML)

- To fix the bias in standard ML, modern software uses Restricted Maximum Likelihood (REML).
- REML mathematically removes the fixed effects from the likelihood equation before estimating the variance components.
- By accounting for the degrees of freedom lost in estimating fixed effects, REML provides *unbiased* estimates of σ^2 and σ^2_α.
- Additionally, REML handles unbalanced data effortlessly and prevents the impossible "negative variance" estimates that can sometimes occur when calculating variance using classical EMS methods.

Estimating variance components in R

- If we are primarily interested in the variance components σ^2_α, we will want to get the REML estimate for this component, as it is unbiased.
- We will shift from our `aov()` workflow and use the `lmer` function from the `lme4` package to do this in R:

```
# install.packages("lme4")
library(lme4)
lmer(response ~ 1 + (1 | randomEffect), data = data)
```

Example: Penicillin

- We will first consider the built-in `Penicillin` data in R, which contains the reduced diameter of bacterial growth, after penicillin application. Each measurement is taken on a specific `plate`, which is our random effect.

```
data("Penicillin")

output <- lmer(
  diameter ~ 1 + (1 | plate),
  data = Penicillin, REML = TRUE   # default
)
```

Example: Penicillin II

- What is the notation here doing? `1 + (1 | plate)`.
  - The model we are fitting is $y_{ij} = \mu + \alpha_i + \epsilon_{ij}$.
  - The `1` in the beginning is not strictly needed, but represents the grand mean μ.
  - The `(1 | plate)` represents the random effect. In `lme4`, the syntax `(A | B)` implies that B is the random effect grouping variable, and A is the effect that is random. For a simple random effects model, the mean (`1`) is always the thing that is random.
- Now let's look at the output.

Example: Penicillin III

```
summary(output)

Linear mixed model fit by REML ['lmerMod']
Formula: diameter ~ 1 + (1 | plate)
   Data: Penicillin

REML criterion at convergence: 613.3

Scaled residuals:
     Min       1Q   Median       3Q      Max
-2.41583 -0.55523  0.01212  0.59232  1.95241

Random effects:
 Groups   Name        Variance Std.Dev.
 plate    (Intercept) 0.09509  0.3084
 Residual             4.03333  2.0083
Number of obs: 144, groups:  plate, 24
```

Confidence intervals

- We can get confidence intervals for our estimates using the `confint` function.
- The two primary methods will be `profile` (default) and `boot`.
- This computes profile likelihood confidence intervals, which are effectively the same as a **likelihood ratio test** (LRT).

```
set.seed(123)
confint(output, signame = FALSE, level = 0.95, method = "boot")
```

```
                          2.5 %     97.5 %
sd_(Intercept)|plate  0.000000  0.8138926
sigma                 1.771368  2.2218709
(Intercept)          22.656619 23.3378052
```

Hypothesis testing

- For random effects, the hypothesis that we would like to test is:

$$H_0 : \sigma_\alpha^2 = 0, \quad H_1 : \sigma_\alpha^2 > 0$$

- This is always one-sided, because σ_α^2 can't be negative.
- Notice: Our assumption was $\alpha_i \sim N(0, \sigma_\alpha^2)$. If H_0 is true, then $\sigma_\alpha^2 = 0$, and $\alpha_i = 0$ for all i.
- Thus, for a simple random effects model, we have the same null hypothesis and tests!

$$H_0 : \sigma_\alpha^2 = 0 \iff \alpha_i = 0 \text{ for all } i$$

Hypothesis testing II

- Because of this, we *could* just use a standard F-test for random effects.
- However: this approach does not generalize well to more complex models with random effects.
- Rather than using an F-test, we will introduce the idea of a Likelihood Ratio Test.

Likelihood Ratio Tests (LRT)

- Instead of using exact F-tests derived from ratios of Mean Squares, modern mixed modeling often uses the Likelihood Ratio Test (LRT) for hypothesis testing.
- The LRT explicitly compares two competing models:
  1. A **Full Model** that includes the random effect.
  2. A **Reduced Model** (Null Model) that removes the random effect.
- The test measures how much the likelihood (fit) of the data drops when the random effect is removed.

$$\Lambda = \frac{\text{Likelihood}_{\text{reduced}}}{\text{Likelihood}_{\text{Full}}}, \quad \Longrightarrow \quad -2\log\Lambda \sim \chi^2_d$$

Likelihood Ratio Tests (LRT) II

- Using properties of logarithms, we often consider differences in log-likelihood:

$$\ell_{\text{full}} - \ell_{\text{reduced}} \sim \frac{1}{2}\chi^2_d,$$

where d is the difference in number parameters between the models.

Likelihood Ratio Tests (LRT) III

- The test can be extended for restricted likelihood ratio tests (RLRT); the idea is similar: fit two models (full vs reduced) using REML, compare differences in the restricted log-likelihoods.
- Rather than fitting two models manually and comparing, we will use the `lmerTest` package to do this automatically for us.

Likelihood Ratio Tests (LRT) IV

```
# install.packages("lmerTest")
lmerTest::ranova(output)  # Will do RLRT or LRT, depending on ob

ANOVA-like table for random-effects: Single term deletions

Model:
diameter ~ 1 + (1 | plate)
             npar  logLik    AIC      LRT Df Pr(>Chisq)
<none>          3 -305.81 617.63
(1 | plate)     2 -305.86 615.71 0.082153  1     0.7744
```

Factorial Random Effects Models

Factorial models

- Just like with fixed effects, we may want to consider random effects with multiple random treatment factors.
- Again, we use random effects when we aren't interested in the specific treatment effects chosen, but the larger population of treatment effects.

Example: Carton Factory (continued)

- Let's expand our motivating example of the Carton Factory.

Two-way random effects model

Suppose the carton factory has multiple machines and would like to know the variance in the effect of these machines, but there are also different machine operators, and their effect may be random.

- We randomly select machines and operators, and have each operator work on each each machine and produce $n = 4$ cartons each.
- We have a two-way, completely randomized factorial design.

Example: Carton Factory (continued) II

- In the design proposed above, a two-way random effects model is natural:

$$y_{ijk} = \mu + \alpha_i + \beta_j + (\alpha\beta)_{ij} + \epsilon_{ijk},$$

with

$$\alpha_i \overset{iid}{\sim} N(0, \sigma^2_\alpha) \quad \beta_j \overset{iid}{\sim} N(0, \sigma^2_\beta)$$
$$(\alpha\beta)_{ij} \overset{iid}{\sim} N(0, \sigma^2_{\alpha\beta}) \quad \epsilon_{ijk} \overset{iid}{\sim} N(0, \sigma^2)$$

- The variance of each observation is

$$\mathrm{Var}(y_{ijk}) = \sigma^2_\alpha + \sigma^2_\beta + \sigma^2_{\alpha\beta} + \sigma^2.$$

Example: Carton Factory (continued) III

- This idea once again can generalize to higher-order factorial models

Parameter estimation

- The parameter estimation will still happen in a similar way: we can use ML or REML.
- If we are primarily interested in the error components, REML is unbiased, and generally preferred.

```
carton <- read.csv('carton.csv')
output <- lmer(
  thickness ~ 1 + (1 | Machine) + (1 | Operator) + (1 | Machine:
  # OR: 1 + (1 | Machine * Operator)
  data = carton,
  REML = TRUE
)
```

```
summary(output)

Linear mixed model fit by REML ['lmerMod']
Formula:
thickness ~ 1 + (1 | Machine) + (1 | Operator) + (1 | Machine:Op
   Data: carton

REML criterion at convergence: 317.2

Scaled residuals:
     Min       1Q   Median       3Q      Max
-1.58412 -0.78134 -0.07557  0.72636  2.15223

Random effects:
 Groups           Name        Variance Std.Dev.
 Machine:Operator (Intercept)  3.782   1.945
 Machine          (Intercept)  2.083   1.443
 Operator         (Intercept) 11.024   3.320
```

```
 Residual                        1.465   1.210
Number of obs: 80, groups:
Machine:Operator, 20; Machine, 5; Operator, 4

Fixed effects:
            Estimate Std. Error t value
(Intercept)   46.671      1.838   25.39
```

Confidence intervals

- We can obtain confidence intervals the same way as before. This can be very slow.

```
set.seed(444)
confint(output, method = 'boot', signame = FALSE)
```

```
                                    2.5 %    97.5 %
sd_(Intercept)|Machine:Operator  1.0920933  2.746098
sd_(Intercept)|Machine           0.0000000  3.031105
sd_(Intercept)|Operator          0.1061933  5.646589
sigma                            0.9914768  1.422556
(Intercept)                     43.4999935 50.636162
```

Hypothesis testing

- We can also do hypothesis testing, but it can get tricky due to the interaction effect.
- The `lmerTest::ranova` function just drops the term from the model, and compares REML (or ML). This may not be meaningful for all variables in the presence of an interaction effect.
- Instead, we often will rely on the confidence intervals.

Hypothesis testing II

```
# May not be meaningful for main (random) effects if interaction

lmerTest::ranova(output)

ANOVA-like table for random-effects: Single term deletions

Model:
thickness ~ 1 + (1 | Machine) + (1 | Operator) + (1 | Machine:Op
                        npar  logLik    AIC    LRT Df
<none>                     5 -158.62 327.24
(1 | Machine)              4 -159.67 327.35  2.105  1
(1 | Operator)             4 -164.39 336.77 11.531  1
(1 | Machine:Operator)     4 -180.10 368.19 42.952  1
                        Pr(>Chisq)
<none>
(1 | Machine)            0.1468521
(1 | Operator)           0.0006846 ***
```

Summary

- We fit a random effects model if we are interested in the variance components specifically, rather than specific treatment effects.
- Random effects are often very useful for **block-designs**. Though we may want to block to reduce variance, we are not often interested in the exact block effect; rather, we want to know how the general population of possible blocks might impact results.
- Though they can be useful, random effects are often coupled with fixed effects, to create a mixed effects model, which is covered in our next module.

Summary II

- Common situation: A fixed effect of interest, in a block-design; we will treat the block effect as random, giving a mixed effect model.

References and Acknowledgements

Oehlert GW (2000). *A First Course in Design and Analysis of Experiments*. W. H. Freeman. ISBN 0-7167-3510-5.

- Compiled on October 22, 2026 using R version 4.5.2.
- Licensed under the Creative Commons Attribution-NonCommercial license. Please share and remix non-commercially, mentioning its origin.
- We acknowledge students and instructors for previous versions of this course / slides.