Random Effects Models

Jesse Wheeler

Contents

1 Overview **1**

2 Models for Random Effects **2**

3 Estimation: REML and LRT **3**

1 Overview

Objectives and overview

- This material is based Chapter 11–12 on the textbook "A First Course in Design and Analysis of Experiments" (Oehlert, 2000).
- Chapter outcomes include:
  - Students will be able to distinguish between a random effect and a fixed effect.
  - Students will be able to identify when random effects are appropriate.
  - Students will know how to implement random effects and mixed effects models.
  - Students will gain some familiarity with the `lme4` package in R.

Introduction

- So far, the treatment effects that we have been working with can be classified as *fixed-effects*: we wish to estimate the effect of specific (fixed) treatments.
- What if the treatments themselves are a random selection from a larger population?
- **Example:** A farmer has 100 pieces of equipment, and would like to know how the equipment used may impact crop-yield. Rather than testing all 100 pieces of equipment, they *randomly select* g pieces of equipment, and performs a CRD with these equipment.

Random Effects

A random effect is appropriate to model treatments that are a *random sample* from a population of potential treatments.
Random effects are useful if the focus of inference is on the *population*, not the individual treatment effects.

2 Models for Random Effects

Example: Carton factory

- We will build intuition of Random Effects models and designs using an example.

Carboard company
Suppose a company makes cardboard cartons for storing goods. They have 50 machines to make cartons, and want to understand the variation in quality for finished products.
They choose 10 machines at random from the 50, and make 40 cartons on each machine. The resulting cartons are tested for strength.

- This is a completely randomized design (CRD), with $g = 10$ treatments (the different machines), and $n_i = 40$, for a total of $N = 400$ EUs.
- The model we have been using for this type of data is:

$$y_{ij} = \mu + \alpha_i + \epsilon_{ij}, \quad \epsilon_{ij} \overset{iid}{\sim} N(0, \sigma^2).$$

- We are still going to use a very similar model. However, we are not as interested in the exact α_i for the 10 randomly chosen machines. Instead, we want to make inference on the larger population.
- We could proceed as usual, but that only tells us about the 10 specific machines that were randomly chosen, and we would get an entirely different set of machines if we repeated the experiment.
- Thus, we want to introduce a *random effect*: let the effect α_i represent a random draw from a population of possible effects:

$$y_{ij} = \mu + \alpha_i + \epsilon_{ij}, \qquad \alpha_i \overset{iid}{\sim} N(0, \sigma^2_\alpha), \quad \epsilon_{ij} \overset{iid}{\sim} N(0, \sigma^2).$$

- We no longer assume $\sum_i \alpha_i = 0$, rather $E[\alpha_i] = 0$.
- We have introduced a new parameter, σ^2_α, which describes the spread of the effects.
- *We assume that the two random components are independent.* That is, α_i and ϵ_{ij} are independent.
- If they are independent, the variance of the observed data is increased:

$$\text{Var}(y_{ij}) = \text{Var}(\mu + \alpha_i + \epsilon_{ij}) = \sigma^2_\alpha + \sigma^2.$$

- The parameters σ^2_α and σ^2 are called *variance components*.
- If observations belong to the same treatment group, then they now have non-zero correlation:

$$\text{Cor}(y_{ij}, y_{ik}) = \begin{cases} 1 & j = k \\ \sigma^2_\alpha/(\sigma^2_\alpha + \sigma^2) & j \neq k \end{cases}$$

- We call this within treatment group correlation the *intraclass* correlation.

Key summary

- In a random effects model, the parameters are μ, σ^2, and σ^2_α.
- We are not interested in making inferences about the specific α_i and ϵ_{ij}, which are treated as random variables, not parameters.
- Inference wise, we are now interested in confidence intervals and hypothesis tests regarding μ and σ^2.

3 Estimation: REML and LRT

Beyond ANOVA: Maximum Likelihood

- Classical random effects estimation relies on Expected Mean Squares (EMS) from ANOVA tables (See Oehlert (2000, Sections 11.3–11.6)).
- These approaches can be computed by hand, and would naturally extend our workflow with the `aov()` function and the `emmeans` package.
- Unfortunately, these approaches have their weaknesses for random (and mixed) effects models.
- Our models will remain the same, but how we analyze them will shift, keeping in-line with modern practices.
- For the type of models considered so far (fixed-effects), the two approaches are the same.
- Modern software relies on *Likelihood* methods.
- **Maximum Likelihood (ML)** finds the parameter estimates (mean and variances) that make the observed data most probable; confidence intervals are obtained using likelihood ratio tests.
- However, standard ML has a flaw when estimating variance components: it does not account for the degrees of freedom used to estimate the fixed effects (like the grand mean).
- This makes standard ML variance estimates *biased* (they underestimate the true variance).

  Example: For a normal distribution model, $y_{ij} \sim N(\mu_i, \sigma^2)$, the MLE of (μ, σ^2) is:

$$\left(\hat{\mu}_{MLE}, \hat{\sigma}^2_{MLE}\right) = \left(\bar{Y}_i, \frac{1}{n}\sum_{i=1}^{n}(x_i - \bar{x})^2\right).$$

  However, you may recall from you intro stats class that we usually don't estimate variance that way, instead we divide by n_1:

$$s^2 = \frac{1}{n-1}\sum_{i=1}^{n}(x_i - \bar{x})^2$$

  That's because the MLE for the variance component is biased low: on average, our estimate will be slightly smaller than the truth.

Restricted Maximum Likelihood (REML)

- To fix the bias in standard ML, modern software uses *Restricted Maximum Likelihood (REML).*
- REML mathematically removes the fixed effects from the likelihood equation before estimating the variance components.
- By accounting for the degrees of freedom lost in estimating fixed effects, REML provides *unbiased* estimates of σ^2 and σ^2_α.

  Example: building on the normally distributed model from before, the MLE for σ^2 was biased low:

$$\hat{\sigma}^2_{MLE} = \frac{1}{n}\sum_{i=1}^{n}(x_i - \bar{x})^2.$$

  The REML estimate for this model is unbiased:

$$\hat{\sigma}^2_{REML} = s^2 = \frac{1}{n-1}\sum_{i=1}^{n}(x_i - \bar{x})^2$$

- Additionally, REML handles unbalanced data effortlessly and prevents the impossible "negative variance" estimates that can sometimes occur when calculating variance using classical EMS methods.

Estimating variance components in R

- If we are primarily interested in the variance components σ^2_α, we will want to get the REML estimate for this component, as it is unbiased.
- We will shift from our `aov()` workflow and use the `lmer` function from the `lme4` package to do this in R:

```
# install.packages("lme4")
library(lme4)
lmer(response ~ 1 + (1 | randomEffect), data = data)
```

Example: Penicillin

- We will first consider the built-in `Penicillin` data in R, which contains the reduced diameter of bacterial growth, after penicillin application. Each measurement is taken on a specific `plate`, which is our random effect.

```
data("Penicillin")

output <- lmer(
  diameter ~ 1 + (1 | plate),
  data = Penicillin, REML = FALSE  # default
)
```

- What is the notation here doing? `1 + (1 | plate)`.
  - The model we are fitting is $y_{ij} = \mu + \alpha_i + \epsilon_{ij}$.
  - The `1` in the beginning is not strictly needed, but represents the grand mean μ.
  - The `(1 | plate)` represents the random effect. In `lme4`, the syntax `(A | B)` implies that B is the random effect grouping variable, and A is the effect that is random. For a simple random effects model, the mean (`1`) is always the thing that is random.
- Now let's look at the output.

```
summary(output)

Linear mixed model fit by REML ['lmerMod']
Formula: diameter ~ 1 + (1 | plate)
   Data: Penicillin

REML criterion at convergence: 613.3

Scaled residuals:
     Min       1Q   Median       3Q      Max
-2.41583 -0.55523  0.01212  0.59232  1.95241
```

```
Random effects:
 Groups   Name        Variance Std.Dev.
 plate    (Intercept) 0.09509  0.3084
 Residual             4.03333  2.0083
Number of obs: 144, groups:  plate, 24

Fixed effects:
            Estimate Std. Error t value
(Intercept)  22.9722     0.1788   128.5
```

Confidence intervals

- We can get confidence intervals for our estimates using the `confint` function.
- The two primary methods will be `profile` (default) and `boot`.
- This computes *profile likelihood* confidence intervals, which are effectively the same as a **likelihood ratio test** (LRT).
- Mathematically, we can obtain a confidence interval using profiled restricted likelihoods, but unfortunately there's not an immediate way to do this in R; we can use `method = "boot"` for a bootstrap interval for REML estimates, or `method = "profile"` for a profile likelihood. Both should give very similar output.

```
set.seed(123)
confint(output, signame = FALSE, level = 0.95, method = "boot")

                          2.5 %     97.5 %
sd_(Intercept)|plate  0.000000  0.8138926
sigma                 1.771368  2.2218709
(Intercept)          22.656619 23.3378052
```

Hypothesis testing

- For random effects, the hypothesis that we would like to test is:

$$H_0 : \sigma_\alpha^2 = 0, \quad H_1 : \sigma_\alpha^2 > 0$$

- This is always one-sided, because σ_α^2 can't be negative.
- *Notice:* Our assumption was $\alpha_i \sim N(0, \sigma_\alpha^2)$. If H_0 is true, then $\sigma_\alpha^2 = 0$, and $\alpha_i = 0$ for all i.
- Thus, for a simple random effects model, we have the same null hypothesis and tests!

$$H_0 : \sigma_\alpha^2 = 0 \iff \alpha_i = 0 \text{ for all } i$$

- Because of this, we *could* just use a standard F-test for random effects.
- *However:* this approach does not generalize well to more complex models with random effects.
- Rather than using an F-test, we will introduce the idea of a Likelihood Ratio Test.

Likelihood Ratio Tests (LRT)

- Instead of using exact F-tests derived from ratios of Mean Squares, modern mixed modeling often uses the *Likelihood Ratio Test (LRT)* for hypothesis testing.
- The LRT explicitly compares two competing models:
  1. A **Full Model** that includes the random effect.
  2. A **Reduced Model** (Null Model) that removes the random effect.
- The test measures how much the likelihood (fit) of the data drops when the random effect is removed.

$$\Lambda = \frac{\text{Likelihood}_{\text{reduced}}}{\text{Likelihood}_{\text{Full}}}, \quad \Longrightarrow \quad -2\log\Lambda \sim \chi^2_d$$

- Using properties of logarithms, we often consider differences in log-likelihood:

$$\ell_{\text{full}} - \ell_{\text{reduced}} \sim \frac{1}{2}\chi^2_d,$$

  where d is the difference in number parameters between the models.
- The test can be extended for restricted likelihood ratio tests (RLRT); the idea is similar: fit two models (full vs reduced) using REML, compare differences in the restricted log-likelihoods.
- Rather than fitting two models manually and comparing, we will use the `lmerTest` package to do this automatically for us.

```
# install.packages("lmerTest")
lmerTest::ranova(output)

ANOVA-like table for random-effects: Single term deletions

Model:
diameter ~ 1 + (1 | plate)
            npar  logLik    AIC     LRT Df Pr(>Chisq)
<none>         3 -306.63 619.26
(1 | plate)    2 -306.71 617.43 0.17402  1     0.6766
```

Acknowledgments

- Compiled on October 22, 2026 using R version 4.5.2.

- Licensed under the Creative Commons Attribution-NonCommercial license.  Please share and remix non-commercially, mentioning its origin.

- We acknowledge students and instructors for previous versions of this course / slides.

References

Oehlert GW (2000). *A First Course in Design and Analysis of Experiments*. W. H. Freeman. ISBN 0-7167-3510-5. 1, 4